arXiv:hep-th/0211232v2 30 Dec 2002

MIFP-02-09
OU-HET 418
hep-th/0211232
November 2002

Open Wilson Lines as States of Closed String

Koichi MURAKAMI*

George P. and Cynthia W. Mitchell Institute for Fundamental Physics,
Texas A&M University, College Station, TX 77843-4242, USA

Toshio NAKATSU†

Department of Physics, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560-0043, Japan

Abstract

System of a D-brane in bosonic string theory on a constant B field background is studied in order to obtain further insight into the bulk-boundary duality. Boundary states which describe arbitrary numbers of open-string tachyons and gluons are given. UV behaviors of field theories on the non-commutative world-volume are investigated by using these states. We take zero-slope limits of generating functions of one-loop amplitudes of gluons (and open-string tachyons) in which the region of the small open-string proper time is magnified. Existence of B field allows the limits to be slightly different from the standard field theory limits of closed-string. They enable us to capture world-volume theories at a trans-string scale. In this limit the generating functions are shown to be factorized by two curved open Wilson lines (and their analogues) and become integrals on the space of paths with a Gaussian distribution around straight lines. These indicate a possibility that field theories on the non-commutative world-volume are topological at such a trans-string scale. We also give a proof of the Dhar-Kitazawa conjecture by making an explicit correspondence between the closed-string states and the paths. Momentum eigenstates of closed-string or momentum loops also play an important role in these analyses.

---

[*] E-mail: kmurakami@physics.tamu.edu
[†] E-mail: nakatsu@het.phys.sci.osaka-u.ac.jp

1 Introduction

Systems of interacting open- and closed-strings play important roles in several aspects of string theory. It has been found out that theories even formulated as pure closed-strings have open-string sectors when they are accompanied by D-branes [1]. One of the most important features of such open-closed mixed systems is a duality between open- and closed-strings. This duality becomes manifest, for instance, by seeing one-loop diagrams of open-string as tree propagations of closed-string through modular transformations on the string world-sheets. In the systems of D-branes, this duality should be a rationale for correspondence between gauge theory on the world-volumes (open-string sector) and gravity theory in the bulk space-time (closed-string sector). AdS/CFT correspondence [2][3][4] may be regarded as one of the most remarkable examples of such bulk-boundary correspondence. This viewpoint was emphasized also in [5].

Nevertheless it becomes very difficult to establish the correspondence between the two. This is essentially because light particles (IR effects) in the one sector are realized by summing up whole massive towers (UV effects) in the other sector. When a constant B field background of closed-string is turned on, however, we expect that the situation could be drastically changed. On this background, the world-volumes of D-branes become non-commutative [6][7][8][9][10]. Low-energy effective theories of the open-string sector, therefore, become field theories on the non-commutative world-volumes. It was pointed out [11] that in such theories there happens a mixing of the UV and the IR. This UV/IR mixing gives [11] us a chance to capture some effects of light particles in the bulk gravity theories, e.g. gravitons, by investigating non-commutative field theories on the world-volumes. To pursue such possibility, non-planar one-loop amplitudes of open-string were studied on this background [12][13][14][15][16]. Higher loops were investigated as well in [17].

Coupling of non-commutative D-branes to closed-string in the bulk has attracted much interest particularly from the viewpoint of bulk-boundary correspondence [18][19]. It was pointed out in [20][21] that the generalized star-products arise in disk amplitudes consisting of a closed-string vertex operator and open-string vertex ones on a constant B field background. The generalized star products are also found to appear in straight open Wilson lines by the expansions in powers of non-commutative gauge fields [22]. These observations were combined in [23] and it was shown there that in the zero-slope limit of [10] disk amplitudes of a closed-string tachyon and arbitrary numbers of gauge fields on this background give rise to a straight open Wilson line. Thus it has been clarified that open Wilson lines play a role in the correspondence between the bulk gravity and the non-commutative gauge theories.

Open Wilson lines found in [24] are remarkable gauge invariant objects in non-commutative gauge theories. Gauge invariant operators in non-commutative gauge theories can be constructed in the forms of local operators smeared along straight Wilson lines [24][25][26]. It was also shown in [23] that coupling of closed-string graviton to non-commutative gauge theories actually is an operator of this form.

Open Wilson lines need not be straight in order to be gauge invariant. Taking account of the above role played by straight open Wilson lines in the bulk-boundary duality, Dhar and Kitazawa [27] conjectured that curved open Wilson lines fluctuating around a straight path should be brought about by higher level (stringy) states of closed-string. They showed that the coupling of graviton state is in fact the gauge invariant operator which appears in the leading coefficients in harmonic expansions of curved open Wilson lines around a straight path.

It is well-known that the so-called boundary states [28][29][30] provide a description of D-branes in closed-string theory. Open- and closed-strings interact on the branes in space-time. Therefore it is further expected that boundary states admit to describe these interactions. In this article, receiving the above development and understanding, we study the system of a D-brane in bosonic string theory on a constant B field background. In order to obtain further insight into the bulk-boundary duality, we first exploit the boundary state formalism to include states which describe arbitrary numbers of open-string tachyons and gluons. Open-string legs of these boundary states, that is, tachyons and gluons, need not to be on-shell. Thereby it becomes possible to study the duality.

On a constant B field background, boundary state without open-string legs has been constructed in [28][29][30]. It is briefly reviewed in the next section. Let us denote the state by $|B\rangle$. This state does not lead any couplings of a closed-string to open-string excitations on the world-volume. We may regard $|B\rangle$ as a perturbative vacuum of closed-string in the presence of the brane. It is also possible to interpret the state as a Bogolubov transform of the $SL_2(\mathbb{C})$-invariant vacuum of closed-string : $|B\rangle = g|\mathbf{0}\rangle$, where g is a suitable generator of the transformation.

In Section 3 we construct boundary states of open-string tachyons. Basic idea of the construction may be explained as follows. We first examine an insertion of a closed-string tachyon into the brane. Let us suppose that the closed-string world-sheet is an infinite semi-cylinder (σ, τ) with $\tau \geq 0$ and that the brane or the state $|B\rangle$ is located at $\tau = 0$. We may describe the insertion by $\lim_{\tau \to 0+} V_T(\sigma, \tau)|B\rangle$, where V_T is a closed-string tachyon vertex operator. However, there occurs a singularity within V_T at the world-sheet boundary. This originates in correlations between the chiral and the anti-chiral sectors which are caused by the brane. We then regularize such a singularity and define a renormalized tachyon vertex operator V_T^{ren}. It turns out surprisingly that the renormalized operator becomes an open-string tachyon vertex at the boundary. In our prescription of the renormalization, disk Green's function and the transform $g^{-1}V_T g$ play essential roles. Boundary state of a single open-string tachyon is given by $\lim_{\tau\to 0+} V_T^{ren}(\sigma,\tau)|B\rangle$. Boundary states of arbitrary numbers of open-string tachyons turn out to be realized by the successive insertions of the above renormalized operators. As for boundary states of gluons, they can be obtained by taking the same steps as illustrated above, except that we need to introduce a suitable local operator of closed-string in place of V_T and renormalize it in a suitable fashion. These are presented in Section 7. Consistency of the constructions is examined from several aspects at some length. In particular, we compute closed-string tree propagations between these boundary states and make sure that they reproduce the corresponding open-string one-loop amplitudes. These are presented separately in Section 3 for the tachyons and in Section 7 for the gluons. It is also worth noting that the present constructions are relevant on a vanishing B field background.

Low-energy description of world-volume theories of D-branes is obtained by taking a zero-slope limit $\alpha' \to 0$ so that it makes all perturbative stringy states (mass$^2 \sim 1/\alpha'$) of open-string infinitely heavy and decouple from the light states. As regards one-loop amplitudes of open-string, this can be achieved by taking the limit with $\alpha'\tau^{(o)}$ fixed. Here $\tau^{(o)}$ is the proper time on the world-sheet of open-string. The fixed parameter $s^{(o)} \equiv \alpha'\tau^{(o)}$ becomes the Schwinger parameter of one-loop amplitudes of low-energy effective world-volume theories. It is shown [12][13][14][15][16][32] that one-loop amplitudes of open-string tachyons on a constant B field background reduce to those of a non-commutative (tachyonic) scalar field theory on the world-volume. The above zero-slope limit can be interpreted as magnification of the string amplitudes in the vicinity of $\tau^{(o)} = +\infty$. In Section 4 we investigate a possible UV behavior of this non-commutative field theory. In order to know the UV behavior one needs to focus on the region $s^{(o)} \approx 0$. For the description we take the following route. We introduce a parameter $s^{(c)} \equiv \alpha'|\tau^{(c)}|$, where $\tau^{(c)}$ is the closed-string proper time. $s^{(o)}$ and $s^{(c)}$ are related to each other through the modular transformation of the world-sheet by $s^{(c)} = 2\alpha'^2/s^{(o)}$. Then we take a zero-slope limit with $s^{(c)}$ fixed. Existence of B field allows us to make the limit slightly different from the standard field theory limit of closed-string (gravity limit). This enables us to capture the world-volume theories. This zero-slope limit is magnification of the string amplitudes in the vicinity of $\tau^{(o)} = 0$ and hence the region $s^{(o)} \approx 0$. Strictly speaking, this is a trans-string scale of the world-volume theories. Generating function of one-loop amplitudes of open-string tachyons is found out to be factorized at the limit into two (analogues of) straight open Wilson lines exchanging closed-string tachyons. In general, one-loop amplitudes of open-string are factorized by a tower of closed-string states. In gravity limit the propagations of closed-string tachyons become dominant. Although the present limit is slightly different from gravity limit, these zero-slope limits share the common property. UV behavior of the non-commutative gauge theory is explored in Section 8 by following the same route. Generating function of one-loop amplitudes of gluons is also found out to be factorized at this limit into two straight open Wilson lines exchanging closed-string tachyons. These analyses indicate a possibility that field theories on the non-commutative world-volume become topological at such a trans-string scale of the world-volume.

The conjecture [27] leads us to expect that curved open Wilson lines somehow factorize generating functions of one-loop amplitudes of gluons or open-string tachyons. The standard factorization made by particle states of closed-string seems to be little use. Instead, momentum eigenstates of closed-string are used for the factorization. Closed-string momentum $P(\sigma)$ is a momentum loop (a loop in the momentum space) while closed-string coordinate $X(\sigma)$ is a space-time loop. Section 5 is devoted to introduction of coordinate and momentum eigenstates of closed-string. We also provide some observations on their relations with boundary states. Factorizations made by the momentum eigenstates are examined in Section 6 for the tachyons and in Section 9 for the gluons. It turns out that in the zero-slope limit (the previous UV limit of the world-volume theories) momentum loops become curves in the non-commutative world-volume and that open Wilson lines along these curves appear in the factorizations. The generating functions are factorized by two curved open Wilson lines. More precisely, they become integrals on the space of curves with a Gaussian distribution around straight lines. As regards fluctuations from the straight lines, width of the distribution becomes so sharp that the integrals reduce to the previous factorizations by straight open Wilson lines.

We start Section 9 by giving a proof of the conjecture made by Dhar and Kitazawa. We introduce the closed-string state $|\Omega(P)\rangle =: \exp\left(i\int_0^{2\pi} d\sigma P_\mu(\sigma)\hat{X}^\mu(\sigma)\right) : |\mathbf{0}\rangle$, where $P(\sigma)$ is a momentum loop. This state is not an eigenstate of closed-string momentum but serves as a generating function of (generally off-shell) closed-string states. Overlap between $|\Omega(P)\rangle$ and boundary states with open-string legs is a generating function of couplings of all the closed-string states to the non-commutative D-brane. The overlap with the boundary states of gluons is shown to become a curved open Wilson line in the zero-slope limit. We make an explicit correspondence between the closed-string states and the coefficients of harmonic expansions of the curve. These provide the proof. We also show that in the zero-slope limit the momentum eigenstate is identified with the state $|\Omega(P)\rangle$ after some manipulation. This accounts for the previous factorizations by curved open Wilson lines.

Wilson line is invariant under reparametrizations of the path. The corresponding transformations on the string world-sheet are reparametrizations of the boundary. The reparametrization invariance of boundary states is fulfilled by imposing the Ishibashi condition [31] or equivalently the BRST invariance on these states. As regards the boundary states of open-string tachyon and gluon it is shown in Sections 3 and 7 that their reparametrization invariance is equivalent to the on-shell conditions. On the other hand we do not require any condition on gluons to obtain open Wilson lines. This puzzle is solved by seeing that the Ishibashi condition or the BRST invariance becomes null in the present zero-slope limit.

It is observed in Section 10 that all the boundary states constructed so far in this article are eigenstates of the closed-string momentum operators. The eigenvalues are essentially delta functions on the world-sheet boundary. Their boundary actions are computed by following the prescription given in [30]. It turns out that they are the standard boundary actions used in the path-integral formalism of the world-sheet theory. After a speculation based on these observations we finally make a conjecture on the duality between open- and closed- strings.

In Appendix A world-volume and space-time tensors used in the text are summarized including their relations. In Appendix B some formulae of creation and annihilation modes are described. These are necessary for our computations of several string amplitudes in the text. Oscillator realizations of coordinate and momentum eigenstates are presented in Appendix C.

2 A Short Course on Boundary States

Let us consider the system of Dp-brane in bosonic string theory. We take the following closed-string background :

$$\begin{aligned} ds^2 &= g_{MN}dx^M dx^N = g_{\mu\nu}dx^\mu dx^\nu + g_{ij}dx^i dx^j \ , \\ B &= \frac{1}{2}B_{\mu\nu}dx^\mu \wedge dx^\nu \ . \end{aligned} \tag{2.1}$$

Here g_{MN} is a flat space-time metric, which we refer to as closed-string metric. Two-form gauge field $B_{\mu\nu}$ is constant. We divide the space-time directions into two pieces, $x^M = (x^\mu, x^i)$, where $\mu = 0, 1, \cdots, p$ and $i = p+1, \cdots, D-1$. The directions x^μ are supposed to be parallel to the Dp-brane. The directions x^i are perpendicular to the brane.

Closed-string may capture the brane. Relevant action of a closed-string takes the form :

$$S[X] = \frac{1}{4\pi\alpha'}\int_\Sigma d\tau d\sigma \Big\{\partial_a X^M \partial^a X^N g_{MN} - i2\pi\alpha'\epsilon^{ab}\partial_a X^\mu \partial_b X^\nu B_{\mu\nu}\Big\} \ , \tag{2.2}$$

where ϵ^{ab} is the antisymmetric tensor on the world-sheet with $\epsilon^{\tau\sigma} \equiv 1$. The world-sheet Σ is a disk or an infinite semi-cylinder, and we use the cylinder coordinates (τ, σ) ($\tau \geq 0$ and $0 \leq \sigma < 2\pi$). Closed-string interacts with the brane at the boundary of the world-sheet, that is, at $\tau = 0$. The second term of the action is an integration of the two-form (its pull-back) on Σ. Since it is an exact two-form, by applying the Stokes theorem we can recast it into a boundary integral :

$$S[X] = \frac{1}{4\pi\alpha'} \int_\Sigma d\tau d\sigma \partial_a X^M \partial^a X^N g_{MN} - \frac{i}{2} \int_{\partial\Sigma} d\sigma B_{\mu\nu} X^\mu \partial_\sigma X^\nu \ . \tag{2.3}$$

Energy-momentum tensor $T_{ab}(\sigma, \tau)$ is obtained from the action. Since the integration of the two-form is independent of the world-sheet metric, T_{ab} acquires the standard form without the B-field.

Closed-string coordinates $X^M(\sigma, \tau)$ have mode expansions of the form,

$$X^M(\sigma, \tau) = \hat{x}_0^M - i\alpha' \hat{p}_0^M \tau + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \left(\frac{\alpha_n^M}{n} e^{-n(\tau + i\sigma)} + \frac{\tilde{\alpha}_n^M}{n} e^{-n(\tau - i\sigma)} \right) \ , \tag{2.4}$$

and the standard first quantization requires the following commutation relations :

$$\begin{aligned} &[\hat{x}_0^M, \hat{p}_0^N] = ig^{MN} \ , \quad [\hat{x}_0^M, \hat{x}_0^N] = [\hat{p}_0^M, \hat{p}_0^N] = 0 \ , \\ &[\alpha_m^M, \alpha_n^N] = mg^{MN}\delta_{m+n} \ , \quad [\tilde{\alpha}_m^M, \tilde{\alpha}_n^N] = mg^{MN}\delta_{m+n} \ , \quad [\alpha_m^M, \tilde{\alpha}_n^N] = 0 \ . \end{aligned} \tag{2.5}$$

The energy-momentum tensor generates reparametrizations of the world-sheet. Their infinitesimal forms turn out to be the Virasoro algebras with the central charges equal to D. The Virasoro generators are given by the expansions, $T_{zz}(z) = \sum_{m \in \mathbb{Z}} L_m z^{-m-2}$ and $T_{\overline{z}\overline{z}}(\overline{z}) = \sum_{m \in \mathbb{Z}} \tilde{L}_m \overline{z}^{-m-2}$. Here we use the complex coordinates $(z, \bar{z})$ instead of the cylinder coordinates. They are related by $(z, \bar{z}) = (e^{\tau + i\sigma}, e^{\tau - i\sigma})$. L_m and $\tilde{L}_m$ are generators of the chiral and the anti-chiral sectors respectively. These have the following representations in terms of the oscillator modes :

$$L_m = \frac{1}{2} : \sum_{n \in \mathbb{Z}} g_{MN} \alpha_n^M \alpha_{m-n}^N : \ , \quad \tilde{L}_m = \frac{1}{2} : \sum_{n \in \mathbb{Z}} g_{MN} \tilde{\alpha}_n^M \tilde{\alpha}_{m-n}^N : \ , \tag{2.6}$$

where $\alpha_0^M = \tilde{\alpha}_0^M = \sqrt{\frac{\alpha'}{2}} p_0^M$. $: \ :$ denotes the standard normal ordering with respect to the $SL_2(\mathbb{C})$-invariant vacuum of closed-string $|\mathbf{0}\rangle$:

$$: \hat{x}_0^M \hat{p}_0^N : \ = \ : \hat{p}_0^N \hat{x}_0^M : \ = \ \hat{x}_0^M \hat{p}_0^N \ , \qquad : \alpha_{-n}^M \alpha_n^N : \ = \ : \alpha_n^N \alpha_{-n}^M : \ = \ \alpha_{-n}^M \alpha_n^N \ , \tag{2.7}$$

for $n \geq 1$ and the similar prescription for the anti-chiral modes.

Closed-string can find the brane through the boundary conditions imposed at $\tau = 0$. We take a variation of the action :

$$\begin{aligned}\delta S[X] &= -\frac{1}{2\pi\alpha'}\int_{\Sigma} d\tau d\sigma \delta X^M g_{MN}\Big(\partial_\tau\partial_\tau + \partial_\sigma\partial_\sigma\Big)X^N \\ &\quad +\frac{1}{2\pi\alpha'}\int_{\partial\Sigma} d\sigma \left[\delta X^\mu\Big(g_{\mu\nu}\partial_\tau X^\nu - 2\pi i\alpha' B_{\mu\nu}\partial_\sigma X^\nu\Big) + \delta X^i g_{ij}\partial_\tau X^j\right] \ . \end{aligned} \tag{2.8}$$

Vanishing of the first term (bulk term) for an arbitrary $\delta X^M(\sigma,\tau)$ gives the equation of motions of the string-coordinates. Vanishing of the second term (boundary term) needs to be considered separately for $\delta X^\mu(\sigma,0)$ and $\delta X^i(\sigma,0)$. The interaction with the brane is possible everywhere on its world-volume. Thus $\delta X^\mu(\sigma,\tau)$ are arbitrary at the boundary. We need to impose the Neumann boundary condition on X^μ [1]. On the other hand $\delta X^i(\sigma,\tau)$ need to vanish at the boundary. For the vanishing we impose the Dirichlet boundary condition on X^i :

$$\Big[g_{\mu\nu}\partial_\tau X^\nu - 2\pi i\alpha' B_{\mu\nu}\partial_\sigma X^\nu\Big](\sigma,0) = 0 \ , \qquad X^i(\sigma,0) = x_0^i \ . \tag{2.9}$$

Here x_0^i is a position of the brane in the space-time. Owing to these boundary conditions we refer the directions x^μ and x^i respectively as the Neumann and the Dirichlet directions.

The above conditions are used to introduce a boundary state in the first quantized picture of closed-string. Let us denote it by $|B\rangle$. It is a state which satisfies the following conditions :

$$\Big[g_{\mu\nu}\partial_\tau X^\nu - 2\pi i\alpha B_{\mu\nu}\partial_\sigma X^\nu\Big](\sigma,0)|B\rangle = 0 \ , \qquad X^i(\sigma,0)|B\rangle = x_0^i|B\rangle \ . \tag{2.10}$$

These are linear constraints on $|B\rangle$ and determine the state modulo its normalization. It turns out that the state satisfies the Ishibashi condition [31] as follows :

$$\Big(L_n - \tilde{L}_{-n}\Big)\,|B\rangle = 0 \qquad \text{for } \forall n \ . \tag{2.11}$$

The above state can be interpreted as a perturbative vacuum of closed-string in the presence of the brane. Since there are no correlations between string-coordinates of the Neumann and the Dirichlet directions, we can factorize the state into a product :

$$|B\rangle = |B_N\rangle \otimes |B_D\rangle, \tag{2.12}$$

where the subscripts N and D denote the corresponding boundary conditions. We may refer to these two states respectively as the Neumann and the Dirichlet boundary states for short. They satisfy the following constraints :

$$\Big[g_{\mu\nu}\partial_\tau X^\nu - 2\pi i\alpha' B_{\mu\nu}\partial_\sigma X^\nu\Big](\sigma,0)|B_N\rangle = 0 \ , \quad X^i(\sigma,0)|B_D\rangle = x_0^i|B_D\rangle \ . \tag{2.13}$$

---

[1] Precisely speaking, this is a mixed boundary condition. We refer to this boundary condition as the Neumann boundary condition because it reduces to the Neumann boundary condition in the absence of B field. In this paper we use this terminology otherwise stated.

These constraints may be handled in the oscillator representations. For the Neumann boundary state it can be read as follows :

$$\hat{p}_{0\mu}|B_N\rangle = 0 \ , \quad \Big(E_{\mu\nu}\alpha_n^\nu + E_{\mu\nu}^T\tilde{\alpha}_{-n}^\nu\Big)|B_N\rangle = 0 \qquad \text{for } \forall n \neq 0, \tag{2.14}$$

where $E_{\mu\nu}$ and its transpose $E_{\mu\nu}^T$ are tensors defined as

$$E_{\mu\nu} = g_{\mu\nu} + 2\pi\alpha' B_{\mu\nu} \ , \quad E_{\mu\nu}^T = g_{\mu\nu} - 2\pi\alpha' B_{\mu\nu} \ . \tag{2.15}$$

As regards the Dirichlet boundary state the constraint can be read as follows :

$$\hat{x}_0^i|B_D\rangle = x_0^i|B_D\rangle \ , \quad \left(\alpha_n^i - \tilde{\alpha}_{-n}^i\right)|B_D\rangle = 0 \qquad \text{for } \ \forall n \neq 0 \ . \tag{2.16}$$

Boundary state $|B_D\rangle$

Let us describe these boundary states including their normalizations. The above constraints can be solved without any difficulty in the oscillator representations. Determinations of their normalization factors need to be cared. We start with the Dirichlet boundary state.

The Dirichlet boundary state turns out to be given by

$$|B_D\rangle \equiv \left(\frac{(2\pi^2\alpha')^d}{\det g_{ij}}\right)^{\frac{1}{4}} \prod_{n=1}^{\infty} \exp\left\{\frac{1}{n} g_{ij}\alpha_{-n}^i\tilde{\alpha}_{-n}^j\right\}|x_{0D}\rangle, \tag{2.17}$$

where we put $d = D - p - 1$. The state $|x_{0D}\rangle = |x_{0D} = (x_0^i)\rangle$ is an eigenstate of the zero modes $\hat{x}_0^i$ with eigenvalues x_0^i. Momentum representation of $|x_{0D}\rangle$ is given by

$$|x_{0D}\rangle = \frac{1}{(2\pi)^{d/2}}\int d^dk\,|k_D\rangle e^{-ik_ix_0^i} \ , \tag{2.18}$$

where $|k_D\rangle = |k_D = (k_i)\rangle$ denotes an eigenstate of the momentum zero modes $\hat{p}_{0i}$ with eigenvalues k_i defined as

$$|k_D\rangle = e^{ik_i\hat{x}_0^i}|\mathbf{0}\rangle \ . \tag{2.19}$$

The dual state can be obtained by taking the BPZ conjugation :

$$\langle B_D| = \left(\frac{(2\pi^2\alpha')^d}{\det g_{ij}}\right)^{\frac{1}{4}} \langle x_{0D}| \prod_{n=1}^{\infty} \exp\left\{\frac{1}{n} g_{ij}\alpha_{-n}^i\tilde{\alpha}_{-n}^j\right\} . \tag{2.20}$$

It is easy to see that the state (2.17) satisfies the conditions (2.16). Since they are linear conditions, normalization factor of the state should be determined by other means. We fixed



Figure 1: Evolution of closed-string by an imaginary time $\pi|\tau^{(c)}|$.

it to be $\left(\frac{(2\pi^2\alpha')^d}{\det g_{ij}}\right)^{\frac{1}{4}}$ in the above. It is determined by claiming that closed-string propagations along the Dirichlet directions reproduce the vacuum one-loop amplitude of open-string which satisfies the Dirichlet-Dirichlet (D-D) boundary conditions.

We parametrize propagations of closed-string by $\tau^{(c)} \in i\mathbb{R}_{\geq 0}$. Closed-string evolves by an imaginary time $\pi|\tau^{(c)}|$ with the Hamiltonian $L_0 + \tilde{L_0}$. See Figure 1. Closed-string propagations along the Dirichlet directions are measured by $\langle B_D|q_c^{\frac{L_0+\tilde{L}_0}{2}}|B_D\rangle$, where we put $q_c = e^{2\pi i\tau^{(c)}}$. This amplitude can be easily calculated by using the formula in Appendix B. It turns out to be

$$\begin{aligned}\langle B_D|q_c^{\frac{L_0+\tilde{L}_0}{2}}|B_D\rangle &= \left(\frac{\alpha'}{2}\right)^{\frac{d}{2}}\int\frac{dk}{\sqrt{\det g_{ij}}}q_c^{\frac{\alpha'}{4}k_i g^{ij}k_j}\prod_{n=1}^{\infty}(1-q_c^n)^{-d},\\ &= e^{\frac{\pi i\tau^{(c)}d}{12}}\left(-i\tau^{(c)}\right)^{-\frac{d}{2}}\eta\left(\tau^{(c)}\right)^{-d},\end{aligned} \qquad (2.21)$$

where $\eta(\tau) \equiv q^{1/24}\prod_{n=1}^{\infty}(1-q^n)$ with $q = e^{2\pi i\tau}$. Contribution of the world-sheet reparametrization ghosts is excluded in Eq.(2.21). It will be included in string propagations along the Neumann directions. The corresponding amplitude of open-string is given by $\mathrm{Tr}_{D-D}e^{-\tau^{(o)}L_0}$. Open-string propagates by an imaginary time $\tau^{(o)}$ ($\in \mathbb{R}_{\geq 0}$) with the Hamiltonian L_0. See Figure 2. The trace is taken over the sector of open-string which satisfies the D-D boundary conditions. The amplitude becomes as follows :

$$\mathrm{Tr}_{D-D}e^{-\tau^{(o)}L_0} = e^{-\frac{\tau^{(o)}d}{24}}\eta\left(-\frac{\tau^{(o)}}{2\pi i}\right)^{-d}. \qquad (2.22)$$



Figure 2: Evolution of open-string by an imaginary time $\tau^{(o)}$. Two bold horizontal lines are identified.

The standard argument allows us to interpret these open-string amplitudes as the free propagations of closed-string. Two imaginary times are related with each other by $\tau^{(o)} = 2\pi i/\tau^{(c)}$. This leads us to write Eq.(2.22) as

$$\mathrm{Tr}_{D-D} e^{-\tau^{(o)} L_0} = e^{-\frac{\pi i d}{12\tau^{(c)}}} \left(-i\tau^{(c)}\right)^{-\frac{d}{2}} \eta\left(\tau^{(c)}\right)^{-d} \Bigg|_{\tau^{(c)}=\frac{2\pi i}{\tau^{(o)}}} , \tag{2.23}$$

where the modular transformation $\eta\left(-\frac{1}{\tau^{(c)}}\right)^3 = (-i\tau^{(c)})^{3/2}\eta\left(\tau^{(c)}\right)^3$ is used. A comparison between Eqs.(2.21) and (2.23) gives the identity :

$$\langle B_D | q_c^{\frac{L_0+\tilde{L}_0}{2}} | B_D \rangle = e^{\frac{\pi i d}{12}\left(\tau^{(c)}+\frac{1}{\tau^{(c)}}\right)} \, \mathrm{Tr}_{D-D} e^{-\tau^{(o)} L_0} \Bigg|_{\tau^{(o)}=\frac{2\pi i}{\tau^{(c)}}} . \tag{2.24}$$

Therefore the propagations of closed-string reproduce correctly the corresponding one-loop amplitude of open-string except the factor $e^{\frac{\pi i d}{12}(\tau^{(c)}+1/\tau^{(c)})}$. This factor turns out to be canceled by a similar one appearing in closed-string propagations along the Neumann directions.

Boundary state $|B_N\rangle$

The Neumann boundary state is given [28] by

$$|B_N\rangle \equiv \left(\frac{\det^2 E_{\mu\nu}}{(2\alpha')^{p+1}(-\det g_{\mu\nu})}\right)^{\frac{1}{4}} \prod_{n=1}^{\infty} \exp\left\{-\frac{1}{n} N_{\mu\nu} \alpha^{\mu}_{-n} \tilde{\alpha}^{\nu}_{-n}\right\} |\mathbf{0}\rangle , \tag{2.25}$$

where $N_{\mu\nu}$ is a tensor defined by

$$N_{\mu\nu} = \left(g\frac{1}{E}E^T \right)_{\mu\nu} \ . \tag{2.26}$$

The dual state is obtained by the BPZ conjugation. Its explicit form is as follows :

$$\langle B_N| = \left(\frac{\det^2 E_{\mu\nu}}{(2\alpha')^{p+1}(-\det g_{\mu\nu})} \right)^{\frac{1}{4}} \langle \mathbf{0}| \prod_{n=1}^{\infty} \exp\left\{ -\frac{1}{n} N_{\mu\nu}\alpha_n^{\mu}\tilde{\alpha}_n^{\nu} \right\} \ . \tag{2.27}$$

The above normalization factor of the state is chosen to reproduce the related one-loop amplitudes of open-string. This will be shown in the next section.

3 Open-String Tachyons in Closed-String Theory

As can be seen in Eqs.(2.17) and (2.25) these boundary states are the Bogolubov transforms of perturbative vacua. Generators of the transformations are $\prod_{n=1}^{\infty} \exp\left\{ \frac{1}{n} g_{ij}\alpha_{-n}^{i}\tilde{\alpha}_{-n}^{j} \right\}$ for the Dirichlet boundary and $\prod_{n=1}^{\infty} \exp\left\{ -\frac{1}{n} N_{\mu\nu}\alpha_{-n}^{\mu}\tilde{\alpha}_{-n}^{\nu} \right\}$ for the Neumann boundary. We can expect that information on the boundary conditions are all encoded in these generators.

Let $\mathcal{O}(\sigma,\tau)$ be a local operator of closed-string. Action of this operator on a boundary state $g|\mathbf{0}\rangle$, where g is a generator, can be written as follows :

$$\mathcal{O}(\sigma,\tau)g|\mathbf{0}\rangle = g \times g^{-1}\mathcal{O}(\sigma,\tau)g|\mathbf{0}\rangle. \tag{3.1}$$

One may expect that $g^{-1}\mathcal{O}(\sigma,\tau)g$ describes the Bogolubov transform of this local operator and ask the physical implication particularly from the viewpoint of boundary conformal field theories. However, story is not so simple. In general, $g^{-1}\mathcal{O}(\sigma,\tau)g$ turns out to be singular. More precisely it becomes singular at $\tau = 0$, where the boundary state resides. This reflects the fact that the system under consideration is actually a system of closed- and open-strings. We wish to make an idea of the Bogolubov transformations of these local operators rigorous. For this purpose, we have to perform a regularization by which the above singularity becomes tractable. This leads us to define $\mathrm{ad}_{g^{-1}}\mathcal{O}(\sigma,\tau)$, which becomes regular at the world-sheet boundary. It is a local operator and interpreted as the adjoint transform of $\mathcal{O}(\sigma,\tau)$ by g^{-1}. We will find out their physical interpretation. In this section we concentrate on tachyon vertex operators. Since we are mainly interested in the world-volume theory of p-brane, we restrict ourselves to the Bogolubov transform associated with the Neumann boundary state. We denote the generator by g_N,

$$g_N \equiv \prod_{n=1}^{\infty} \exp\left\{ -\frac{1}{n} N_{\mu\nu}\alpha_{-n}^{\mu}\tilde{\alpha}_{-n}^{\nu} \right\} . \tag{3.2}$$

3.1 Bogolubov transformation and renormalization

Let $V_T(\sigma,\tau;k)$ be closed-string tachyon vertex operator of momentum k. Momentum k is supposed to have only components along the Neumann directions. An explicit form is given by

$$\begin{aligned}
V_T(\sigma,\tau;k) &= :e^{ik_\mu X^\mu(\sigma,\tau)}:, \\
&= \exp(ik_\mu \hat{x}_0^\mu) \times |z|^{\alpha' k_\mu \hat{p}_0^\mu} \times \prod_{n=1}^{\infty} \exp\left[\sqrt{\frac{\alpha'}{2}}\frac{1}{n}k_\mu(\alpha_{-n}^\mu z^n + \tilde{\alpha}_{-n}^\mu \bar{z}^n)\right] \\
&\quad \times \prod_{n=1}^{\infty} \exp\left[-\sqrt{\frac{\alpha'}{2}}\frac{1}{n}k_\mu(\alpha_n^\mu z^{-n} + \tilde{\alpha}_n^\mu \bar{z}^{-n})\right]. \qquad (3.3)
\end{aligned}$$

To discuss the Bogolubov transform it is convenient to write down the transforms of oscillator modes of the string coordinates. These can be read as :

$$g_N^{-1}\alpha_n^\mu g_N = \alpha_n^\mu - (g^{-1}N)^\mu{}_\nu \tilde{\alpha}_{-n}^\nu, \qquad g_N^{-1}\tilde{\alpha}_n^\mu g_N = \tilde{\alpha}_n^\mu - (g^{-1}N^T)^\mu{}_\nu \alpha_{-n}^\nu, \qquad (3.4)$$

for $n \geq 1$. The modes α_n^μ and $\tilde{\alpha}_n^\mu$ for $n \leq 0$ are kept intact. (We put $\alpha_0^\mu = \tilde{\alpha}_0^\mu \equiv \sqrt{\alpha'/2}\hat{p}_0^\mu$.) The above mixture of the creation- and annihilation-modes makes the transform $g_N^{-1}V_T(\sigma,\tau;k)g_N$ singular at $\tau = 0$ or equivalently, since we put $z = e^{\tau+i\sigma}$, at $|z| = 1$. It can be written in the following form for $|z| > 1$:

$$g_N^{-1}V_T(\sigma,\tau;k)g_N = \left(\frac{|z|^2}{|z|^2-1}\right)^{-\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu} \times \mathrm{ad}_{g_N^{-1}}V_T(\sigma,\tau;k). \qquad (3.5)$$

Here we introduce a local operator $\mathrm{ad}_{g_N^{-1}}V_T$ which we interpret as the Bogolubov transform of V_T. It takes the form of

$$\mathrm{ad}_{g_N^{-1}}V_T(\sigma,\tau;k) = \mathcal{N}(\bar{z};k)\tilde{\mathcal{N}}(z;k)V_T(\sigma,\tau;k), \qquad (3.6)$$

where $\mathcal{N}(\bar{z};k)$ and $\tilde{\mathcal{N}}(z;k)$ are operators consisting only of the creation modes :

$$\begin{aligned}
\mathcal{N}(\bar{z};k) &= \prod_{n=1}^{\infty} \exp\left\{\sqrt{\frac{\alpha'}{2}}\frac{1}{n}k_\mu(g^{-1}N^T)^\mu{}_\nu \alpha_{-n}^\nu \bar{z}^{-n}\right\}, \\
\tilde{\mathcal{N}}(z;k) &= \prod_{n=1}^{\infty} \exp\left\{\sqrt{\frac{\alpha'}{2}}\frac{1}{n}k_\mu(g^{-1}N)^\mu{}_\nu \tilde{\alpha}_{-n}^\nu z^{-n}\right\}. \qquad (3.7)
\end{aligned}$$

Singularity of $g_N^{-1}V_T(\sigma,\tau;k)g_N$ comes from the factor $\left(\frac{|z|^2}{|z|^2-1}\right)^{-\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu}$ in Eq.(3.5). Because of this factor, the transform (3.5) becomes singular at $\tau = 0$ where the boundary state $|B_N\rangle$ is located. As will be seen soon, this factor should be subtracted in our construction of open-string tachyon vertex operator from a closed-string local operator. This factor can therefore be regarded as a renormalization factor of open-string tachyon vertex operator under its interpretation in terms of closed-string. Putting this factor aside for a while, let us examine the operators $\mathrm{ad}_{g_N^{-1}} V_T$. We first discuss their operator product expansion (OPE). It is convenient to recall the OPE between closed-string tachyon vertex operators. It can be read from the expansion (3.3) as follows :

$$V_T(\sigma_1,\tau_1;k^{(1)})V_T(\sigma_2,\tau_2;k^{(2)}) = |z_1-z_2|^{\alpha' k^{(1)}_\mu g^{\mu\nu} k^{(2)}_\nu} \times :V_T(\sigma_1,\tau_1;k^{(1)})V_T(\sigma_2,\tau_2;k^{(2)}): \ , \tag{3.8}$$

for $|z_1| \geq |z_2|$ and $z_1 \neq z_2$. The OPE under consideration changes from Eq.(3.8). The modification comes from operator products between $\mathcal{N}$, $\tilde{\mathcal{N}}$ and V_T. It can be calculated by using the expansions (3.7). We finally obtain :

$$\begin{aligned}&\mathrm{ad}_{g_N^{-1}} V_T(\sigma_1,\tau_1;k^{(1)})\mathrm{ad}_{g_N^{-1}} V_T(\sigma_2,\tau_2;k^{(2)}) \\ &\quad = \exp\Big\{-k^{(1)}_\mu \mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)k^{(2)}_\nu\Big\}\times :\mathrm{ad}_{g_N^{-1}} V_T(\sigma_1,\tau_1;k^{(1)})\mathrm{ad}_{g_N^{-1}} V_T(\sigma_2,\tau_2;k^{(2)}): \ ,\end{aligned} \tag{3.9}$$

for $|z_1| \geq |z_2|, |z_1z_2| \geq 1$ and $z_1 \neq z_2$. Here $\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)$ is Green's function on the unit disk $|z| \geq 1$ in the presence of a constant B field. It is defined by

$$\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2) \equiv \frac{\langle \mathbf{0}|X^\mu(\sigma_1,\tau_1)X^\nu(\sigma_2,\tau_2)|B_N\rangle}{\langle \mathbf{0}|B_N\rangle} - \frac{\langle \mathbf{0}|\hat{x}^\mu_0\hat{x}^\nu_0|\mathbf{0}\rangle}{\langle \mathbf{0}|\mathbf{0}\rangle}. \tag{3.10}$$

The RHS can be evaluated by using the Bogolubov transforms (3.4) and written down explicitly as follows :

$$\begin{aligned}\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2) &= -\frac{\alpha'}{2}g^{\mu\nu}\ln|z_1-z_2|^2 - \frac{\alpha'}{2}(g^{-1}Ng^{-1})^{\mu\nu}\ln\left(1-\frac{1}{z_1\bar{z}_2}\right) \\ &\quad -\frac{\alpha'}{2}(g^{-1}N^Tg^{-1})^{\mu\nu}\ln\left(1-\frac{1}{\bar{z}_1 z_2}\right).\end{aligned} \tag{3.11}$$

Let us recall that a system only of closed-strings admits a holomorphic factorization. Particularly there is no correlation between chiral and anti-chiral pieces, $X^\mu(z)$ and $\tilde{X}^\mu(\bar{z})$, of string coordinates $X^\mu(\sigma,\tau) = X^\mu(z) + \tilde{X}^\mu(\bar{z})$. The factorized term of $\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)$ in Eq.(3.11) is a sum of the correlations $\langle X^\mu(z_1)X^\nu(z_2)\rangle$ and $\langle \tilde{X}^\mu(\bar{z}_1)\tilde{X}^\nu(\bar{z}_2)\rangle$. The second and third terms are not factorized and they are respectively the correlations $\langle X^\mu(z_1)\tilde{X}^\nu(\bar{z}_2)\rangle$ and $\langle \tilde{X}^\mu(\bar{z}_1)X^\nu(z_2)\rangle$. These correlations are characteristic of open-string theory. Green's function provides a nice description of the singular factor in Eq.(3.5). Using this terminology we can write it as follows:

$$\left(\frac{|z|^2}{|z|^2-1}\right)^{-\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu} = \frac{\langle k|\,V_T(\sigma,\tau;k)\,|B_N\rangle}{\langle \mathbf{0}\,|B_N\rangle} = \exp\Big\{ik_\mu ik_\nu\langle X^\mu(z)\tilde{X}^\nu(\bar{z})\rangle\Big\} \ , \tag{3.12}$$

where the first equality follows from Eq.(3.5) and $\langle k|\mathrm{ad}_{g_N^{-1}}V_T(\sigma,\tau;k)|\mathbf{0}\rangle/\left\langle \mathbf{0}\,|B_N\right\rangle = 1$. Taking account of this expression, the transform (3.5) indicates that boundary states give rise to extra correlations, i.e. correlations between the chiral and the anti-chiral sectors, by the amount of $\langle X^\mu(z_1)\tilde{X}^\nu(\bar{z}_2)\rangle$. In the presence of the world-sheet boundary, or equivalently in the open-closed mixed system, correlations between the chiral and anti-chiral sectors exist even in the closed string sector, and we need to take care of them. As pointed out in [29], this is a direct result of the fact that boundary states on the unit circle reflect a vertex operator at z making its mirror image at $1/\bar{z}$: $e^{ik_\mu X^\mu(z)}|B_N\rangle = e^{ik_\mu (g^{-1}N)^\mu{}_\nu \tilde{X}^\nu\left(\frac{1}{z}\right)}|B\rangle$ (see also [14]). This brings about a short distance singularity at the boundary $z = 1/\bar{z}$. Since we intend to construct open-string vertex operators in terms of closed-string, we need to carry out a renormalization to manage this type of singularity.

The OPE (3.9) itself strongly suggests an interpretation of $\mathrm{ad}_{g_N^{-1}}V_T$ from the open-string viewpoint. To pursue such a possibility we introduce a renormalized tachyon vertex operator V_T^{ren} by subtracting the above singular factor :

$$V_T^{ren}(\sigma,\tau;k) \equiv \left(\frac{|z|^2}{|z|^2-1}\right)^{\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu} \times V_T(\sigma,\tau;k). \tag{3.13}$$

In other words we have

$$g_N^{-1}V_T^{ren}(\sigma,\tau;k)g_N = \mathrm{ad}_{g_N^{-1}}V_T(\sigma,\tau;k). \tag{3.14}$$

We call $V_T^{ren}(\sigma,\tau;k)$ renormalized *open-string* tachyon vertex operator with momentum k.

We need to explain why we call it open-string tachyon vertex operator. Let us consider an action of diffS^1, the group of diffeomorphisms of a circle, on V_T^{ren}. We take $L_n - \tilde{L}_{-n}$ $(n \in \mathbb{Z})$ as the generators. It is convenient to recall that L_n act on V_T in the following manner :

$$\Big[L_n, V_T(\sigma,\tau;k)\Big] = \left\{z^{n+1}\partial_z + \frac{\alpha'}{4}k_\mu g^{\mu\nu}k_\nu(n+1)z^n\right\}V_T(\sigma,\tau;k). \tag{3.15}$$

As regards $\tilde{L}_n$ their actions are the same as above except replacing z by $\bar{z}$. The action of diffS^1 on V_T^{ren} becomes slightly different from a sum of the above Virasoro actions. The modification comes from an existence of the renormalization factor in Eq.(3.13). It can be read as follows :

$$\begin{aligned}
&\Big[L_n - \tilde{L}_{-n}, V_T^{ren}(\sigma,\tau;k)\Big] \\
&\quad = \left\{z^{n+1}\partial_z - \bar{z}^{-(n-1)}\partial_{\bar{z}} + \frac{\alpha'}{4}k_\mu g^{\mu\nu}k_\nu\left((n+1)z^n + (n-1)\bar{z}^{-n}\right)\right. \\
&\qquad \left. + \frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu\frac{z^n - \bar{z}^{-n}}{|z|^2-1}\right\}V_T^{ren}(\sigma,\tau;k).
\end{aligned} \tag{3.16}$$

An implication may be seen by taking the limit $\tau \rightarrow 0+$ in the equation. By using the cylinder coordinates instead of $(z, \bar{z})$ the limit turns out to be as follows :

$$\begin{aligned}&\left[L_n - \tilde{L}_{-n}, \lim_{\tau\rightarrow 0+} V_T^{ren}(\sigma, \tau; k)\right] \\ &\qquad = \left\{-ie^{in\sigma}\frac{d}{d\sigma} + n\alpha' k_\mu G^{\mu\nu} k_\nu e^{in\sigma}\right\} \lim_{\tau\rightarrow 0+} V_T^{ren}(\sigma, \tau; k), \end{aligned} \tag{3.17}$$

where we translate the closed-string background tensors $g_{\mu\nu}$ and $B_{\mu\nu}$ into the open-string background tensors $G_{\mu\nu}$ and $\theta^{\mu\nu}$. We may call these two kinds of background tensors respectively closed- and open-string tensors for short. They are related [10] with each other by

$$\left(E^{-1}\right)^{\mu\nu} = G^{\mu\nu} + \frac{1}{2\pi\alpha'}\theta^{\mu\nu} \ . \tag{3.18}$$

In an actual derivation of Eq.(3.17) we use the following identities :

$$\begin{aligned} g^{\mu\nu} &= G^{\mu\nu} - \frac{1}{(2\pi\alpha')^2}\theta^{\mu\rho}G_{\rho\rho'}\theta^{\rho'\nu}, \\ g^{\mu\rho}N_{\rho\rho'}g^{\rho'\nu} &= \left(G^{\mu\nu} + \frac{1}{(2\pi\alpha')^2}\theta^{\mu\rho}G_{\rho\rho'}\theta^{\rho'\nu}\right) + \frac{1}{\pi\alpha'}\theta^{\mu\nu}. \end{aligned} \tag{3.19}$$

These can be obtained from Eq.(3.18). Related formulas are summarized in Appendix A.

The action (3.17) is completely the same as the action of the Virasoro algebra on open-string tachyon vertex operator with the same momentum. In addition to the OPE (3.9) this is the reason why we identify V_T^{ren} with renormalized open-string tachyon vertex operators.

We can also make the above consideration in terms of boundary states. Let us consider the state $V_T^{ren}(\sigma, \tau; k)|B_N\rangle$. By using the relation (3.14) we can write it in a form, $g_N\ \mathrm{ad}_{g_N^{-1}} V_T(\sigma, \tau; k)|\mathbf{0}\rangle$. An explicit form of $\mathrm{ad}_{g_N^{-1}} V_T$ has been given in Eq.(3.6). Thereby we can express the state in the oscillator representation and then take the limit $\tau \rightarrow 0+$ without ambiguity. We call thus obtained state $|B_N; (\sigma, k)\rangle$. It turns out to have the following form :

$$\begin{aligned} |B_N; (\sigma, k)\rangle &\equiv \lim_{\tau\rightarrow 0+} V_T^{ren}(\sigma, \tau; k)|B_N\rangle \\ &= \left(\frac{\det^2 E_{\mu\nu}}{(2\alpha')^{(p+1)}(-\det g_{\mu\nu})}\right)^{1/4} g_N \times \lim_{\tau\rightarrow 0+} \mathrm{ad}_{g_N^{-1}} V_T^{ren}(\sigma, \tau; k)|\mathbf{0}\rangle \\ &= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{(p+1)}}\right)^{1/4} \prod_{n=1}^{\infty} \exp\left[\sqrt{2\alpha'}\frac{1}{n}k_\mu\left\{\left(\frac{1}{E^T}g\right)^\mu_{\ \nu}\alpha^\nu_{-n}e^{in\sigma} + \left(\frac{1}{E}g\right)^\mu_{\ \nu}\tilde{\alpha}^\nu_{-n}e^{-in\sigma}\right\}\right] g_N|k\rangle. \end{aligned} \tag{3.20}$$

Here we express the normalization factor of $|B_N\rangle$ in terms of the open-string tensors. This translation is done by using the identity, $G_{\mu\nu} = E^T_{\mu\rho}g^{\rho\rho'}E_{\rho'\nu}$, which also follows from Eq.(3.18).

Action of diffS^1 on this state can be obtained from Eq.(3.17). It can be read as

$$\left(L_n - \tilde{L}_{-n}\right)|B_N;(\sigma,k)\rangle = (-i)\left\{e^{in\sigma}\frac{d}{d\sigma} + in\alpha' k_\mu G^{\mu\nu}k_\nu e^{in\sigma}\right\}|B_N;(\sigma,k)\rangle\ , \tag{3.21}$$

for $n \in \mathbb{Z}$. The Ishibashi condition imposed on $|B_N;(\sigma,k)\rangle$, i.e. vanishing of the RHS of the above equation for an arbitrary n, (strictly speaking, modulo a total derivative with respect to σ), requires $\alpha' k_\mu G^{\mu\nu} k_\nu = 1$. It is the on-shell condition of open-string tachyon.

3.2 String field theory viewpoint

The action of diffS^1 on the boundary state has an interpretation in terms of string field theory.

Fundamental ingredient in string field theory is a BRST charge Q. It is a grassmann-odd operator obeying the usual relations, $Q^2 = 0$ and $T_{ab}(\sigma,\tau) = \{Q, b_{ab}(\sigma,\tau)\}$, where T_{ab} and b_{ab} are respectively total energy-momentum tensor and anti-ghost field of a world-sheet theory. In the case of bosonic closed-string field theory [33] world-sheet theory consists of closed-string coordinates X^I (a matter system) and the world-sheet reparametrization ghosts. The ghost system is described by (b,c) for the chiral part and $(\tilde{b},\tilde{c})$ for the anti-chiral part. Closed-string BRST charge Q_c is decomposed into $Q_c = q + \tilde{q}$. The chiral part q has the following form :

$$q = \sum_n c_n L^X_{-n} + c_0 L_0^{(b,c)} + \sum_{m,n\neq 0} n c_n c_m b_{-n-m}, \tag{3.22}$$

where $L_n^{X,(b,c)}$ are the Virasoro generators of the matter and the ghost systems. c_n and b_n are the Fourier modes of ghost and anti-ghost fields : $c(\sigma,\tau) = \sum_n c_n e^{-n(\tau+i\sigma)}$ and $b(\sigma,\tau) = \sum_n b_n e^{-n(\tau+i\sigma)}$. The anti-chiral part $\tilde{q}$ has the same form as the above except replacing the chiral quantities with the anti-chiral ones. These q and $\tilde{q}$ are nilpotent independently and satisfy the relations, $\{q, b_n\} = L_n$ and $\left\{\tilde{q}, \tilde{b}_n\right\} = \tilde{L}_n$. Here $L_n = L_n^X + L_n^{(b,c)}$ and $\tilde{L}_n = \tilde{L}_n^X + \tilde{L}_n^{(b,c)}$ are the Virasoro generators of the total system.

Let $\mathcal{H}_c^X$ and $\mathcal{H}_c^{ghost}$ be respectively the Hilbert spaces of matter and ghost systems. We put $\mathcal{H}^{aux} = \mathcal{H}_c^X \otimes \mathcal{H}_c^{ghost}$. Closed-string Hilbert space $\mathcal{H}_c$ consists of vectors ψ of $\mathcal{H}^{aux}$ which satisfy the conditions,

$$(b_0 - \tilde{b}_0)\psi = 0, \qquad (L_0 - \tilde{L}_0)\psi = 0. \tag{3.23}$$

Necessity of the two conditions is explained in [33] from the perspective of two-dimensional conformal field theories.

Now we want to interpret the state $|B_N;(\sigma,k)\rangle$ as a state of the closed-string Hilbert space. Since it is a boundary state of the Neumann directions we first extend it to a vector of $\mathcal{H}_c^X$

by tensoring the boundary state (2.17) of the Dirichlet directions. As for the ghost sector an appropriate state is known. It is given [29] by

$$|B\rangle_{ghost} = \prod_{n=1}^{\infty} \exp\left\{c_{-n}\tilde{b}_{-n} + \tilde{c}_{-n}b_{-n}\right\}(c_0 + \tilde{c}_0)c_1\tilde{c}_1|\mathbf{0}\rangle, \tag{3.24}$$

and satisfies the following boundary conditions :

$$(c_n + \tilde{c}_{-n})|B\rangle_{ghost} = 0, \qquad (b_n - \tilde{b}_{-n})|B\rangle_{ghost} = 0, \tag{3.25}$$

for $n \in \mathbb{Z}$. It can be seen from these conditions that $|B\rangle_{ghost}$ satisfies the Ishibashi condition. Further tensoring the ghost boundary state we obtain $(|B_N;(\sigma,k)\rangle \otimes |B_D\rangle) \otimes |B\rangle_{ghost}$ of $\mathcal{H}^{aux}$. We call it $|B;(\sigma,k)\rangle_{tot}$. This state satisfies the first condition in Eq.(3.23) because of Eq.(3.25). As for the second, the $n=0$ case of Eq.(3.21) gives $(L_0 - \tilde{L}_0)|B;(\sigma,k)\rangle_{tot} = (-i)\partial_\sigma|B;(\sigma,k)\rangle_{tot}$. This means that we need to integrate over σ to regard the state as a vector of $\mathcal{H}_c$.

Physical states of closed-string are identified with the BRST invariant states of $\mathcal{H}_c$. In order to describe how Q_c acts on the boundary state we note a formula related with the ghost boundary state : Let $|\mathcal{O}\rangle_X$ be a state of $\mathcal{H}_c^X$. Action of Q_c on $|\mathcal{O}\rangle_X \otimes |B\rangle_{ghost}$ is expressed as

$$Q_c|\mathcal{O}\rangle_X \otimes |B\rangle_{ghost} = \sum_n c_{-n}(L_n^X - \tilde{L}_{-n}^X)|\mathcal{O}\rangle_X \otimes |B\rangle_{ghost} \tag{3.26}$$

This can be derived by the standard calculation. The action on the boundary state can be read as follows :

$$\begin{aligned} Q_c &\left[\int_0^{2\pi} d\sigma |B;(\sigma,k)\rangle_{tot}\right] \\ &= \int_0^{2\pi} d\sigma \sum_n c_{-n}(L_n^X - \tilde{L}_{-n}^X)|B;(\sigma,k)\rangle_{tot} \\ &= (-i)\int_0^{2\pi} d\sigma \sum_n e^{in\sigma}\left(\frac{d}{d\sigma} + in\alpha' k_{mu}G^{\mu\nu}k_\nu\right)c_{-n}|B;(\sigma,k)\rangle_{tot}, \end{aligned} \tag{3.27}$$

where Eq.(3.21) is used in the second equality. The BRST invariance requires, as we stated previously, the on-shell condition of open-string tachyon.

The above action can be identified with the BRST transformation of open-string tachyon vertex operator. c_n in Eq.(3.27) correspond to the same modes of ghost field appearing in open-string field theory. Actually Eq.(3.27) is an example of the formula given in [34]. It states that the closed-string BRST charge Q_c acts on boundary states as the generator of the BRST transformation of open-string field. It is used there to show a macroscopic description of the cubic open-string field theory is given by a boundary open-string field theory.

3.3 Boundary states with M open-string tachyons

The previous construction can be generalized to those of boundary states relevant to the description of M open-string tachyons. Let $V_T^{ren}(\sigma_r,\tau_r;k^{(r)})$ $(1\le r\le M)$ be the renormalized open-string tachyon vertex operators. Also let $z_r\equiv e^{\tau_r+i\sigma_r}$ be distinct points on the infinite semi-cylinder satisfying the condition, $|z_1|\ge|z_2|\ge\cdots|z_M|\ge 1$. We start with the following state :

$$V_T^{ren}(\sigma_1,\tau_1;k^{(1)})\cdots V_T^{ren}(\sigma_M,\tau_M;k^{(M)})|B_N\rangle. \tag{3.28}$$

We translate the state into a form, $g_N\prod_r \mathrm{ad}_{g_N^{-1}}V_T^{ren}(\sigma_r,\tau_r;k^{(r)})\ |\mathbf{0}\rangle$ by the relation (3.14). We use the OPE (3.9) for a further evaluation. Finally we arrive at the following representation :

$$\begin{aligned}
&V_T^{ren}(\sigma_1,\tau_1;k^{(1)})V_T^{ren}(\sigma_2,\tau_2;k^{(2)})\cdots V_T^{ren}(\sigma_M,\tau_M;k^{(M)})|B_N\rangle\\
&=\left(\frac{\det^2E_{\mu\nu}}{(2\alpha')^{p+1}(-\det g_{\mu\nu})}\right)^{1/4}\prod_{r<s}^{M}\exp\Big\{-k_\mu^{(r)}\mathbb{G}^{\mu\nu}(z_r,\bar z_r|z_s,\bar z_s)k_\nu^{(s)}\Big\}\\
&\qquad\times g_N:\mathrm{ad}_{g_N^{-1}}V_T(\sigma_1,\tau_1;k^{(1)})\mathrm{ad}_{g_N^{-1}}V_T(\sigma_2,\tau_2;k^{(2)})\cdots\mathrm{ad}_{g_N^{-1}}V_T(\sigma_M,\tau_M;k^{(M)}):|\mathbf{0}\rangle\\
&=\left(\frac{\det^2E_{\mu\nu}}{(2\alpha')^{p+1}(-\det g_{\mu\nu})}\right)^{1/4}\prod_{r<s}^{M}\exp\Big\{-k_\mu^{(r)}\mathbb{G}^{\mu\nu}(z_r,\bar z_r|z_s,\bar z_s)k_\nu^{(s)}\Big\}\\
&\qquad\times\prod_{n=1}^{\infty}\exp\left[\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\sum_{r=1}^{M}k_\mu^{(r)}\left(z_r^n\delta^\mu_\nu+\bar z_r^{-n}(g^{-1}N^T)^\mu{}_\nu\right)\alpha^\nu_{-n}\right]\\
&\qquad\times\prod_{n=1}^{\infty}\exp\left[\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\sum_{r=1}^{M}k_\mu^{(r)}\left(\bar z_r^n\delta^\mu_\nu+z_r^{-n}(g^{-1}N)^\mu{}_\nu\right)\tilde\alpha^\nu_{-n}\right]g_N\left|\sum_{r=1}^{M}k^{(r)}\right\rangle\quad.
\end{aligned}\tag{3.29}$$

Boundary state relevant to the description of M off-shell open-string tachyons may be obtained from the above state by taking the same steps as for $M=1$. This state will be called $|B_N;(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\rangle$. It is defined by the limit $\forall\tau_r\to 0+$ of Eq.(3.29). In the course of taking the limit we do not meet any difficulty for the oscillator part. What we need to take care are the products $\prod_{r<s}^M\exp\left\{-k_\mu^{(r)}\mathbb{G}^{\mu\nu}(z_r,\bar z_r|z_s,\bar z_s)k_\nu^{(s)}\right\}$. Behavior of Green's function at the boundary turns out to be as follows :

$$\begin{aligned}
&\lim_{\tau_{r,s}\to 0+}\mathbb{G}^{\mu\nu}(z_r,\bar z_r|z_s,\bar z_s)\\
&\qquad=-2\alpha'G^{\mu\nu}\ln|e^{i\sigma_r}-e^{i\sigma_s}|+\frac{i}{2\pi}\theta^{\mu\nu}\left\{\sigma_r-\sigma_s-\pi\epsilon(\sigma_r-\sigma_s)\right\},
\end{aligned}\tag{3.30}$$

where $\epsilon(x)$ is the sign function defined as $\epsilon(x)=1$ for $x>0$ and $=-1$ for $x<0$. Taking account of this behavior of Green's function, the limit $\forall\tau_r\to 0+$ becomes as follows :

$$\big|B_N;(\sigma_1,k^{(1)}),(\sigma_2,k^{(2)}),\cdots,(\sigma_M,k^{(M)})\big\rangle$$

$$
\begin{aligned}
&\equiv \lim_{\forall\tau_r\to 0+} V_T^{ren}(\sigma_1,\tau_1;k^{(1)})V_T^{ren}(\sigma_2,\tau_1;k^{(2)})\cdots V_T^{ren}(\sigma_M,\tau_M;k^{(M)})|B_N\rangle \\
&= \left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{1/4} \prod_{r<s}^{M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)} \\
&\quad \times \prod_{r<s}^{M} e^{-\frac{i}{2\pi}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}(\sigma_r-\sigma_s)} \times \prod_{r<s}^{M} |e^{i\sigma_r}-e^{i\sigma_s}|^{2\alpha' k_\mu^{(r)}G^{\mu\nu}k_\nu^{(s)}} \\
&\quad \times \prod_{n=1}^{\infty} \exp\left[\frac{\sqrt{2\alpha'}}{n}\sum_{r=1}^{M} k_\mu^{(r)}\left\{\left(\frac{1}{E^T}g\right)^\mu{}_\nu \alpha_{-n}^\nu e^{in\sigma_r} + \left(\frac{1}{E}g\right)^\mu{}_\nu \tilde{\alpha}_{-n}^\nu e^{-in\sigma_r}\right\}\right] \\
&\quad \times g_N\left|\sum_{r=1}^{M} k^{(r)}\right\rangle . \qquad (3.31)
\end{aligned}
$$

We wish to interpret these states as boundary states with off-shell open-string tachyons. This is verified by comparing closed-string tree propagations between these states with the corresponding open-string one-loop amplitudes and testing their coincidence. These will be done in the next subsection.

It is worth mentioning that the present construction is also applicable in a vanishing B field background and in particular the boundary states (3.31) become available just by putting $B_{\mu\nu}=0$.

3.3.1 Construction of dual boundary state

To describe the closed-string propagations we need the dual boundary states. Although they are given by the BPZ conjugation, we apply the previous formalism to their construction as well in order to gain some insight into them. Instead of g_N we use the BPZ dual :

$$
g_N^\dagger = \prod_{n=1}^{\infty}\exp\left\{-\frac{1}{n}N_{\mu\nu}\alpha_n^\mu\tilde{\alpha}_n^\nu\right\}. \qquad (3.32)
$$

The transform of tachyon vertex operator is given by $g_N^\dagger V_T(\sigma,\tau;k)\left(g_N^\dagger\right)^{-1}$. It can be written in the following form for $|z|<1$:

$$
g_N^\dagger V_T(\sigma,\tau;k)\left(g_N^\dagger\right)^{-1} = \left(1-|z|^2\right)^{\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu} \times \mathrm{ad}_{g_N^\dagger}V_T(\sigma,\tau;k). \qquad (3.33)
$$

Here we introduce a local operator $\mathrm{ad}_{g_N^\dagger}V_T$, which we interpret as the (dual) Bogolubov transform of $V_T(\sigma,\tau;k)$. It takes the form of

$$
\mathrm{ad}_{g_N^\dagger}V_T(\sigma,\tau;k) = V_T(\sigma,\tau;k)\mathcal{N}_\infty(\bar{z};k)\tilde{\mathcal{N}}_\infty(z;k), \qquad (3.34)
$$

where $\mathcal{N}_\infty(\bar{z};k)$ and $\tilde{\mathcal{N}}_\infty(z;k)$ are operators consisting only of the annihilation modes :

$$\begin{aligned}\mathcal{N}_\infty(\bar{z};k) &= \prod_{n=1}^{\infty}\exp\left\{-\sqrt{\frac{\alpha'}{2}}\frac{1}{n}k_\mu(g^{-1}N^T)^\mu{}_\nu\alpha_n^\nu\bar{z}^n\right\},\\ \tilde{\mathcal{N}}_\infty(z;k) &= \prod_{n=1}^{\infty}\exp\left\{-\sqrt{\frac{\alpha'}{2}}\frac{1}{n}k_\mu(g^{-1}N)^\mu{}_\nu\tilde{\alpha}_n^\nu z^n\right\}.\end{aligned} \tag{3.35}$$

Singularity of $g_N^\dagger V_T(\sigma,\tau;k)\left(g_N^\dagger\right)^{-1}$ comes from $(1-|z|^2)^{\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu}$ in Eq.(3.33). It becomes the same as the singular factor appearing in Eq.(3.5) after changing the coordinate z to $1/z$.

Similar calculation to what is made to obtain Eq.(3.9) leads to the OPE :

$$\begin{aligned}&\mathrm{ad}_{g_N^\dagger}V_T(\sigma_1,\tau_1;k^{(1)})\mathrm{ad}_{g_N^\dagger}V_T(\sigma_2,\tau_2;k^{(2)})\\ &\quad= \exp\left\{-k_\mu^{(1)}\mathbb{G}_\infty^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)k_\nu^{(2)}\right\}\times :\mathrm{ad}_{g_N^\dagger}V_T(\sigma_1,\tau_1;k^{(1)})\mathrm{ad}_{g_N^\dagger}V_T(\sigma_2,\tau_2;k^{(2)}): \quad ,\end{aligned} \tag{3.36}$$

for $|z_1|\geq|z_2|, |z_1z_2|\leq 1$ and $z_1\neq z_2$. Here $\mathbb{G}_\infty^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)$ is Green's function on the unit disk $|z|\leq 1$. It is defined by

$$\mathbb{G}_\infty^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)\equiv\frac{\langle B_N|X^\mu(\sigma_1,\tau_1)X^\nu(\sigma_2,\tau_2)|\mathbf{0}\rangle}{\langle B_N|\mathbf{0}\rangle}-\frac{\langle\mathbf{0}|\hat{x}_0^\mu\hat{x}_0^\nu|\mathbf{0}\rangle}{\langle\mathbf{0}|\mathbf{0}\rangle}, \tag{3.37}$$

and written down explicitly as follows :

$$\begin{aligned}\mathbb{G}_\infty^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2) &= -\frac{\alpha'}{2}g^{\mu\nu}\ln|z_1-z_2|^2-\frac{\alpha'}{2}(g^{-1}Ng^{-1})^{\mu\nu}\ln(1-z_1\bar{z}_2)\\ &\quad -\frac{\alpha'}{2}(g^{-1}N^Tg^{-1})^{\mu\nu}\ln(1-\bar{z}_1z_2)\,.\end{aligned} \tag{3.38}$$

Similarly to Eq.(3.12), the singular factor appearing in Eq.(3.33) is expressed as the self-contraction of $V_T(\sigma,\tau;k)$ between its chiral and anti-chiral parts. This may be seen as follows. Let $\langle X^\mu(z_1)\tilde{X}^\nu(\bar{z}_2)\rangle_\infty$ be the chiral–anti-chiral correlation of Green's function $\mathbb{G}_\infty^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)$: $\langle X^\mu(z_1)\tilde{X}^\nu(\bar{z}_2)\rangle_\infty\equiv-\frac{\alpha'}{2}(g^{-1}Ng^{-1})^{\mu\nu}\ln(1-z_1\bar{z}_2)$. Using this quantity, we can write the singular factor as

$$\left(1-|z|^2\right)^{\frac{\alpha'}{2}k_\mu\left(g^{-1}Ng^{-1}\right)^{\mu\nu}k_\nu}=\frac{\langle B_N|\,V_T(\sigma,\tau;k)\,|{-k}\rangle}{\langle B_N\,|\mathbf{0}\rangle}=\exp\left\{ik_\mu ik_\nu\left\langle X^\mu(z)\bar{X}^\nu(\bar{z})\right\rangle_\infty\right\}\ . \tag{3.39}$$

Dual of the renormalized open-string tachyon operator, which we call $V_{T\infty}^{ren}$, is given by subtracting the singular factor appearing in Eq.(3.33) as follows:

$$V_{T\infty}^{ren}(\sigma,\tau;k)=\left(1-|z|^2\right)^{-\frac{\alpha'}{2}k_\mu(g^{-1}Ng^{-1})^{\mu\nu}k_\nu}\times V_T(\sigma,\tau;k). \tag{3.40}$$

We remark that the diffS^1 action on the dual vertex operator reduces to Eq.(3.17) after we take the limit $\tau \to 0-$.

Let us describe the dual boundary state. It is obtained by following the same route as taken for Eq.(3.31). Let $V^{ren}_{T\infty}(\sigma_r, \tau_r; k^{(r)})$ $(1 \leq r \leq M)$ be the dual vertex operators. Let $z_r = e^{\tau_r + i\sigma_r}$ be distinct points on the infinite semi-cylinder satisfying the condition, $1 \geq |z_1| \geq |z_2| \geq \cdots \geq |z_M|$. We consider the state $\langle B_N | V^{ren}_{T\infty}(\sigma_1, \tau_1; k^{(1)}) \cdots V^{ren}_{T\infty}(\sigma_M, \tau_M; k^{(M)})$ and take the limit $\forall \tau_r \to 0-$. Thus obtained state, which we call $\big\langle B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \big|$, is the dual. Explicitly it is given by

$$\begin{aligned}
&\big\langle B_N; (\sigma_1, k^{(1)}), (\sigma_2, k^{(2)}), \cdots, (\sigma_M, k^{(M)}) \big| \\
&\quad \equiv \lim_{\forall \tau_r \to 0-} \langle B_N | V^{ren}_{T\infty}\left(\sigma_1, \tau_1; k^{(1)}\right) V^{ren}_{T\infty}\left(\sigma_2, \tau_2; k^{(2)}\right) \cdots V^{ren}_{T\infty}(\sigma_M, \tau_M; k^{(M)}) \\
&\quad = \left(\frac{-\mathrm{det} G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{1/4} \prod_{r<s}^{M} e^{-\frac{i}{2} k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu \epsilon(\sigma_r - \sigma_s)} \\
&\qquad \times \prod_{r<s}^{M} e^{\frac{i}{2\pi} k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu (\sigma_r - \sigma_s)} \times \prod_{r<s}^{M} |e^{i\sigma_r} - e^{i\sigma_s}|^{2\alpha' k^{(r)}_\mu G^{\mu\nu} k^{(s)}_\nu} \\
&\qquad \times \left\langle - \sum_{r=1}^{M} k^{(r)} \right| g_N^\dagger \prod_{n=1}^{\infty} \exp\left[- \frac{\sqrt{2\alpha'}}{n} \sum_{r=1}^{M} k^{(r)}_\mu \left\{ \left(\frac{1}{E^T} g \right)^{\mu}_{\ \nu} \alpha^\nu_n e^{-in\sigma_r} + \left(\frac{1}{E} g \right)^{\mu}_{\ \nu} \tilde{\alpha}^\nu_n e^{in\sigma_r} \right\} \right].
\end{aligned} \tag{3.41}$$

3.3.2 Closed-string propagation

The hamiltonian operator of closed-string is $L_0 + \tilde{L}_0 - 2$. We can conveniently parametrize propagations of closed-string by $\tau^{(c)} \in i\mathbb{R}_{\geq 0}$. The evolution by an imaginary time $\pi |\tau^{(c)}|$ is given by the operator $q_c^{\frac{1}{2}(L_0 + \tilde{L}_0 - 2)}$, where we put $q_c = e^{2\pi i \tau^{(c)}}$. In this subsection we only consider the propagations along the Neumann directions. Propagations of closed-string between the boundary states (3.31) are measured by the following amplitudes :

$$\big\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \big| \, q_c^{\frac{1}{2}(L_0 + \tilde{L}_0 - 2)} \, \big| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \big\rangle \,. \tag{3.42}$$

We calculate these amplitudes by using the oscillator representations. It is useful to recall the following equalities :

$$q_c^{-\frac{1}{2}(L_0 + \tilde{L}_0 - 2)} \alpha^\mu_n q_c^{\frac{1}{2}(L_0 + \tilde{L}_0 - 2)} = q_c^{\frac{n}{2}} \alpha^\mu_n, \qquad q_c^{-\frac{1}{2}(L_0 + \tilde{L}_0 - 2)} \tilde{\alpha}^\mu_n q_c^{\frac{1}{2}(L_0 + \tilde{L}_0 - 2)} = q_c^{\frac{n}{2}} \tilde{\alpha}^\mu_n, \tag{3.43}$$

and

$$\langle -k | q_c^{\frac{1}{2}(L_0 + \tilde{L}_0)} | k' \rangle = q_c^{\frac{\alpha'}{4} g_{\mu\nu} k_\mu k_\nu} \delta^{(p+1)}(k + k'), \tag{3.44}$$

where we put $\delta^{(p+1)}(k) \equiv \prod_\mu \delta(k_\mu)$. First we plug the representations (3.31) and (3.41) into Eq.(3.42). We then evaluate the amplitudes by taking account of the above equalities. They turn out to have the following factorized form :

$$
\begin{aligned}
&\left\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right| q_c^{\frac{1}{2}(L_0+\tilde{L}_0-2)} \left|B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \\
&= \left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{(p+1)}}\right)^{1/2} \delta^{(p+1)}\left(\sum_{r=1}^{M+N} k^{(r)}\right) \times q_c^{-1-\frac{\alpha'}{4}\sum_{r=1}^{M}\sum_{s=M+1}^{M+N} k_\mu^{(r)} g^{\mu\nu} k_\nu^{(s)}} \\
&\quad \times \prod_{1\leq r<s\leq M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)} \times \prod_{M+1\leq r<s\leq M+N} e^{-\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)} \\
&\quad \times \prod_{1\leq r<s\leq M} e^{-\frac{i}{2\pi}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}(\sigma_r-\sigma_s)} \times \prod_{M+1\leq r<s\leq M+N} e^{\frac{i}{2\pi}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}(\sigma_r-\sigma_s)} \\
&\quad \times \prod_{1\leq r<s\leq M} \left|e^{i\sigma_r} - e^{i\sigma_s}\right|^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} \times \prod_{M+1\leq r<s\leq M+N} \left|e^{i\sigma_r} - e^{i\sigma_s}\right|^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} \\
&\quad \times F\left(q_c, \{\sigma_r\}, \{k^{(r)}\}\right) \quad ,
\end{aligned}
\tag{3.45}
$$

where F denotes contributions from the massive modes of closed-string. It is given by the infinite products :

$$
\begin{aligned}
&F\left(q_c, \{\sigma_r\}, \{k^{(r)}\}\right) \\
&\equiv \prod_{n=1}^{\infty} \langle 0| \exp\left[-\frac{q_c^n}{n}\alpha_n^\mu N_{\mu\nu}\tilde{\alpha}_n^\nu \right.\\
&\qquad\qquad \left. -\frac{\sqrt{2\alpha'}q_c^{n/2}}{n}\sum_{r=M+1}^{M+N} k_\mu^{(r)} \left\{ \left(\frac{1}{E^T}g\right)^\mu_{\ \nu} \alpha_n^\nu e^{-in\sigma_r} + \left(\frac{1}{E}g\right)^\mu_{\ \nu} \tilde{\alpha}_n^\nu e^{in\sigma_r} \right\} \right] \\
&\quad \times \exp\left[-\frac{1}{n}\alpha_{-n}^\mu N_{\mu\nu}\tilde{\alpha}_{-n}^\nu \right.\\
&\qquad\qquad \left. +\frac{\sqrt{2\alpha'}}{n}\sum_{r=1}^{M} k_\mu^{(r)} \left\{ \left(\frac{1}{E^T}g\right)^\mu_{\ \nu} \alpha_{-n}^\nu e^{in\sigma_r} + \left(\frac{1}{E}g\right)^\mu_{\ \nu} \tilde{\alpha}_{-n}^\nu e^{-in\sigma_r} \right\} \right] |0\rangle \quad .
\end{aligned}
\tag{3.46}
$$

Further evaluations of the infinite products are carried out in Appendix B. We just quote the result obtained there. The contributions from the massive modes turn out to be as follows:

$$
\begin{aligned}
&F\left(q_c, \{\sigma_r\}, \{k^{(r)}\}\right) \\
&= \prod_{n=1}^{\infty} (1-q_c^n)^{-p-1} \\
&\quad \times \prod_{1\leq r<s\leq M} \left[\frac{\prod_{n=1}^{\infty}\left(1-e^{i(\sigma_r-\sigma_s)}q_c^n\right)\left(1-e^{-i(\sigma_r-\sigma_s)}q_c^n\right)}{\prod_{n=1}^{\infty}(1-q_c^n)^2}\right]^{2\alpha' k_\mu^{(r)} G_{\mu\nu} k_\nu^{(s)}}
\end{aligned}
$$

$$\times \prod_{M+1\le r<s\le M+N} \left[\frac{\prod_{n=1}^{\infty}\left(1-e^{i(\sigma_r-\sigma_s)}q_c^{n}\right)\left(1-e^{-i(\sigma_r-\sigma_s)}q_c^{n}\right)}{\prod_{n=1}^{\infty}\left(1-q_c^{n}\right)^2}\right]^{2\alpha' k_\mu^{(r)} G_{\mu\nu} k_\nu^{(s)}}$$

$$\times \prod_{r=1}^{M}\prod_{s=M+1}^{M+N} \left[\frac{\prod_{n=0}^{\infty}\left(1-e^{i(\sigma_r-\sigma_s)}q_c^{n+1/2}\right)\left(1-e^{-i(\sigma_r-\sigma_s)}q_c^{n+1/2}\right)}{\prod_{n=1}^{\infty}\left(1-q_c^{n}\right)^2}\right]^{2\alpha' k_\mu^{(r)} G_{\mu\nu} k_\nu^{(s)}} . \tag{3.47}$$

For the comparison with open-string one-loop calculation it is convenient to write down the amplitudes (3.45) by using the elliptic θ-functions. Appropriate θ-functions are $\theta_{1,4}(\nu|\tau)$. They have the following representations :

$$\begin{aligned} \theta_1(\nu|\tau) &= 2q^{\frac{1}{8}}\sin\pi\nu\prod_{n=1}^{\infty}(1-q^n)(1-e^{2\pi i\nu}q^n)(1-e^{-2\pi i\nu}q^n), \\ \theta_4(\nu|\tau) &= \prod_{n=1}^{\infty}(1-q^n)\prod_{n=0}^{\infty}(1-e^{2\pi i\nu}q^{n+\frac{1}{2}})(1-e^{-2\pi i\nu}q^{n+\frac{1}{2}}), \end{aligned} \tag{3.48}$$

where we put $q=e^{2\pi i\tau}$.

We first pay attention to $|e^{i\sigma_r}-e^{i\sigma_s}|^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}}$ in Eq.(3.45). It can be combined with infinite products in Eq.(3.47) and gives rise to $\left[\eta(\tau^{(c)})^{-3}\theta_1\left(\frac{|\sigma_r-\sigma_s|}{2\pi}\,\middle|\,\tau^{(c)}\right)\right]^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}}$. Here we have taken into account that $\sigma_{r,s}$ run only from 0 to 2π. We next consider $q_c^{-\frac{\alpha'}{4}\sum_{r=1}^{M}\sum_{s=M+1}^{M+N}k_\mu^{(r)} g^{\mu\nu} k_\nu^{(s)}}$ in Eq.(3.45). Using the first relation of (3.19) it can be written in terms of the open-string tensors as follows :

$$\begin{aligned} & q_c^{-\frac{\alpha'}{4}\sum_{r=1}^{M}\sum_{s=M+1}^{M+N}k_\mu^{(r)} g^{\mu\nu} k_\nu^{(s)}} \\ &\quad = q_c^{\frac{1}{16\pi^2\alpha'}\sum_{r=1}^{M}\sum_{s=M+1}^{M+N}k_\mu^{(r)} (\theta G\theta)^{\mu\nu} k_\nu^{(s)}} \times q_c^{-\frac{\alpha'}{4}\sum_{r=1}^{M}\sum_{s=M+1}^{M+N}k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} . \end{aligned} \tag{3.49}$$

Let us introduce K_μ as the total momentum of the *in*-state : $K \equiv \sum_{r=1}^{M} k^{(r)}$. The first factor can be written as $q_c^{-\frac{1}{16\pi^2\alpha'}K_\mu(\theta G\theta)^{\mu\nu}K_\nu}$ due to the momentum conservation. We then manage $q_c^{-\frac{\alpha'}{4}k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}}$ in the second factor. It can be combined with another infinite products in Eq.(3.47) and gives rise to $\left[\eta(\tau^{(c)})^{-3}\theta_4\left(\frac{(\sigma_r-\sigma_s)}{2\pi}\,\middle|\,\tau^{(c)}\right)\right]^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}}$. Gathering these expressions we finally find out :

$$\begin{aligned} & \left\langle B_N;(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right| q_c^{\frac{1}{2}(L_0+\tilde{L}_0-2)} \left|B_N;(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle \\ & = \left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{(p+1)}}\right)^{1/2} \delta^{p+1}\left(\sum_{r=1}^{M+N}k^{(r)}\right) \end{aligned}$$

$$
\begin{aligned}
&\times \prod_{1\le r<s\le M} e^{\frac{i}{2}k^{(r)}_{\mu}\theta^{\mu\nu}k^{(s)}_{\nu}\epsilon(\sigma_r-\sigma_s)} \times \prod_{M+1\le r<s\le M+N} e^{-\frac{i}{2}k^{(r)}_{\mu}\theta^{\mu\nu}k^{(s)}_{\nu}\epsilon(\sigma_r-\sigma_s)} \\
&\times q_c^{\frac{p-25}{24}}\,\eta(\tau^{(c)})^{-p+1} \\
&\times q_c^{-\frac{1}{16\pi^2\alpha'}K_{\mu}(\theta G\theta)^{\mu\nu}K_{\nu}} \times \prod_{r=1}^{M}\prod_{s=M+1}^{M+N}\left[\eta(\tau^{(c)})^{-3}\theta_4\left(\left.\frac{\sigma_r-\sigma_s}{2\pi}\right|\tau^{(c)}\right)\right]^{2\alpha' k^{(r)}_{\mu}G^{\mu\nu}k^{(s)}_{\nu}} \\
&\times \prod_{r=1}^{M} e^{\frac{\sigma_r}{2\pi i}k^{(r)}_{\mu}\theta^{\mu\nu}K_{\nu}} \times \prod_{1\le r<s\le M}\left[\eta(\tau^{(c)})^{-3}\theta_1\left(\left.\frac{|\sigma_r-\sigma_s|}{2\pi}\right|\tau^{(c)}\right)\right]^{2\alpha' k^{(r)}_{\mu}G^{\mu\nu}k^{(s)}_{\nu}} \\
&\times \prod_{r=M+1}^{M+N} e^{\frac{\sigma_r}{2\pi i}k^{(r)}_{\mu}\theta^{\mu\nu}K_{\nu}} \times \prod_{M+1\le r<s\le M+N}\left[\eta(\tau^{(c)})^{-3}\theta_1\left(\left.\frac{|\sigma_r-\sigma_s|}{2\pi}\right|\tau^{(c)}\right)\right]^{2\alpha' k^{(r)}_{\mu}G^{\mu\nu}k^{(s)}_{\nu}} ,
\end{aligned}
\tag{3.50}
$$

where the products of $e^{\frac{\sigma_r}{2\pi i}k^{(r)}_{\mu}\theta^{\mu\nu}K_{\nu}}$ come from the following translations of the corresponding terms in Eq.(3.45) :

$$
\begin{aligned}
\prod_{1\le r<s\le M} e^{-\frac{i}{2\pi}k^{(r)}_{\mu}\theta^{\mu\nu}k^{(s)}_{\nu}(\sigma_r-\sigma_s)} &= \prod_{r=1}^{M} e^{\frac{\sigma_r}{2\pi i}k^{(r)}_{\mu}\theta^{\mu\nu}\left(\sum_{s=1}^{M}k^{(s)}_{\nu}\right)}, \\
\prod_{M+1\le r<s\le M+N} e^{\frac{i}{2\pi}k^{(r)}_{\mu}\theta^{\mu\nu}k^{(s)}_{\nu}(\sigma_r-\sigma_s)} &= \prod_{r=M+1}^{M+N} e^{\frac{\sigma_r}{2\pi i}k^{(r)}_{\mu}\theta^{\mu\nu}\left(-\sum_{s=M+1}^{M+N}k^{(s)}_{\nu}\right)}.
\end{aligned}
\tag{3.51}
$$

We also note that contribution of the world-sheet reparametrization ghosts, which turns out to be $\prod_{n=1}^{\infty}(1-q_c^n)^2$, has been included in Eq.(3.50). Together with the ghost contribution, q_c^{-1} in Eq.(3.45) and $\prod_{n=1}^{\infty}(1-q_c^n)^{-p-1}$ in F give rise to $q_c^{\frac{p-25}{24}}\eta(\tau^{(c)})^{-p+1}$ in Eq.(3.50).

3.4 Comparison with open-string one-loop calculation

The amplitudes (3.50), combined with closed-string propagations along the Dirichlet directions, should be compared with open-string tachyon one-loop amplitudes. We begin this subsection with a brief exposition on the open-string calculation in a constant B field background of closed-string.

3.4.1 Open-string one-loop calculation

Calculation of open-string one-loop amplitudes in a constant B field background is a direct extension of that in the case of $B=0$ (see e.g. [35]) except a quantization of open-string. Let us first describe the quantization. Action of a bosonic open-string in this background is given by

$$S=\frac{1}{\pi\alpha'}\int\frac{d\bar{\rho}\wedge d\rho}{2i}\Big\{g_{MN}\bar{\partial}_\rho X^M(\rho,\bar{\rho})\partial_\rho X^N(\rho,\bar{\rho})+2\pi\alpha' B_{\mu\nu}\bar{\partial}_\rho X^\mu(\rho,\bar{\rho})\partial_\rho X^\nu(\rho,\bar{\rho})\Big\}, \tag{3.52}$$

where the upper half-plane, Im$\rho \geq 0$, is taken as the world-sheet. An infinite strip (τ,σ) $(0\leq\sigma\leq\pi)$ can be mapped to the upper half-plane by $\rho=e^{\tau+i\sigma}$. Taking variation of the action, one can find that the Neumann and the Dirichlet boundary conditions are imposed on open-string coordinates X^μ and X^i respectively:

$$g_{\mu\nu}(\bar{\partial}_\rho-\partial_\rho)X^\nu+2\pi\alpha' B_{\mu\nu}(\bar{\partial}_\rho+\partial_\rho)X^\nu\Big|_{\rho=\bar{\rho}}=0, \qquad X^i\big|_{\rho=\bar{\rho}}=x_0^i. \tag{3.53}$$

These boundary conditions can be managed in the mode expansions of X^M. They turn out to be as follows :

$$\begin{aligned} X^\mu(\rho,\bar{\rho}) &= \hat{x}_0^\mu - i\alpha'\hat{p}_0^\mu\ln|\rho|^2 + i\sqrt{\frac{\alpha'}{2}}\sum_{n\neq0}\frac{\alpha_n^\mu}{n}(\rho^{-n}+\bar{\rho}^{-n}) \\ &\quad -\frac{i}{2\pi}(\theta G)^\mu{}_\nu\left\{\hat{p}_0^\nu\ln\left(\frac{\rho}{\bar{\rho}}\right)-\frac{1}{\sqrt{2\alpha'}}\sum_{n\neq0}\frac{\alpha_n^\nu}{n}(\rho^{-n}-\bar{\rho}^{-n})\right\}, \\ X^i(\rho,\bar{\rho}) &= x_0^i + i\sqrt{\frac{\alpha'}{2}}\sum_{n\neq0}\frac{\alpha_n^i}{n}(\rho^{-n}-\bar{\rho}^{-n}), \end{aligned} \tag{3.54}$$

where we translate the closed-string tensors into the open-string ones. The quantization is prescribed in [8][9]. It turns out to be realized by the following commutation relations[2] :

$$\begin{aligned} &[\hat{x}_0^\mu,\hat{x}_0^\nu]=i\theta^{\mu\nu}, \quad [\hat{p}_0^\mu,\hat{p}_0^\nu]=0, \quad [\hat{x}_0^\mu,\hat{p}_0^\nu]=iG^{\mu\nu}, \\ &[\alpha_m^\mu,\alpha_n^\nu]=mG^{\mu\nu}\delta_{m+n}, \quad [\alpha_m^i,\alpha_n^j]=mg^{ij}\delta_{m+n}. \end{aligned} \tag{3.55}$$

The commutation relation of $\hat{x}_0^\mu$ indicates the non-commutativity of the world-volume [6][7].

Related with a prescription of normal orderings relative to the $SL_2(\mathbb{R})$-invariant vacuum of open-string, it is convenient to introduce [36] commutative zero-modes $\tilde{x}_0^\mu \equiv \hat{x}_0^\mu+\frac{1}{2}(\theta G)^\mu{}_\nu\hat{p}_0^\nu$ instead of the non-commutative ones. They satisfy the standard canonical relations ; $[\tilde{x}_0^\mu,\tilde{x}_0^\nu]=[\hat{p}_0^\mu,\hat{p}_0^\nu]=0$ and $[\tilde{x}_0^\mu,\hat{p}_0^\nu]=iG^{\mu\nu}$. In what follows we adopt the standard normal ordering of $(\tilde{x}_0^\mu,\hat{p}_0^\mu,\alpha_n^\mu,\alpha_n^i)$. It is also denoted by : :.

The Virasoro generators L_n can be obtained from the energy-momentum tensor $T(\rho)=-\frac{1}{\alpha'}g_{MN}\partial_\rho X^M\partial_\rho X^N$ by $T(\rho)=\sum_m L_m\rho^{-m-2}$. They turn out to have the following forms :

$$L_0 = \alpha' G^{\mu\nu}\hat{p}_0^\mu\hat{p}_0^\nu+\left(\sum_{m=1}^{\infty}G_{\mu\nu}\alpha_{-m}^\mu\alpha_m^\nu+g_{ij}\alpha_{-m}^i\alpha_m^j\right),$$

---

[2] Notations used for open-string are similar to those for closed-string as far as they do not cause any confusion.

$$L_n = \frac{1}{2}\sum_{m\in\mathbb{Z}}\left(G_{\mu\nu}\alpha^\mu_{n-m}\alpha^\nu_m + g_{ij}\alpha^i_{n-m}\alpha^j_m\right) \qquad \text{for} \;\; n \neq 0, \tag{3.56}$$

where we put $\alpha^\mu_0 \equiv \sqrt{2\alpha'}\hat{p}^\mu_0$. Among these generators open-string propagator is given by $\Delta \equiv \frac{1}{L_0-1}$. Evolution of open-string on the upper half-plane is described by ξ^{L_0-2} due to an integral representation $\Delta = \int_0^1 d\xi\, \xi^{L_0-2}$.

It is useful to observe the open-string coordinates, particularly $X^\mu(\rho,\bar{\rho})$, at the boundary of the world-sheet. We may use the radial coordinates (ξ,σ), where $\xi \equiv |\rho|$, as a parametrization of the upper half-plane. They can be read as follows :

$$\begin{aligned} X^\mu(\rho,\bar{\rho})|_{\sigma=0} &= \tilde{x}^\mu_0 - \frac{1}{2}(\theta G)^\mu{}_\nu \hat{p}^\nu_0 - 2i\alpha'\hat{p}^\mu_0 \ln\xi + i\sqrt{2\alpha'}\sum_{n\neq 0}\frac{\alpha^\mu_n}{n}\xi^{-n}, \\ X^\mu(\rho,\bar{\rho})|_{\sigma=\pi} &= \tilde{x}^\mu_0 + \frac{1}{2}(\theta G)^\mu{}_\nu \hat{p}^\nu_0 - 2i\alpha'\hat{p}^\mu_0 \ln\xi + i\sqrt{2\alpha'}\sum_{n\neq 0}\frac{\alpha^\mu_n}{n}(-\xi)^{-n}. \end{aligned} \tag{3.57}$$

At both boundaries $\sigma = 0$ and $\sigma = \pi$, effect of the non-commutativity of the world-volume (the θ-dependence) is encoded only in the zero modes while the massive part acquires the standard mode expansion. Open-string amplitudes are correlation functions between local operators inserted at the boundaries. It follows from Eq.(3.57) that θ-dependence of local operators such as tachyon- and gluon-vertices are encoded only in their zero modes. One-loop amplitudes can be obtained in the operator formalism simply by taking a trace of these operators with the propagators inserting among them. The integral representation of the propagator may be used. The zero mode dependent part decouples from the others in ξ^{L_0-2}. Therefore, for the particular cases of tachyons and gluons, only the zero-modes of open-string are influenced by the non-commutativity of the world-volume.

Let us consider one-loop amplitudes of tachyon vertex operators. We introduce open-string tachyon vertex operator of momentum k by

$$V^{open}_T(\xi,\sigma;k) \equiv \,:e^{ik_\mu X^\mu(\rho,\bar{\rho})}:\Big|_{\rho=\xi e^{i\sigma}} . \tag{3.58}$$

We consider the scattering process of $M+N$ tachyons with momenta $k^{(r)}$. It is worth noting that the momenta $k^{(r)}$ have only components along the Neumann directions. It is because the propagation of open-string is restricted along the D-brane world-volume. Diagram describing the one-loop scattering process can be drawn on the upper half-plane as depicted in Figure 3. An open-string evolves along the radial direction from the outer semi-circle to the inner one and interacts with the tachyon vertices at its ends. Two semi-circles are identified with each other and thereby the diagrams is interpreted as an open-string one-loop. The corresponding



Figure 3: Open-string one-loop diagram drawn on the upper half-plane. Two semi-circles are identified with each other.

tachyon amplitude, which we call $I_T\left((k^{(1)},\cdots,k^{(M)});(k^{(M+1)},\cdots,k^{(M+N)})\right)$, is given by a sum of traces of their products arranged in cyclically distinct orders with keeping their partial cyclic orderings at the both ends :

$$\begin{aligned} & I_T\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\ & \quad \equiv \mathrm{Tr}\left\{\Delta V_T^{open}\left(1,0;k^{(1)}\right)\cdots\Delta V_T^{open}\left(1,0;k^{(M)}\right)\right. \\ & \qquad\qquad \left.\times\,\Delta V_T^{open}\left(1,\pi;k^{(M+1)}\right)\cdots\Delta V_T^{open}\left(1,\pi;k^{(M+N)}\right)\right\} \\ & \qquad + \quad \cdots\cdots. \end{aligned} \tag{3.59}$$

We can evaluate the amplitude in the standard manner. Let us recall that the tachyon vertex operator enjoys the properties :

$$\begin{aligned} \xi^{L_0}V_T^{open}(1,0;k)\xi^{-L_0} &= \xi^{\alpha' k_\mu G^{\mu\nu}k_\nu}V_T^{open}(\xi,0;k), \\ \xi^{L_0}V_T^{open}(1,\pi;k)\xi^{-L_0} &= \xi^{\alpha' k_\mu G^{\mu\nu}k_\nu}V_T^{open}(\xi,\pi;k). \end{aligned} \tag{3.60}$$

Combined with the integral form of Δ, this enables us to write the amplitude as follows :

$$\begin{aligned} & I_T\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\ & = \int_{0\le\xi_{M+N}\le\xi_{M+N-1}\le\cdots\le\xi_1\le 1}\prod_{r=1}^{M+N}\frac{d\xi_r}{\xi_r}\times\prod_{r=1}^{M+N}\xi_r^{\alpha' k_\mu^{(r)}G_{\mu\nu}k_\nu^{(r)}} \\ & \qquad \times\mathrm{Tr}\left\{V_T^{open}\left(\xi_1,0;k^{(1)}\right)\cdots V_T^{open}\left(\xi_M,0;k^{(M)}\right)\right. \\ & \qquad\qquad \left.\times V_T^{open}\left(\xi_{M+1},\pi;k^{(M+1)}\right)\cdots V_T^{open}\left(\xi_{M+N},\pi;k^{(M+N)}\right)\xi_{M+N}^{L_0-1}\right\} \\ & \quad + \quad \cdots\cdots. \end{aligned} \tag{3.61}$$

The coordinates ξ_r, which appear in the RHS as insertion points of the tachyon vertices, can be thought to provide a parametrization of the diagram. Another parametrization may be obtained by mapping the diagram to a cylinder with width π. See Figure 4. Correspondingly ξ_r are mapped to ν_r by $\nu_r = -\ln\xi_r$. We put $\tau^{(o)} \equiv -\ln\xi_{M+N}$. We may use $\tau^{(o)}$ instead of ν_{M+N}. The evolution of open-string becomes manifest in this parameterization. In the figure an open-string with width π evolves along the real axis from the origin to $\tau^{(0)}$, interacting with the tachyon vertices inserted at ν_r or $\nu_r + i\pi$.

After a little calculation of the traces in the RHS of Eq.(3.61) we can obtain the amplitude in an integral form. In terms of $\tau^{(o)}$ and ν_r this integral can be written as follows :

$$\begin{aligned} & I_T\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\ & = \sqrt{-\mathrm{det}G_{\mu\nu}}\,\delta^{(p+1)}\left(\sum_{r=1}^{M+N}k^{(r)}\right) \end{aligned}$$



Figure 4: Open-string one-loop diagram drawn on the u-plane. Two bold vertical lines are identified with each other.

$$
\begin{aligned}
&\times \prod_{1\le r<s\le M} e^{-\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}} \times \prod_{M+1\le r<s\le M+N} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}} \\
&\times \int d\tau^{(o)} \prod_{r=1}^{M+N-1} d\nu_r \left(\frac{\pi}{\alpha'\tau^{(o)}}\right)^{\frac{p+1}{2}} e^{\frac{26-D}{24}\tau^{(o)}} \eta\left(-\frac{\tau^{(o)}}{2\pi i}\right)^{-D+2} \\
&\quad \times e^{\frac{1}{4\alpha'\tau^{(o)}}K_\mu(\theta G\theta)^{\mu\nu}K_\nu} \times \prod_{r=1}^{M+N} e^{-i\frac{\nu_r}{\tau^{(o)}}k_\mu^{(r)}\theta^{\mu\nu}K_\nu} \\
&\quad \times \prod_{r=1}^{M}\prod_{s=M+1}^{M+N}\left[e^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\frac{(\nu_s+i\pi)-\nu_r}{2\pi i}\middle|\frac{-\tau^{(o)}}{2\pi i}\right)\right]^{2\alpha' k_\mu^{(r)}G^{\mu\nu}k_\nu^{(s)}} \\
&\quad \times \prod_{1\le r<s\le M}\left[ie^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\frac{\nu_s-\nu_r}{2\pi i}\middle|\frac{-\tau^{(o)}}{2\pi i}\right)\right]^{2\alpha' k_\mu^{(r)}G^{\mu\nu}k_\nu^{(s)}} \\
&\quad \times \prod_{M+1\le r<s\le M+N}\left[ie^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\frac{\nu_s-\nu_r}{2\pi i}\middle|\frac{-\tau^{(o)}}{2\pi i}\right)\right]^{2\alpha' k_\mu^{(r)}G^{\mu\nu}k_\nu^{(s)}} ,
\end{aligned}
\tag{3.62}
$$

where the integral is performed over the region :

$$
\begin{aligned}
&\tau^{(o)} \in \mathbb{R}_{\ge 0}, \\
&0 \le \nu_1 \le \cdots \le \nu_M \le \tau^{(0)}, \\
&0 \le \nu_{M+1} \le \cdots \le \nu_{M+N-1} \le \tau^{(0)}.
\end{aligned}
\tag{3.63}
$$

Contribution of the world-sheet reparametrization ghosts, which is equal to $\prod_{n=1}^{\infty}(1-e^{-n\tau^{(o)}})^2$, has been included. We note that K_μ used in the above is given by $K_\mu = \sum_{r=1}^{M} k_\mu^{(r)}$.

3.4.2 Comparison with one-loop amplitudes of open-string

To compare the open-string one-loop amplitude (3.62) with Eq.(3.50) we map the cylinder drawn on the u-plane (Figure 4) to a cylinder with width $2\pi^2/\tau^{(o)}$ on the v-plane by the conformal transformation $v = 2\pi u/\tau^{(o)}$. See Figure 5. ν_r are mapped to σ_r by $\sigma_r = 2\pi\nu_r/\tau^{(o)}$. We put $\tau^{(c)} \equiv 2\pi i/\tau^{(o)}$. These provide a new parametrization of the diagram, which makes an evolution of closed-string manifest. A closed-string with circumference 2π evolves along the imaginary axis, starting from the real axis where it interacts with M open-string tachyons and ending on $\mathrm{Im}v = \pi|\tau^{(c)}|$ where it interacts with N open-string tachyons. We note that the insertion point of the $(M+N)$-th open-string tachyon is fixed at $\sigma_{M+N} = 2\pi$.

Let us describe the one-loop scattering amplitude of $M+N$ tachyons as an integral with respect to $\tau^{(c)}$ and σ_r. By making use of the modular transform, $\theta_1(\nu/\tau|-1/\tau) = -i\sqrt{-i\tau}e^{\pi i\nu^2/\tau}\theta_1(\nu|\tau)$,



Figure 5: Open-string one-loop diagram drawn on the v-plane. Two bold vertical lines are identified with each other.

we can express several correlations appearing in Eq.(3.62) by means of $\tau^{(c)}$ and σ_r :

$$\begin{aligned}
&e^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\frac{(\nu_s+i\pi)-\nu_r}{2\pi i}\middle|\,\frac{-\tau^{(o)}}{2\pi i}\right)=(-i\tau^{(c)})^{-1}\eta(\tau^{(c)})^{-3}\theta_4\left(\frac{\sigma_s-\sigma_r}{2\pi}\middle|\,\tau^{(c)}\right),\\
&ie^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\frac{\nu_s-\nu_r}{2\pi i}\middle|\,\frac{-\tau^{(o)}}{2\pi i}\right)=(-i\tau^{(c)})^{-1}\eta(\tau^{(c)})^{-3}\theta_1\left(\frac{\sigma_s-\sigma_r}{2\pi}\middle|\,\tau^{(c)}\right).
\end{aligned}\tag{3.64}$$

Other ingredients can be also expressed in terms of these parameters. The integral turns out to be as follows :

$$\begin{aligned}
&I_T\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right)\\
&=\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{1/2}\delta^{(p+1)}\left(\sum_{r=1}^{M+N}k^{(r)}\right)\\
&\quad\times\prod_{1\le r<s\le M}e^{-\frac{i}{2}k^{(r)}_\mu\theta^{\mu\nu}k^{(s)}_\nu}\times\prod_{M+1\le r<s\le M+N}e^{\frac{i}{2}k^{(r)}_\mu\theta^{\mu\nu}k^{(s)}_\nu}\\
&\quad\times(-2\pi i)\int_{i0}^{i\infty}d\tau^{(c)}\int\prod_{r=1}^{M+N-1}d\sigma_r\;e^{\frac{26-D}{24}\frac{2\pi i}{\tau^{(c)}}}\left(-i\tau^{(c)}\right)^{-\frac{d}{2}}\eta(\tau^{(c)})^{-D+2}\\
&\qquad\times\prod_{r=1}^{M+N}\left(-i\tau^{(c)}\right)^{\alpha'k^{(r)}_\mu G^{\mu\nu}k^{(r)}_\nu-1}\\
&\qquad\times e^{\frac{\tau^{(c)}}{8\pi i\alpha'}K_\mu(\theta G\theta)^{\mu\nu}K_\nu}\times\prod_{r=1}^{M}\prod_{s=M+1}^{M+N}\left[\eta(\tau^{(c)})^{-3}\theta_4\left(\frac{\sigma_s-\sigma_r}{2\pi}\middle|\,\tau^{(c)}\right)\right]^{2\alpha'k^{(r)}_\mu G^{\mu\nu}k^{(s)}_\nu}\\
&\qquad\times\prod_{r=1}^{M}e^{\frac{\sigma_r}{2\pi i}k^{(r)}_\mu\theta^{\mu\nu}K_\nu}\times\prod_{1\le r<s\le M}\left[\eta(\tau^{(c)})^{-3}\theta_1\left(\frac{\sigma_s-\sigma_r}{2\pi}\middle|\,\tau^{(c)}\right)\right]^{2\alpha'k^{(r)}_\mu G^{\mu\nu}k^{(s)}_\nu}\\
&\qquad\times\prod_{r=M+1}^{M+N}e^{\frac{\sigma_r}{2\pi i}k^{(r)}_\mu\theta^{\mu\nu}K_\nu}\times\prod_{M+1\le r<s\le M+N}\left[\eta(\tau^{(c)})^{-3}\theta_4\left(\frac{\sigma_s-\sigma_r}{2\pi}\middle|\,\tau^{(c)}\right)\right]^{2\alpha'k^{(r)}_\mu G^{\mu\nu}k^{(s)}_\nu},
\end{aligned}\tag{3.65}$$

where the integration region becomes as follows :

$$\begin{aligned}
&\tau^{(c)}\in i\mathbb{R}_{\ge0},\\
&0\le\sigma_1\le\cdots\le\sigma_M\le2\pi,\\
&0\le\sigma_{M+1}\le\cdots\le\sigma_{M+N-1}\le2\pi.
\end{aligned}\tag{3.66}$$

The above integral form allows us to express the one-loop amplitude of $M+N$ open-string tachyons by using the closed-string amplitude given in Eq.(3.50). We need to include the contribution of closed-string propagations along the Dirichlet directions. This is carried out by multiplying Eq.(3.50) by $\langle B_D|q_c^{\frac{L_0+\tilde{L}_0}{2}}|B_D\rangle = q_c^{d/24}(-i\tau^{(c)})^{-d/2}\eta(\tau^{(c)})^{-d}$. Thus we finally obtain the following equality :

$$
\begin{aligned}
&I_T\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right)\\
&=(-2\pi i)\int_{i0}^{i\infty}d\tau^{(c)}\;e^{2\pi i\left(\tau^{(c)}+\frac{1}{\tau^{(c)}}\right)\frac{26-D}{24}}\prod_{r=1}^{M+N}\left(-i\tau^{(c)}\right)^{\alpha' k^{(r)}_{\mu}G^{\mu\nu}k^{(r)}_{\nu}-1}\\
&\qquad\left[\int\prod_{r=M+1}^{M+N-1}d\sigma_r\,\langle B_D|\otimes\left\langle B_N;(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right|\right]\\
&\qquad\qquad\times q_c^{\frac{L_0+\tilde{L}_0-2}{2}}\left[\int\prod_{r=1}^{M}d\sigma_r\left|B_N;(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle\otimes|B_D\rangle\right],
\end{aligned}
\tag{3.67}
$$

where the integration is performed on the region (3.66). Additional two factors appearing in the RHS vanish when we impose the on-shell conditions on $k^{(r)}$ at the critical dimensions ($D = 26$). Thus we showed that our boundary states (3.31) correctly reproduce the corresponding open-string one-loop amplitudes. We note that this computation implies as well that normalization factor used in Eq.(2.25) is properly chosen to reproduce the corresponding amplitudes.

4 UV limit of Non-Commutative Scalar Field Theory

Low-energy description of world-volume theories of p-brane can be obtained by taking a zero-slope limit $\alpha' \to 0$ so that it makes all perturbative stringy states (mass$^2 \sim 1/\alpha'$) of open-string infinitely heavy and decouple from the light states. Low-energy effective theory relevant to open-string tachyons in the presence of a constant B field is a scalar field theory on the non-commutative world-volume. In general, quantum field theories on a non-commutative space suffer the UV-IR mixing [11] which originates in the non-commutativity. It causes [37] a serious problem on the renormalization prescription of these theories. In this section, based on the results obtained so far, we investigate the UV behavior of the non-commutative scalar field theory. For this sake it is convenient to start with a brief description of the above zero-slope limit of the one-loop amplitudes of $M + N$ tachyons. Our study in this section is restricted to the case of 25-brane in the critical dimensions.

The amplitude is described in (3.62) as an integral, where $\tau^{(o)}$ and ν_r are used as the integral variables. Since these parameters make an evolution of open-string manifest in the scattering process, we call them open-string parameters. For the same reason we call $\tau^{(c)}$ and σ_r closed-string parameters. The zero-slope limit of the amplitude (3.62) will be a field theory one-loop amplitude, particularly written in the Schwinger representation [12][13][14][15][16][32]. Open-string parameters are translated to the Schwinger parameters in the low-energy world-volume theory. For the dimensional reasons we put

$$s^{(o)} \equiv \alpha' \tau^{(o)}, \qquad T_r \equiv \alpha' \nu_r. \tag{4.1}$$

In order to obtain a proper field theory limit of open-string the field theory parameters $s^{(o)}$ and T_r need to be fixed under the zero-slope limit. Simultaneously we also need to fix open-string tensors $G_{\mu\nu}$ and $\theta^{\mu\nu}$ since they describe a classical geometry of the world-volume. In an actual derivation of the zero-slope limit we first rewrite the integral in terms of these field theory parameters and then pick up the dominant contribution of the α'-expansion. Let us take a look at correlations appearing in Eq.(3.62). Terms which become dominant after the above translations can be read as follows :

$$\begin{aligned}
&\left[e^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}} \eta \left(\frac{-\tau^{(o)}}{2\pi i} \right)^{-3} \theta_1 \left(\frac{(\nu_s + i\pi) - \nu_r}{2\pi i} \middle| \frac{-\tau^{(o)}}{2\pi i} \right) \right]^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} \\
&\qquad\qquad \approx \left[e^{-\frac{(T_s-T_r)^2}{s^{(o)}} + |T_s - T_r|} \right]^{k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} , \\
&\left[i e^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}} \eta \left(\frac{-\tau^{(o)}}{2\pi i} \right)^{-3} \theta_1 \left(\frac{\nu_s - \nu_r}{2\pi i} \middle| \frac{-\tau^{(o)}}{2\pi i} \right) \right]^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} \\
&\qquad\qquad \approx \left[e^{-\frac{(T_s-T_r)^2}{s^{(o)}} + |T_s - T_r|} \right]^{k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} .
\end{aligned} \tag{4.2}$$

We also note that the dominant contribution of $\eta(-\tau^{(o)}/2\pi i)^{-24}$ in Eq.(3.62) is $e^{s^{(o)}/\alpha'}$. Gathering these estimations the following integral [12][13][14][15][16][32] turns out to be the zero-slope limit :

$$\begin{aligned}
&I_T \left(\left(k^{(1)}, \cdots, k^{(M)} \right); \left(k^{(M+1)}, \cdots, k^{(M+N)} \right) \right) \\
&\approx \frac{\pi^{13}}{\alpha'^{M+N}} \sqrt{-\mathrm{det} G_{\mu\nu}} \delta^{26} \left(\sum_{r=1}^{M+N} k^{(r)} \right) \prod_{1 \le r < s \le M} e^{-\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)}} \times \prod_{M+1 \le r < s \le M+N} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)}} \\
&\quad \times \int ds^{(o)} (s^{(o)})^{-13} \exp \left\{ \frac{s^{(o)}}{\alpha'} + \frac{K_\mu (\theta G \theta)^{\mu\nu} K_\nu}{4 s^{(o)}} \right\} \int \prod_{r=1}^{M+N-1} dT_r \\
&\qquad\qquad \times \prod_{r=1}^{M+N} e^{-i \frac{T_r}{s^{(0)}} k_\mu^{(r)} \theta^{\mu\nu} K_\nu} \times \prod_{r<s}^{M+N} \left[e^{-\frac{(T_r - T_s)^2}{s^{(o)}} + |T_r - T_s|} \right]^{k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} ,
\end{aligned} \tag{4.3}$$

where the integral is performed over the region :

$$\begin{aligned}
&s^{(o)} \in \mathbb{R}_{\geq 0}, \\
&0 \leq T_1 \leq \cdots \leq T_M \leq s^{(o)}, \\
&0 \leq T_{M+1} \leq \cdots T_{M+N-1} \leq s^{(o)}.
\end{aligned} \tag{4.4}$$

Eq.(4.3) can be identified with one-loop amplitude of open-string tachyon ϕ (a scalar particle of mass$^2 = -1/\alpha'$) living on the non-commutative world-volume. It is the amplitude obtained from the corresponding one-loop Feynman diagram consisting of $M+N$ trivalent vertices. Each of the vertices represents the following cubic interaction :

$$\mathcal{L}_{int} = \frac{1}{3}\phi \star \phi \star \phi(x). \tag{4.5}$$

Here we introduce the Moyal product (an associative non-commutative product) by

$$f \star g(x) \equiv \lim_{y \to x} \exp\left\{ \frac{i}{2}\theta^{\mu\nu}\frac{\partial}{\partial x^\mu}\frac{\partial}{\partial y^\nu} \right\} f(x)g(y). \tag{4.6}$$

The Feynman rule becomes a little complicated due to the above Moyal products. But the Feynman integral can be evaluated by using the standard technique [39] and translated to Eq.(4.3).

We can find $\exp\left\{\frac{s^{(o)}}{\alpha'} + \frac{K_\mu(\theta G\theta)^{\mu\nu}K_\nu}{4s^{(o)}}\right\} = \exp\left\{\frac{s^{(o)}}{\alpha'}\right\} \times \exp\left\{\frac{K_\mu(\theta G\theta)^{\mu\nu}K_\nu}{4s^{(o)}}\right\}$ in the integral of Eq.(4.3). The first term comes from the Schwinger representation of the tachyon propagator, $\exp\left\{-s^{(0)}(p^2 - \frac{1}{\alpha'})\right\}$. This is the standard term which appears in ordinary field theory one-loop amplitudes. The second term can be understood [11] as a UV regularization (a regularization of the integration near $s^{(o)} = 0$). It is a curious regularization since it depends on the external momentum, $K = \sum_{r=1}^{M} k^{(r)}$. This feature can be thought of as a characteristic of quantum field theories on a non-commutative space and makes field theoretical description of physics at high energy scale difficult. It should be also noted that integrations in Eq.(4.3) have potential singularities at $T_r = T_s$ $(r \neq s)$. These singularities can be already seen in the integral (3.62). Although we do not describe their regularization here, prescription used [34] for the cubic open-string field theory will be effective.

4.1 UV limit of non-commutative scalar field theory

Both parameters $(\tau^{(o)}, \nu_r)$ and $(\tau^{(c)}, \sigma_r)$ are regarded as coordinates of the moduli space of conformal classes of cylinder with $M+N$ punctures at the boundaries. For each values of the closed-string parameters we obtain a graph as depicted in Figure 5. Set of these graphs is the above moduli space. Strictly speaking, these graphs are representatives of the conformal classes and thus it is possible to make different choices. For each values of the open-string parameters we obtain a graph as depicted in Figure 4. It can be identified in the standard manner with a metrized trivalent one-loop ribbon graph. Metric of the graph is given by the open-string parameters. These metrized ribbon graphs can be chosen as the representatives. Therefore the open-string parameters give another set of coordinates of the moduli space.

The above moduli space has several ends. Typically we have two ends at $\tau^{(o)} = 0$ and $\tau^{(o)} = +\infty$. The previous zero-slope limit of open-string is related with the end at the infinity as can be seen from Eq.(4.1). Actual procedure to obtain the zero-slope limit (4.3) shows that the field theory amplitude is a suitable magnification of the integral (3.62) on an infinitesimal neighbourhood around this end. We want to know the UV behavior of the world-volume theory. For such a purpose we need to focus on the region $s^{(o)} \approx 0$. One possible resolution may be obtained by taking a zero-slope limit such that it magnifies an infinitesimal neighbourhood around another end located at $\tau^{(o)} = 0$. Near this end use of the closed-string parameters will be effective as the open-string parameters near the infinity. Let us examine a zero-slope limit which will be taken by fixing the following parameters :

$$s^{(c)} \equiv \alpha' |\tau^{(c)}|, \qquad \sigma_r. \tag{4.7}$$

In the course of taking the limit we also fix open-string tensors $G_{\mu\nu}$ and $\theta^{\mu\nu}$ in order to capture the world-volume theory. It follows from Eq.(3.19) that this makes the limit different from the standard field theory limit of closed-string. Transformation between the open- and the closed-string parameters leads $s^{(c)} = 2\pi\alpha' \cdot \frac{\alpha'}{s^{(o)}}$ and $\sigma_r = 2\pi \frac{T_r}{s^{(o)}}$. For $s^{(c)}$ and σ_r to remain finite the original field theory parameters need to satisfy $s^{(o)} \ll \alpha'$ and $T_r \approx s^{(o)}$. We cannot neglect effects of all the perturbative stringy states of open-string at such a trans-string scale. Hence the limit in question includes their effects.

Relevant integral form of one-loop amplitude of $M+N$ open-string tachyons is given by Eq.(3.65) using the closed-string parameters or equivalently by Eq.(3.67) using the boundary states. We generalize the amplitude by dropping out the factor $\prod_{r=1}^{M+N} (-i\tau^{(c)})^{\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(r)} - 1}$ in the integral and introduce the following quantity [3] :

$$\begin{aligned} & J_T\left(\left(k^{(1)}, \cdots, k^{(M)}\right); \left(k^{(M+1)}, \cdots, k^{(M+N)}\right)\right) \\ & \equiv (-2\pi i) \int_{i0}^{i\infty} d\tau^{(c)} \left[\int \prod_{r=M+1}^{M+N-1} d\sigma_r \left\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \right| \right] \end{aligned}$$

---

[3] We put $p = 25$ and $D = 26$.

$$\times\, q_c^{\frac{L_0+\tilde{L}_0-2}{2}} \left[\int \prod_{r=1}^{M} d\sigma_r \left| B_N ; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \right], \tag{4.8}$$

where the integral is performed over the region (3.66). σ_{M+N} is fixed to 2π. When $k^{(r)}$ satisfy the on-shell conditions the above J_T coincides with the original amplitude. It is an off-shell generalization slightly different from the original one.

Let us take the zero-slope limit. We first look at correlations which appear in Eq.(4.8). They can be read from Eq.(3.65). Terms which become dominant after the translations (4.7) are as follows :

$$\begin{aligned} \left[\eta\left(\tau^{(c)}\right)^{-3} \theta_4 \left(\frac{\sigma_s - \sigma_r}{2\pi} \middle| \tau^{(c)} \right) \right]^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} &\approx e^{\frac{\pi}{2} s^{(c)} k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}}, \\ \left[\eta\left(\tau^{(c)}\right)^{-3} \theta_1 \left(\frac{\sigma_s - \sigma_r}{2\pi} \middle| \tau^{(c)} \right) \right]^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} &\approx 1. \end{aligned} \tag{4.9}$$

We also note that the dominant contribution of $\eta(\tau^{(c)})^{-24}$ in Eq.(3.65) is $e^{\frac{2\pi}{\alpha'} s^{(c)}}$. Gathering these estimations the following integral turns out to be the zero-slope limit :

$$\begin{aligned} & J_T\left(\left(k^{(1)}, \cdots, k^{(M)}\right); \left(k^{(M+1)}, \cdots, k^{(M+N)}\right) \right) \\ & \approx \frac{2\pi}{\alpha'} \delta^{26}\left(\sum_{r=1}^{M+N} k^{(r)} \right) \left[\int ds^{(c)} \exp\left\{ -\frac{\pi s^{(c)}}{2} \left(K_\mu g^{\mu\nu} K_\nu - \frac{4}{\alpha'} \right) \right\} \right] \\ & \quad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{1/4} \int \prod_{r=1}^{M} d\sigma_r \prod_{1 \le r < s \le M} e^{-\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)}} \prod_{r=1}^{M} e^{-i k_\mu^{(r)} \theta^{\mu\nu} K_\nu \frac{\sigma_r}{2\pi}} \\ & \quad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{1/4} \int \prod_{r=M+1}^{M+N-1} d\sigma_r \prod_{M+1 \le r < s \le M+N} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)}} \prod_{r=M+1}^{M+N} e^{-i k_\mu^{(r)} \theta^{\mu\nu} K_\nu \frac{\sigma_r}{2\pi}}, \end{aligned} \tag{4.10}$$

where the integral is performed over the region : (σ_{M+N} is fixed to 2π.)

$$\begin{aligned} & s^{(c)} \in \mathbb{R}_{\ge 0}, \\ & 0 \le \sigma_1 \le \cdots \le \sigma_M \le 2\pi, \\ & 0 \le \sigma_{M+1} \le \cdots \le \sigma_{M+N-1} \le 2\pi. \end{aligned} \tag{4.11}$$

Here the closed-string metric $g^{\mu\nu}$ should be understood as $g^{\mu\nu} \approx -\frac{(\theta G \theta)^{\mu\nu}}{(2\pi\alpha')^2}$ by using Eq.(3.19).

It is important to compare the above zero-slope limit with the previous one. In Eq.(4.10) the Schwinger propagator $\exp\left\{ -\frac{\pi s^{(c)}}{2} \left(K_\mu g^{\mu\nu} K_\nu - \frac{4}{\alpha'} \right) \right\}$ is integrated and gives rise to the propagator of *closed-string tachyon* of momentum K, while the counterpart in Eq.(4.3) consists of two parts one of which is standard in field theories and the other is a curious regularization factor originated in the non-commutativity. The factors $\left[\exp\left\{-\frac{(T_r-T_s)^2}{s^{(o)}}+|T_r-T_s|\right\}\right]^{k_\mu^{(r)}G^{\mu\nu}k_\nu^{(s)}}$ found in Eq.(4.3) describe correlations between two open-string tachyons inserted at T_r and T_s in the Schwinger time. These correlations are made by propagations of open-string tachyon between them and particularly have their origin in kinetic energy p^2 of the propagating open-string tachyon. These correlations are lost in Eq.(4.10). This indicates that open-string tachyon becomes topological at the trans-string scale and that it is described by a one-dimensional topological field theory. Vanishing of the correlations originates in the modular transforms (3.64). So this is a stringy effect.

4.2 Factorization by straight open Wilson lines

We now consider generating function of amplitudes (4.8) and examine the zero-slope limit. Let $\varphi(k)$ be the Fourier modes of open-string tachyon field : $\phi(x)=\int\frac{dk}{(2\pi)^{13}}\varphi(k)e^{ik_\mu x^\mu}$. Since it is a real scalar field, the Fourier modes satisfy $\overline{\varphi(k)}=\varphi(-k)$, where the overline denotes complex conjugation. We examine the zero-slope limit of the following generating function :

$$\sum_{M=0}^{\infty}\sum_{N=1}^{\infty}\int\prod_{r=1}^{M+N}\frac{dk^{(r)}}{(2\pi)^{13}}J_T\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(-k^{(M+1)},\cdots,-k^{(M+N)}\right)\right)$$
$$\times\prod_{r=1}^{M}\varphi(k^{(r)})\prod_{r=M+1}^{M+N}\varphi(-k^{(r)}). \tag{4.12}$$

Let us start with taking the zero-slope limit term by term in the sum. The total momentum is conserved in each term. It can be translated to the following integral :

$$\begin{aligned}
\delta^{26}\left(\sum_{r=1}^{M}k^{(r)}-\sum_{r=M+1}^{M+N}k^{(r)}\right)&=\int dp\,\delta^{26}\left(p+\sum_{r=1}^{M}k^{(r)}\right)\delta^{26}\left(p+\sum_{r=M+1}^{M+N}k^{(r)}\right)\\
&=\int dp\,\delta^{26}\left(p+\sum_{r=1}^{M}k^{(r)}\right)e^{-\frac{i}{2}p_\mu\theta^{\mu\nu}\sum_{r=1}^{M}k_\nu^{(r)}}\,\delta^{26}\left(p+\sum_{r=M+1}^{M+N}k^{(r)}\right)e^{\frac{i}{2}p_\mu\theta^{\mu\nu}\sum_{r=M+1}^{M+N}k_\nu^{(r)}}\\
&=\int dp\left(\int\frac{dx}{(2\pi)^{26}}e^{ip_\mu x^\mu}e^{ik_\mu^{(r)}x^\mu}\right)e^{-\frac{i}{2}p_\mu\theta^{\mu\nu}\sum_{r=1}^{M}k_\nu^{(r)}}\\
&\qquad\times\left(\int\frac{dy}{(2\pi)^{26}}e^{-ip_\mu y^\mu}e^{-ik_\mu^{(r)}y^\mu}\right)e^{\frac{i}{2}p_\mu\theta^{\mu\nu}\sum_{r=M+1}^{M+N}k_\nu^{(r)}}.
\end{aligned}\tag{4.13}$$

The zero-slope limit can be obtained by using Eq.(4.10). It turns out to be expressed in terms of the Moyal products. The above rearrangement of the momentum conservation is used in this course. It yields the following products :

$$\int \prod_{r=1}^{M} \frac{dk^{(r)}}{(2\pi)^{13}} e^{ip_\mu x^\mu} e^{-\frac{i}{2}p_\mu \theta^{\mu\nu} \sum_{r=1}^{M} k_\nu^{(r)}} \prod_{1\le r<s\le M} e^{-i\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}} \prod_{r=1}^{M} \varphi(k^{(r)}) e^{ik_\mu^{(r)}\left(x^\mu+\theta^{\mu\nu}p_\nu \frac{\sigma_r}{2\pi}\right)}$$
$$= e^{ip_\mu x^\mu} \star \phi(x_0(\sigma_1)) \star \phi(x_0(\sigma_2)) \star \cdots \star \phi(x_0(\sigma_M)). \tag{4.14}$$

Here we introduce straight line $x_0(\sigma)$ in the world-volume by $x_0^\mu(\sigma) \equiv x^\mu + \theta^{\mu\nu}p_\nu \frac{\sigma}{2\pi}$ with $0 \le \sigma \le 2\pi$, and the Moyal products are taken with respect to x. After these replacements the zero-slope limit turns out to be :

$$\int \prod_{r=1}^{M+N} \frac{dk^{(r)}}{(2\pi)^{13}} J_T\left(\left(k^{(1)}, \cdots, k^{(M)}\right) ; \left(-k^{(M+1)}, \cdots, -k^{(M+N)}\right)\right)$$
$$\times \prod_{r=1}^{M} \varphi(k^{(r)}) \prod_{r=M+1}^{M+N} \varphi(-k^{(r)})$$
$$\approx \frac{2\pi}{\alpha'} \int dp \left[\int_0^\infty ds^{(c)} \exp\left\{ -\frac{\pi s^{(c)}}{2} \left(p_\mu g^{\mu\nu} p_\nu - \frac{4}{\alpha'} \right) \right\} \right]$$
$$\times \left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{4}} \int \frac{dx}{(2\pi)^{26}} e^{ip_\mu x^\mu} \star \left[\int \prod_{r=1}^{M} d\sigma_r \phi(x_0(\sigma_1)) \star \cdots \star \phi(x_0(\sigma_M)) \right]$$
$$\times \overline{\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{4}} \int \frac{dx}{(2\pi)^{26}} e^{ip_\mu x^\mu} \star \left[\int \prod_{r=M+1}^{M+N-1} d\sigma_r \phi(x_0(\sigma_{M+1})) \star \cdots \star \phi(x_0(\sigma_{M+N})) \right]} \quad , \tag{4.15}$$

where the integrations on σ_r are performed over the region (4.11).

The zero-slope limit of the generating function (4.12) can be obtained by summing up Eq.(4.15) with respect to M and N. Let us introduce path-ordered exponential along a line $x^\mu(\sigma)$. It is defined by

$$\mathcal{P}_\star \left[\exp\left\{ \int_0^{2\pi} d\sigma \phi(x(\sigma)) \right\} \right] \equiv \sum_{M=0}^{\infty} \int_{0\le\sigma_1\le\cdots\le\sigma_M\le 2\pi} \prod_{r=1}^{M} d\sigma_r \phi(x(\sigma_1)) \star \cdots \star \phi(x(\sigma_M)) \quad . \tag{4.16}$$

This is an analogue of the Wilson line in gauge theories. Each summation of the Moyal products of Eq.(4.15) gives the path-ordered exponential along the straight line. The zero-slope limit can be written as follows :

$$\sum_{M=0}^{\infty} \sum_{N=1}^{\infty} \int \prod_{r=1}^{M+N} \frac{dk^{(r)}}{(2\pi)^{13}} J_T\left(\left(k^{(1)}, \cdots, k^{(M)}\right) ; \left(-k^{(M+1)}, \cdots, -k^{(M+N)}\right)\right)$$
$$\times \prod_{r=1}^{M} \varphi(k^{(r)}) \prod_{r=M+1}^{M+N} \varphi(-k^{(r)})$$

$$
\begin{aligned}
&\approx \frac{2\pi}{\alpha'}\int dp\left[\int_0^\infty ds^{(c)}\exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_\mu g^{\mu\nu}p_\nu-\frac{4}{\alpha'}\right)\right\}\right]\\
&\qquad\times\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}}\int\frac{dx}{(2\pi)^{26}}e^{ip_\mu x^\mu}\star\mathcal{P}_\star\left[\exp\left\{\int_0^{2\pi}d\sigma\phi(x_0(\sigma))\right\}\right]\\
&\qquad\times\overline{\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}}\int\frac{dx}{(2\pi)^{26}}e^{ip_\mu x^\mu}\star\mathcal{P}_\star\left[\exp\left\{\int_0^{2\pi}d\sigma\phi(x_0(\sigma))\right\}\right]\star\phi(x_0(2\pi))}\ .
\end{aligned}
\tag{4.17}
$$

Insertion of $\phi(x_0(2\pi))$ makes the above equation asymmetric. This term has its origin in one-loop calculation of open-string or tachyon field on the world-volume. The one-loop diagram has a $U(1)$ symmetry which generates a constant simultaneous shift of the Schwinger parameters. This $U(1)$ symmetry is fixed in the previous discussion by putting $\tau^{(o)}=-\ln\xi_{M+N}$. It becomes equivalent to $\sigma_{M+N}=2\pi$. Modulo fixing the $U(1)$ symmetry, Eq.(4.17) shows that the generating function is factorized at the zero-slope limit into a sum of products of $\mathcal{P}_\star\left[e^{\int_0^{2\pi}d\sigma\phi(x_0(\sigma))}\right]$ and that open Wilson lines with the same velocity $\frac{dx_0^\mu(\sigma)}{d\sigma}=\frac{\theta^{\mu\nu}p_\nu}{2\pi}$ interact with each other by exchanging closed-string tachyon with momentum p_μ.

A factorization of one-loop amplitudes of non-commutative scalar field theory is considered in [32][38] and claimed there to be realized at the low-energy of this field theory by using the path-ordered exponentials. The procedure of the zero-slope limit adopted in those papers are quite different form that of the present paper. Our limiting procedure is not the conventional one to obtain the field theory limit of open-string. In fact the present limit would reduce to the field theory limit of closed-string if the B field were vanishing. We would like to emphasize that the existence of the B field makes the limit different from the conventional field theory limit of closed-string.

5 Coordinate and Momentum Eigenstates of Closed-String

We make a digression from the previous discussion on the world-volume theory. In this section we introduce coordinate and momentum eigenstates of closed-string and make some observation on their relation with the boundary states. These eigenstates will play important roles in the subsequent discussions.

We begin with a simple remark on closed-string momentum currents. These currents are introduced by taking functional derivatives of the action : $P_M(\sigma,\tau)=i\frac{\delta S[X]}{\delta\partial_\tau X^M(\sigma,\tau)}$. We have two expressions of the action, (2.2) and (2.3). They provide two momentum currents, which coincide for the Dirichlet directions but become slightly different from each other for the Neumann directions. When the action (2.3) is used one obtains

$$P_\mu(\sigma,\tau) \equiv \frac{i}{2\pi\alpha'} g_{\mu\nu}\partial_\tau X^\nu(\sigma,\tau) \ , \tag{5.1}$$

while the action (2.2) leads to

$$P_\mu^{(B)}(\sigma,\tau) \equiv \frac{i}{2\pi\alpha'}\Big(g_{\mu\nu}\partial_\tau X^\nu(\sigma,\tau) - i2\pi\alpha' B_{\mu\nu}\partial_\sigma X^\nu(\sigma,\tau)\Big) \ . \tag{5.2}$$

Their conserved charges become the same, that is, momentum p_μ, since these are integrals of the currents over the circle. Whichever momentum current one adopts as the conjugate variable of $X^\mu(\sigma,\tau)$, canonical quantization of closed-string leads to the same commutation relations (2.5).

Let $\hat{X}^M(\sigma)$ be closed-string coordinate operators. They are simply given by $\hat{X}^M(\sigma) \equiv X^M(\sigma,0)$. The conjugate momentum operators $\hat{P}_M(\sigma)$ are those operators satisfying $\Big[\hat{X}^M(\sigma), \hat{P}_N(\sigma')\Big] = i\delta^M_N\delta(\sigma-\sigma')$ [4]. These are given by the above momentum currents. Due to the existence of two different currents we have two operators, $\hat{P}_M(\sigma) \equiv P_M(\sigma,0)$ and $\hat{P}^{(B)}_M(\sigma) \equiv P^{(B)}_M(\sigma,0)$. All these operators are periodic with respect to σ. We may provide their mode expansions similar to Eq.(2.4). For the later discussion these turn out to be inconvenient. Instead we use the following mode expansions :

$$\begin{aligned}
\hat{X}^M(\sigma) &= \hat{x}_0^M + \frac{1}{\sqrt{2}}\sum_{n=1}^{\infty}\left(\hat{\chi}_n^M e^{-in\sigma} + \hat{\chi}_n^{\dagger M} e^{in\sigma}\right) \ , \\
\hat{P}_M(\sigma) &= \frac{1}{2\pi}\left[\hat{p}_{0M} + \frac{1}{\sqrt{2}}\sum_{n=1}^{\infty}\left(\hat{\psi}_{nM} e^{-in\sigma} + \hat{\psi}^\dagger_{nM} e^{in\sigma}\right)\right] \ , \\
\hat{P}^{(B)}_M(\sigma) &= \frac{1}{2\pi}\left[\hat{p}_{0M} + \frac{1}{\sqrt{2}}\sum_{n=1}^{\infty}\left(\hat{\varrho}_{nM} e^{-in\sigma} + \hat{\varrho}^\dagger_{nM} e^{in\sigma}\right)\right] \ ,
\end{aligned} \tag{5.3}$$

where $\hat{\chi}_n^{\dagger M}$, $\hat{\psi}^\dagger_{nM}$ and $\hat{\varrho}^\dagger_{nM}$ are the hermitian conjugates of the corresponding ones. Their commutation relations can be read as follows :

$$\begin{aligned}
&[\hat{x}_0^M, \hat{p}_{0N}] = i\delta^M_N \ , \quad [\hat{\chi}_m^M, \hat{\psi}^\dagger_{nN}] = [\hat{\chi}_m^{\dagger M}, \hat{\psi}_{nN}] = 2i\delta^M_N\delta_{m,n} \ , \\
&[\hat{\chi}_m^M, \hat{\varrho}^\dagger_{nN}] = [\hat{\chi}_m^{\dagger M}, \hat{\varrho}_{nN}] = 2i\delta^M_N\delta_{m,n} \ , \quad \text{otherwise} = 0 \ .
\end{aligned} \tag{5.4}$$

---

[4] The delta function on the circle is given by

$$\delta(\sigma) = \frac{1}{2\pi}\sum_{n\in\mathbb{Z}} e^{in\sigma} \qquad (0 \le \sigma < 2\pi) \ .$$

These oscillator modes are linear sums of the standard oscillator modes of $X^M(\sigma,\tau)$. For the reader's convenience we attach below the dictionary :

$$\begin{aligned}
\hat{\chi}_n^M &= i\frac{\sqrt{\alpha'}}{n}\left(\alpha_n^M - \tilde{\alpha}_{-n}^M\right) \quad , \qquad \hat{\chi}_n^{\dagger M} = -i\frac{\sqrt{\alpha'}}{n}\left(\alpha_{-n}^M - \tilde{\alpha}_n^M\right) \ , \\
\hat{\psi}_{nM} &= \frac{1}{\sqrt{\alpha'}}g_{MN}\left(\alpha_n^N + \tilde{\alpha}_{-n}^N\right) \quad , \qquad \hat{\psi}_{nM}^\dagger = \frac{1}{\sqrt{\alpha'}}g_{MN}\left(\alpha_{-n}^N + \tilde{\alpha}_n^N\right) \ , \\
\hat{\varrho}_{nM} &= \frac{1}{\sqrt{\alpha'}}\left(E_{MN}\alpha_n^N + E_{MN}^T\tilde{\alpha}_{-n}^N\right) \quad , \qquad \hat{\varrho}_{nM}^\dagger = \frac{1}{\sqrt{\alpha'}}\left(E_{MN}\alpha_{-n}^N + E_{MN}^T\tilde{\alpha}_n^N\right) \ ,
\end{aligned} \tag{5.5}$$

where the (i,j)-components of E are understood as g_{ij}. Therefore we have $\hat{\psi}_{ni} = \hat{\varrho}_{ni}$ and $\hat{\psi}_{ni}^\dagger = \hat{\varrho}_{ni}^\dagger$.

5.1 Coordinate and momentum eigenstates

Coordinate and momentum eigenstates are respectively eigenstates of $\hat{X}(\sigma)$ and $\hat{P}(\sigma)$ or $\hat{P}^{(B)}(\sigma)$. We describe them in some detail. These are closed-string extensions of those obtained [40] for open-string.

Coordinate eigenstates

We first describe the coordinate eigenstates. They are characterized by two conditions ; (i) they are eigenstates of the string coordinate operators,

$$\hat{X}^M(\sigma)|X\rangle = X^M(\sigma)|X\rangle, \tag{5.6}$$

and (ii) they satisfy the orthonormality condition,

$$\langle X|X'\rangle = \prod_{0\le\sigma<2\pi}\delta(X(\sigma) - X'(\sigma)), \tag{5.7}$$

where we abbreviate the superscript M in the RHS. Dual states $\langle X|$ are defined by the hermitian conjugation. It is done by simultaneous operations of the BPZ and complex conjugations.

The above conditions determine the eigenstates. We factorize them into products of the Neumann and the Dirichlet sectors :

$$|X\rangle = |X_N\rangle \otimes |X_D\rangle, \tag{5.8}$$

where $|X_{N,D}\rangle$ are the coordinate eigenstates of the corresponding directions. Let us provide oscillator representations of these states. Taking account of the mode expansions (5.3) we parametrize the eigenvalues in terms of complex variables χ_n^M ($n \in \mathbb{Z}_{\geq 1}$) besides the zero-modes x_0^M. We put

$$X^M(\sigma) = x_0^M + \frac{1}{\sqrt{2}} \sum_{n=1}^{\infty} \left(\chi_n^M e^{-in\sigma} + \overline{\chi}_n^M e^{in\sigma} \right) \ . \tag{5.9}$$

The eigenstates are given by the following infinite products :

$$|X_N\rangle = |x_{0N}\rangle \otimes \prod_{n=1}^{\infty} |\chi_n, \bar{\chi}_n\rangle_N \ , \quad |X_D\rangle = |x_{0D}\rangle \otimes \prod_{n=1}^{\infty} |\chi_n, \bar{\chi}_n\rangle_D \, . \tag{5.10}$$

Here $|x_{0N}\rangle = |x_{0N} = (x_0^\mu)\rangle$ (resp. $|x_{0D}\rangle$) are the normalized eigenstates of the zero modes $\hat{x}_0^\mu$ (resp. $\hat{x}_0^i$) with eigenvalues $x_0^\mu \in \mathbb{R}^{p+1}$ (resp. x_0^i), and their normalization conditions are

$$\langle x'_{0N} | x_{0N} \rangle = \prod_{\mu=0}^{p} \delta \left(x_0'^\mu - x_0^\mu \right) \ , \quad \langle x'_{0D} | x_{0D} \rangle = \prod_{i=p+1}^{D-1} \delta \left(x_0'^i - x_0^i \right) \ . \tag{5.11}$$

$|\chi_n, \bar{\chi}_n\rangle_N$ (resp. $|\chi_n, \bar{\chi}_n\rangle_D$) are the normalized eigenstates of $(\hat{\chi}_n^\mu, \hat{\chi}_n^{\dagger\mu})$ (resp. $(\hat{\chi}_n^i, \hat{\chi}_n^{\dagger i})$) with eigenvalues $(\chi_n^\mu, \bar{\chi}_n^\mu)$ (resp. $(\chi_n^i, \bar{\chi}_n^i)$). Their normalization conditions are chosen as

$$\begin{aligned} {}_N\langle \chi'_n, \bar{\chi}'_n | \, \chi_n, \bar{\chi}_n \rangle_N &= \prod_{\mu=0}^{p} \delta \left(\mathrm{Re}\chi_n'^\mu - \mathrm{Re}\chi_n^\mu \right) \delta \left(\mathrm{Im}\chi_n'^\mu - \mathrm{Im}\chi_n^\mu \right) \ , \\ {}_D\langle \chi'_n, \bar{\chi}'_n | \, \chi_n, \bar{\chi}_n \rangle_D &= \prod_{i=p+1}^{D-1} \delta \left(\mathrm{Re}\chi_n'^i - \mathrm{Re}\chi_n^i \right) \delta \left(\mathrm{Im}\chi_n'^i - \mathrm{Im}\chi_n^i \right) \ . \end{aligned} \tag{5.12}$$

The orthonormality (5.7) follows from the above conditions imposed on each eigenstates :

$$\begin{aligned} \langle X | X' \rangle &= \langle x_0 | x'_0 \rangle \prod_{n=1}^{\infty} \langle \chi_n, \bar{\chi}_n | \chi'_n, \bar{\chi}'_n \rangle \\ &= \delta(x_0 - x'_0) \prod_{n=1}^{\infty} \delta(\mathrm{Re}\chi_n - \mathrm{Re}\chi'_n) \delta(\mathrm{Im}\chi_n - \mathrm{Im}\chi'_n) \\ &= \prod_{0 \leq \sigma < 2\pi} \delta(X(\sigma) - X'(\sigma)), \end{aligned} \tag{5.13}$$

where we put $|x_0\rangle \equiv |x_{0N}\rangle \otimes |x_{0D}\rangle$ and $|\chi_n, \bar{\chi}_n\rangle \equiv |\chi_n, \bar{\chi}_n\rangle_N \otimes |\chi_n, \bar{\chi}_n\rangle_D$.

The steps to obtain oscillator realizations of eigenstates $|\chi_n, \bar{\chi}_n\rangle_N$ and $|\chi_n, \bar{\chi}_n\rangle_D$ are presented in Appendix C [5]. They turn out to be as follows :

$$
\begin{aligned}
|\chi_n, \bar{\chi}_n\rangle_N &= \left(\frac{n}{\pi\alpha'}\right)^{\frac{p+1}{2}} \sqrt{-\det g_{\mu\nu}} \\
&\times \exp\left[\frac{1}{n}\alpha^{\mu}_{-n} g_{\mu\nu}\tilde{\alpha}^{\nu}_{-n} - \frac{i}{\sqrt{\alpha'}}\left(\chi^{\mu}_n g_{\mu\nu}\alpha^{\nu}_{-n} + \bar{\chi}^{\mu}_n g_{\mu\nu}\tilde{\alpha}^{\nu}_{-n}\right) - \frac{n}{2\alpha'}\bar{\chi}^{\mu}_n g_{\mu\nu}\chi^{\nu}_n\right]|0\rangle \ , \qquad (5.14) \\
|\chi_n, \bar{\chi}_n\rangle_D &= \left(\frac{n}{\pi\alpha'}\right)^{\frac{d}{2}} \sqrt{\det g_{ij}} \\
&\times \exp\left[\frac{1}{n}\alpha^{i}_{-n} g_{ij}\tilde{\alpha}^{\nu}_{j} - \frac{i}{\sqrt{\alpha'}}\left(\chi^{i}_n g_{ij}\alpha^{i}_{-n} + \bar{\chi}^{i}_n g_{ij}\tilde{\alpha}^{j}_{-n}\right) - \frac{n}{2\alpha'}\bar{\chi}^{i}_n g_{ij}\chi^{j}_n\right]|0\rangle \ . \qquad (5.15)
\end{aligned}
$$

The coordinate eigenstates provide a complete basis of the Hilbert space $\mathcal{H}^X_c = \mathcal{H}^X_{cN} \otimes \mathcal{H}^X_{cD}$, where $\mathcal{H}^X_{cN}$ and $\mathcal{H}^X_{cD}$ stand for the Neumann and the Dirichlet sectors. We have the following completeness relation :

$$
\begin{aligned}
1 &= \int [dX] \ |X\rangle\langle X| \\
&= \int d^D x_0 \prod_{n=1}^{\infty}\left(\prod_{M=0}^{D-1}\frac{d\bar{\chi}^M_n d\chi^M_n}{2i}\right)|\chi_n, \bar{\chi}_n\rangle \, |x_0\rangle\langle x_0| \, \langle\chi_n, \bar{\chi}_n| \ . \qquad (5.16)
\end{aligned}
$$

Momentum eigenstates

We start with eigenstates of $\hat{P}(\sigma)$. They are characterized by two conditions ; (i) they are eigenstates of $\hat{P}(\sigma)$,

$$\hat{P}_M(\sigma)|P\rangle = P_M(\sigma)|P\rangle, \tag{5.17}$$

and (ii) they satisfy the orthonormality condition,

$$\langle P|P'\rangle = \prod_{0\le\sigma<2\pi}\delta(P(\sigma) - P'(\sigma)). \tag{5.18}$$

These conditions determine the eigenstates. We factorize them into products of the Neumann and the Dirichlet sectors.

$$|P\rangle = |P_N\rangle \otimes |P_D\rangle. \tag{5.19}$$

---

[5] The following dictionary might be useful for the correspondence :

$$
\begin{aligned}
&\hat{\chi}^M_n = \hat{\phi}^{(\mathrm{I})M}_n + i\hat{\phi}^{(\mathrm{II})M}_n \ , \quad \hat{\psi}_{nM} = \hat{\pi}^{(\mathrm{I})}_{nM} + i\hat{\pi}^{(\mathrm{II})}_{nM} \ , \quad \hat{\varrho}_{nM} = \hat{\varpi}^{(\mathrm{I})}_{nM} + i\hat{\varpi}^{(\mathrm{II})}_{nM} \ , \\
&\hat{\chi}^{\dagger M}_n = \hat{\phi}^{(\mathrm{I})M}_n - i\hat{\phi}^{(\mathrm{II})M}_n \ , \quad \hat{\psi}^{\dagger}_{nM} = \hat{\pi}^{(\mathrm{I})}_{nM} - i\hat{\pi}^{(\mathrm{II})}_{nM} \ , \quad \hat{\varrho}^{\dagger}_{nM} = \hat{\varpi}^{(\mathrm{I})}_{nM} - i\hat{\varpi}^{(\mathrm{II})}_{nM} \ .
\end{aligned}
$$

Let us concentrate on the Neumann directions. Construction of eigenstates of the Dirichlet sector is parallel. We parametrize the eigenvalues in terms of complex variables $\psi_{n\mu}$ ($n \in \mathbb{Z}_{\geq 1}$) besides the zero-modes $p_{0\mu}$. We put

$$P_\mu(\sigma) \;=\; \frac{1}{2\pi}\left[p_{0\mu} + \frac{1}{\sqrt{2}}\sum_{n=1}^{\infty}\left(\psi_{n\mu}e^{-in\sigma} + \overline{\psi}_{n\mu}e^{in\sigma}\right)\right] . \tag{5.20}$$

The eigenstates are given by the following infinite products :

$$|P_N\rangle = |p_{0N}\rangle \otimes \prod_{n=1}^{\infty} |\psi_n, \overline{\psi}_n\rangle_N, \tag{5.21}$$

where $|\psi_n, \overline{\psi}_n\rangle_N$ denote the normalized eigenstates of $(\hat{\psi}_{n\mu}, \hat{\psi}^\dagger_{n\mu})$ with eigenvalues $(\psi_{n\mu}, \bar{\psi}_{n\mu})$, and the normalization conditions are taken as

$$_N\langle\psi'_n, \bar{\psi}'_n|\psi_n, \bar{\psi}_n\rangle_N = \prod_{\mu=0}^{p} \delta\left(\mathrm{Re}\psi'_{n\mu} - \mathrm{Re}\psi_{n\mu}\right)\delta\left(\mathrm{Im}\psi'_{n\mu} - \mathrm{Im}\psi_{n\mu}\right) . \tag{5.22}$$

These conditions ensure the orthonormality of $|P_N\rangle$. Oscillator realizations of $|\psi_n, \bar{\psi}_n\rangle_N$ are given in appendix C. They turn out to be as follows :

$$\begin{aligned} \left|\psi_n, \bar{\psi}_n\right\rangle_N = &\left(\frac{\alpha'}{\pi n}\right)^{\frac{p+1}{2}} \frac{1}{\sqrt{-\det g_{\mu\nu}}} \\ &\times \exp\left[-\frac{1}{n}\alpha^\mu_{-n}g_{\mu\nu}\tilde{\alpha}^\nu_{-n} + \frac{\sqrt{\alpha'}}{n}\left(\psi_{n\mu}\alpha^\mu_{-n} + \bar{\psi}_{n\mu}\tilde{\alpha}^\mu_{-n}\right) - \frac{\alpha'}{2n}\bar{\psi}_{n\mu}g^{\mu\nu}\psi_{n\nu}\right]|0\rangle . \end{aligned} \tag{5.23}$$

Similarly to $|X_N\rangle$, momentum eigenstates $|P_N\rangle$ provide another complete basis of $\mathcal{H}^X_{cN}$. The completeness relation is

$$\begin{aligned} 1 \;&=\; \int [dP_N]\;|P_N\rangle\langle P_N| \\ &=\; \int d^{p+1}p_0 \prod_{n=1}^{\infty}\left(\prod_{\mu=0}^{p}\frac{d\bar{\psi}_{n\mu}d\psi_{n\mu}}{2i}\right)\left|\psi_n, \bar{\psi}_n\right\rangle_N |p_{0N}\rangle\langle p_{0N}|_N\left\langle\psi_n, \bar{\psi}_n\right| . \end{aligned} \tag{5.24}$$

Next we describe eigenstates of $\hat{P}^{(B)}(\sigma)$. They are determined by similar conditions to those imposed on $\hat{P}(\sigma)$ and are factorized into products of the Neumann and the Dirichlet sectors :

$$|P^{(B)}\rangle = |P^{(B)}_N\rangle \otimes |P_D\rangle. \tag{5.25}$$

We also concentrate on the Neumann directions. Let us parametrize the eigenvalues $P^{(B)}_\mu(\sigma)$ by complex variables $\varrho_{n\mu}$ ($n \in \mathbb{Z}_{\geq 1}$) and the zero-modes $p_{0\mu}$ as

$$P^{(B)}_\mu(\sigma) = \frac{1}{2\pi}\left[p_{0\mu} + \frac{1}{\sqrt{2}}\sum_{n=1}^{\infty}\left(\varrho_{n\mu}e^{-in\sigma} + \overline{\varrho}_{n\mu}e^{in\sigma}\right)\right] . \tag{5.26}$$

The eigenstates are given by the following infinite products :

$$|P_N^{(B)}\rangle = |p_{0N}\rangle \otimes \prod_{n=1}^{\infty} |\varrho_n, \bar{\varrho}_n\rangle_N, \tag{5.27}$$

where $|\varrho_n, \bar{\varrho}_n\rangle_N$ are the normalized eigenstates of $(\hat{\varrho}_{n\mu}, \hat{\varrho}^\dagger_{n\mu})$ with eigenvalues $(\varrho_{n\mu}, \bar{\varrho}_{n\mu})$. We normalize them by

$${}_N\langle \varrho'_n, \bar{\varrho}'_n|\varrho_n, \bar{\varrho}_n\rangle_N = \prod_{\mu=0}^{p} \delta(\mathrm{Re}\varrho'_{n\mu} - \mathrm{Re}\varrho_{n\mu})\delta(\mathrm{Im}\varrho'_{n\mu} - \mathrm{Im}\varrho_{n\mu}) \ . \tag{5.28}$$

The orthonormality of $|P_N^{(B)}\rangle$ follows from these conditions. Oscillator realizations of these states are given in the appendix. They turn out to be as follows :

$$\begin{aligned} |\varrho_n, \bar{\varrho}_n\rangle_N = {}& \left(\frac{\alpha'}{n\pi}\right)^{\frac{p+1}{2}} \sqrt{-\det\left(\frac{1}{E^T} g \frac{1}{E}\right)^{\mu\nu}} \\ & \times \exp\left[-\frac{1}{n}\alpha^\mu_{-n} N_{\mu\nu}\tilde{\alpha}^\nu_{-n} + \frac{\sqrt{\alpha'}}{n}\left\{\varrho_{n\mu}\left(\frac{1}{E^T}g\right)^\mu_{\ \nu}\alpha^\nu_{-n} + \bar{\varrho}_{n\mu}\left(\frac{1}{E}g\right)^\mu_{\ \nu}\tilde{\alpha}^\nu_{-n}\right\}\right. \\ & \left. \qquad -\frac{\alpha'}{2n}\bar{\varrho}_{n\mu}\left(\frac{1}{E^T}g\frac{1}{E}\right)^{\mu\nu}\varrho_{n\nu}\right]|0\rangle \ . \end{aligned} \tag{5.29}$$

Momentum eigenstates $|P_N^{(B)}\rangle$ provide a complete basis of $\mathcal{H}^X_{cN}$ as well. The completeness relation reads as follows :

$$\begin{aligned} 1 &= \int \left[dP_N^{(B)}\right] \left|P_N^{(B)}\right\rangle \left\langle P_N^{(B)}\right| \\ &= \int d^{p+1}p_0 \prod_{n=1}^{\infty}\left(\prod_{\mu=0}^{p}\frac{d\bar{\varrho}_{n\mu}d\varrho_{n\mu}}{2i}\right)|\varrho_n, \bar{\varrho}_n\rangle_N\,|p_{0N}\rangle\langle p_{0N}|_N\langle\varrho_n, \bar{\varrho}_n| \ . \end{aligned} \tag{5.30}$$

5.2 Some Observations

The previous constructions give all the eigenstates in terms of infinite products of the correctly normalized eigenstates of each massive modes. For instance the eigenstates $|X_N\rangle$ and $|X_D\rangle$ in (5.10) are given by the infinite products of $|\chi_n, \overline{\chi}_n\rangle_N$ and $|\chi_n, \overline{\chi}_n\rangle_D$. One can find in Eqs.(5.14) and (5.15) that the normalization factors are respectively $\left(\frac{n}{\pi\alpha'}\right)^{\frac{p+1}{2}}\sqrt{-\mathrm{det}g_{\mu\nu}}$ and $\left(\frac{n}{\pi\alpha'}\right)^{\frac{d}{2}}\sqrt{\mathrm{det}g_{ij}}$. Infinite products of these constants become the normalization factors of $|X_N\rangle$ and $|X_D\rangle$. Let us denote them by $\mathcal{C}_X^{(N)} \equiv \prod_{n=1}^{\infty}\left[n^{\frac{p+1}{2}}\left(\frac{-\det g_{\mu\nu}}{(\pi\alpha')^{p+1}}\right)^{\frac{1}{2}}\right]$ and $\mathcal{C}_X^{(D)} \equiv$

$\prod_{n=1}^{\infty}\left[n^{\frac{d}{2}}\left(\frac{\det g_{ij}}{(\pi\alpha')^{d}}\right)^{\frac{1}{2}}\right]$. The eigenstates can be written as follows :

$$\begin{aligned}
&|X_N\rangle \\
&= \mathcal{C}_X^{(N)} \prod_{n=1}^{\infty} \exp\left[\frac{1}{n}\alpha_{-n}^{\mu} g_{\mu\nu}\tilde{\alpha}_{-n}^{\nu} - \frac{i}{\sqrt{\alpha'}}\left(\chi_n^{\mu} g_{\mu\nu}\alpha_{-n}^{\nu} + \bar{\chi}_n^{\mu} g_{\mu\nu}\tilde{\alpha}_{-n}^{\nu}\right) - \frac{n}{2\alpha'}\bar{\chi}_n^{\mu} g_{\mu\nu}\chi_n^{\nu}\right] |x_{0N}\rangle \ , \\
&|X_D\rangle \\
&= \mathcal{C}_X^{(D)} \prod_{n=1}^{\infty} \exp\left[\frac{1}{n}\alpha_{-n}^{i} g_{ij}\tilde{\alpha}_{-n}^{j} - \frac{i}{\sqrt{\alpha'}}\left(\chi_n^{i} g_{ij}\alpha_{-n}^{j} + \bar{\chi}_n^{i} g_{ij}\tilde{\alpha}_{-n}^{j}\right) - \frac{n}{2\alpha'}\bar{\chi}_n^{i} g_{ij}\chi_n^{j}\right] |x_{0D}\rangle \ ,
\end{aligned} \tag{5.31}$$

As is performed in [30], one may evaluate $\mathcal{C}_X^{(N,D)}$ by using the zeta-function regularization. This regularization scheme leads the following identities [6]:

$$\prod_{n=1}^{\infty} a = a^{\zeta(0)} = a^{-\frac{1}{2}} \ , \quad \prod_{n=1}^{\infty} n^{\alpha} = \exp\left(-\alpha\zeta'(0)\right) = (2\pi)^{\frac{\alpha}{2}} \ , \tag{5.32}$$

and thus we obtain

$$\mathcal{C}_X^{(N)} = \left(\frac{(2\pi^2\alpha')^{p+1}}{-\det g_{\mu\nu}}\right)^{\frac{1}{4}} \ , \quad \mathcal{C}_X^{(D)} = \left(\frac{(2\pi^2\alpha')^{d}}{\det g_{ij}}\right)^{\frac{1}{4}} \ . \tag{5.33}$$

It is important to recall that the boundary state $|B_D\rangle$ is an eigenstate of $\hat{X}^i(\sigma)$ with eigenvalues $X^i(\sigma) = x_0^i$. We have $|B_D\rangle \propto |X_D = x_{0D}\rangle$. If one uses the zeta-function regularization and adopts the above $\mathcal{C}_X^{(D)}$ as the normalization constant, these two states become precisely identical :

$$|B_D\rangle = |X_D = x_{0D}\rangle \ . \tag{5.34}$$

Let us also recall that in quantum mechanics of a single particle, coordinate eigenstates $|x\rangle$ are described by $|x\rangle = e^{-ix\hat{p}}|x=0\rangle$, where $\hat{p}$ is the momentum operator. One may find an analogous realization for the string coordinate eigenstates. Taking account of Eq.(5.34) one can infer the following one :

$$|X_D\rangle =: \exp\left(-i\int_0^{2\pi} d\sigma X^i(\sigma)\hat{P}_i(\sigma)\right) : |B_D; x_{0D}=0\rangle \ , \tag{5.35}$$

where $|B_D; x_{0D}=0\rangle$ is the state (5.34) with $x_0^i = 0$. This turns out to be the case. Using the previous parametrization of the eigenvalues one can find

$$\int_0^{2\pi} d\sigma X^i(\sigma)\hat{P}_i(\sigma) = x_0^i\hat{p}_0^i + \frac{g_{ij}}{2\sqrt{\alpha'}}\sum_{n=1}^{\infty}\left\{\left(\chi_n^i\alpha_{-n}^j + \bar{\chi}_n^i\tilde{\alpha}_{-n}^j\right) + \left(\bar{\chi}_n^i\alpha_n^j + \chi_n^i\tilde{\alpha}_n^j\right)\right\} \ . \tag{5.36}$$

---

[6] $\zeta(s) = \sum_{n=1}^{\infty}\frac{1}{n^s}$; $\sum_{n=1}^{\infty} 1 = \zeta(0) = -\frac{1}{2}$, $\sum_{n=1}^{\infty}(-\ln n) = \zeta'(0) = -\frac{1}{2}\ln(2\pi)$ etc.

It is easy to see that the normal ordered exponential in the RHS of Eq.(5.35) reproduces $|X_D\rangle$. The formula (B.1) may be useful.

The above discussion is also applicable to the string momentum eigenstates. We can write these states as follows :

$$
\begin{aligned}
&|P_N\rangle \\
&= \mathcal{C}_{P_N} \prod_{n=1}^{\infty} \exp\left[-\frac{1}{n}\alpha^{\mu}_{-n} g_{\mu\nu} \tilde{\alpha}^{\nu}_{-n} + \frac{\sqrt{\alpha'}}{n}\left(\psi_{n\mu}\alpha^{\mu}_{-n} + \bar{\psi}_{n\mu}\tilde{\alpha}^{\mu}_{-n}\right) - \frac{\alpha'}{2n}\bar{\psi}_{n\mu} g^{\mu\nu}\psi_{n\nu} \right] |p_{0N}\rangle \ , \\
&\left|P_N^{(B)}\right\rangle \\
&= \mathcal{C}_{P_N^{(B)}} \prod_{n=1}^{\infty} \exp\Bigg[-\frac{1}{n}\alpha^{\mu}_{-n} N_{\mu\nu} \tilde{\alpha}^{\nu}_{-n} + \frac{\sqrt{\alpha'}}{n}\left\{ \varrho_{n\mu}\left(\frac{1}{E^T}g\right)^{\mu}{}_{\nu}\alpha^{\nu}_{-n} + \bar{\varrho}_{n\mu}\left(\frac{1}{E}g\right)^{\mu}{}_{\nu}\tilde{\alpha}^{\nu}_{-n} \right\} \\
&\qquad\qquad\qquad - \frac{\alpha'}{2n}\bar{\varrho}_{n\mu}\left(\frac{1}{E^T}g\frac{1}{E}\right)^{\mu\nu}\varrho_{n\nu} \Bigg] |p_{0N}\rangle \ ,
\end{aligned}
\tag{5.37}
$$

where the normalization factors $\mathcal{C}_{P_N}$ and $\mathcal{C}_{P_N^{(B)}}$ are originally given by the infinite products as follow from Eqs.(5.23) and (5.29). They are regularized to

$$
\mathcal{C}_{P_N} = \left(\frac{-\det g_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} \ , \quad \mathcal{C}_{P_N^{(B)}} = \left(\frac{\det^2 E_{\mu\nu}}{(2\alpha')^{p+1}(-\det g_{\mu\nu})} \right)^{\frac{1}{4}} \ . \tag{5.38}
$$

Boundary state $|B_N\rangle$ is an eigenstate of $\hat{P}^{(B)}_{\mu}(\sigma)$ with vanishing eigenvalue. We have $|B_N\rangle \propto |P^{(B)} = 0\rangle$. If we use the above $\mathcal{C}_{P_N^{(B)}}$ as the normalization constant, these two states become identical including their normalizations :

$$
|B_N\rangle = \left| P_N^{(B)} = 0 \right\rangle \ . \tag{5.39}
$$

Momentum eigenstates can be realized in the quantum mechanics by $|p\rangle = e^{ip\hat{x}}|p = 0\rangle$, where $\hat{x}$ is the position operator. We can find an analogous realization of the string momentum eigenstates. It turns out to be as follows :

$$
\begin{aligned}
\left|P_N^{(B)}\right\rangle \ &= \ \prod_{n=1}^{\infty} \exp\left[-\frac{\alpha'}{4n}\bar{\varrho}_{n\mu}\left\{ g^{-1}\left(g - \frac{1}{2}\left(N - N^T\right)\right) g^{-1} \right\}^{\mu\nu} \varrho_{n\nu} \right] \\
&\quad \times : \exp\left(i\int_0^{2\pi} d\sigma P^{(B)}_{\mu}(\sigma)\hat{X}^{\mu}(\sigma) \right) : |B_N\rangle \ .
\end{aligned}
\tag{5.40}
$$

For the later convenience we provide a similar observation for the eigenstates $|P_N\rangle$ as well. Let $|B_N\rangle_{B=0}$ be the state which is obtained from the Neumann boundary state $|B_N\rangle$ by putting $B_{\mu\nu} = 0$:

$$
|B_N\rangle_{B=0} = \left(\frac{-\det g_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} \prod_{n=1}^{\infty} \exp\left(-\frac{1}{n}\alpha^{\mu}_{-n} g_{\mu\nu}\tilde{\alpha}^{\nu}_{-n} \right) |\mathbf{0}\rangle \ . \tag{5.41}
$$

This is the Neumann boundary state in a vanishing B field background. One can readily find that this is an eigenstate of $\hat{P}_\mu(\sigma)$ with zero eigenvalue: $\hat{P}_\mu(\sigma)\left|B_N\right\rangle_{B=0} = 0$ and hence $\left|B_N\right\rangle_{B=0} \propto |P_N = 0\rangle$. Comparing definitions (5.37) and (5.41), we find that these states coincide with each other including their normalizations, if we adopt C_{P_N} in Eq.(5.38) as the normalization of the momentum eigenstates.

Concerning the state $|P_N\rangle$, a similar formula to Eq.(5.40) becomes

$$|P_N\rangle = \prod_{n=1}^{\infty} \exp\left[-\frac{\alpha'}{4n}\bar{\psi}_{n\mu} g^{\mu\nu} \psi_{n\nu}\right] : \exp\left(i\int_0^{2\pi} d\sigma P_\mu(\sigma)\hat{X}^\mu(\sigma)\right) : \left|B_N\right\rangle_{B=0} . \tag{5.42}$$

This has the same form as Eq.(5.40) with putting $B_{\mu\nu} = 0$.

6 Open Wilson Lines in Closed-String Theory (I)

At the zero-slope limit, which is introduced in Eq.(4.7) and its below in order to capture the UV behavior of the world-volume theory, the generating function of one-loop amplitudes of open-string tachyons is shown to exhibit the factorized form (4.17). It is expressed as a sum of products of two Wilson lines (strictly speaking, their analogues) along the same straight lines $x_0(\sigma)$, multiplying the propagators of closed-string tachyon. This shows that closed-string tachyon T has a tadpole interaction with the open Wilson line. It can be written in a form, $\int dx T(x)\mathcal{P}_\star\left[e^{\int_0^{2\pi} d\sigma\phi(x_0(\sigma))}\right]$. Its origin in string theory can be found in Eq.(4.8). The interaction simply comes from the closed-string tachyon modes of boundary states, $\left\langle K \left| B_N; (\sigma_1, k^{(1)}), \right.\ \cdots, (\sigma_M, k^{(M)})\right\rangle$.

Actually these boundary states have all the components of perturbative closed-string states. At the level of string amplitudes all of them propagate between the boundary states and contribute to the amplitudes. It is shown in [23] that the straight open Wilson line can couple with the on-shell graviton and, as will be seen in the later section, it can be generalized to the off-shell. This indicates that all the perturbative closed-string states have tadpole interactions with the open Wilson line. Therefore we may unfasten the zero-slope limit (4.17) so that propagations of all these states are made manifest. In this section we pursue such a possibility. We also restrict to the case of 25-brane in the critical dimensions.

Let us provide a general perspective on this issue before we start calculations. First of all, it can be expected [27] that closed-string propagations including gravitons fluctuate the straight line appearing in Eq.(4.15) and transform it into curved ones. In other words we can expect that there are correlations between their deviations from the straight line and the propagations of closed-string states. These curves will appear as the corresponding Wilson lines, and we may factorize the generating function into a sum of these products at the zero-slope limit. Analogously to the standard factorization of closed-string amplitudes the sum must be taken originally over the perturbative closed-string states. The above correspondence between curves and states will enable us to translate the sum as an integral over the space of curves. This integral may be suppressed by a suitable weight as the straight line is suppressed in Eq.(4.17) by the closed-string tachyon propagator. It is amazing that one can interpret the tachyon propagator as a propagator of the straight open Wilson line by notifying $p_\mu g^{\mu\nu} p_\nu \approx \alpha'^{-2} \dot{x}_0^\mu G_{\mu\nu} \dot{x}_0^\nu$. The factorization (4.17) may be obtained from the aforementioned integral by integrating out the fluctuations. We may say that the straight open Wilson line is the average.

6.1 Factorization by closed-string momentum eigenstates

In order to justify the above perspective let us first factorize the string amplitudes by an insertion of a partition of unity. Use of that constructed from closed-string momentum eigenstates turns out to be relevant. It is given in Eq.(5.24) as $1 = \int [dP_N]\, |P_N\rangle \langle P_N|$, where the eigenvalue $P_\mu(\sigma)$ is parametrized by $P_\mu(\sigma) = \frac{1}{2\pi}\left[p_{0\mu} + \frac{1}{\sqrt{2}} \sum_{n=1}^\infty \left(\psi_{n\mu} e^{-in\sigma} + \overline{\psi}_{n\mu} e^{in\sigma}\right)\right]$.

We factorize the amplitudes (3.42) by an insertion of Eq.(5.24) :

$$\begin{aligned}
&\left\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right| q_c^{\frac{1}{2}(L_0+\tilde{L}_0-2)} \left| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \\
&= \int [dP_N] \left\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right| q_c^{\frac{1}{4}(L_0+\tilde{L}_0-2)} \Big| P_N \Big\rangle \\
&\qquad\qquad \times \Big\langle P_N \Big| q_c^{\frac{1}{4}(L_0+\tilde{L}_0-2)} \left| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle . \qquad (6.1)
\end{aligned}$$

Two factors appearing in the above can be calculated by using the oscillator representations. These are given in Eq.(3.31) (and Eq.(3.41)) for the boundary states, and Eq.(5.37) (and its hermitian conjugate) for the momentum eigenstates. Technical difficulties we may meet in the calculations are evaluations of matrix elements of the following type :

$$\begin{aligned}
&\langle 0| \exp\left[-\frac{1}{n} q_c^{n/2} \alpha_n^\mu g_{\mu\nu} \tilde{\alpha}_n^\nu + \frac{\sqrt{\alpha'}}{n} q_c^{n/4} \left(\overline{\psi}_{n\mu} \alpha_n^\mu + \psi_{n\mu} \tilde{\alpha}_n^\mu\right)\right] \\
&\quad \times \exp\left[-\frac{1}{n} \alpha_{-n}^\mu N_{\mu\nu} \tilde{\alpha}_{-n}^\nu + \frac{\sqrt{2\alpha'}}{n} \sum_{r=1}^M k_\mu^{(r)} \left\{ \left(\frac{1}{E^T} g\right)^\mu{}_\nu \alpha_{-n}^\nu e^{in\sigma_r} + \left(\frac{1}{E} g\right)^\mu{}_\nu \tilde{\alpha}_{-n}^\nu e^{-in\sigma_r} \right\}\right] |0\rangle. \\
&\hfill (6.2)
\end{aligned}$$

These can be computed by using Eq.(B.7) in Appendix B. To describe the result it turns out useful to rescale complex variables ψ_n to $2 q_c^{n/4} \psi_n$. After these rescalings the factorization can be written as follows :

$$
\begin{aligned}
&\int [dP_N] \left\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right| q_c^{\frac{1}{4}(L_0+\tilde{L}_0-2)} \Big| P_N \Big\rangle \\
&\qquad\qquad \times \Big\langle P_N \Big| q_c^{\frac{1}{4}(L_0+\tilde{L}_0-2)} \left| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle . \\
&= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{2}} \int dp_0 \delta^{26} \left(p_0 + \sum_{r=M+1}^{M+N} k^{(r)} \right) \delta^{26} \left(p_0 - \sum_{r=1}^{M} k^{(r)} \right) \times q_c^{\frac{\alpha'}{4} p_{0\mu} g^{\mu\nu} p_{0\nu} - 1} \\
&\quad \times \prod_{1 \le r < s \le M} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \times \prod_{M+1 \le r < s \le M+N} e^{-\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \\
&\quad \times \prod_{1 \le r < s \le M} e^{-\frac{i}{2\pi} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} (\sigma_r - \sigma_s)} \times \prod_{M+1 \le r < s \le M+N} e^{\frac{i}{2\pi} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} (\sigma_r - \sigma_s)} \\
&\quad \times \prod_{1 \le r < s \le M} \left| e^{i\sigma_r} - e^{i\sigma_s} \right|^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} \times \prod_{M+1 \le r < s \le M+N} \left| e^{i\sigma_r} - e^{i\sigma_s} \right|^{2\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)}} \\
&\quad \times \prod_{n=1}^{\infty} \exp \left[-\frac{2\alpha' q_c^{\frac{n}{2}}}{n} \left(\sum_{r=1}^{M} e^{-in\sigma_r} k_\mu^{(r)} \right) \left(\frac{1}{E} g \frac{1}{E - q_c^{\frac{n}{2}} E^T} \right)^{\mu\nu} \left(\sum_{r=1}^{M} e^{in\sigma_r} k_\nu^{(r)} \right) \right. \\
&\qquad\qquad\qquad \left. - \frac{2\alpha' q_c^{\frac{n}{2}}}{n} \left(\sum_{r=M+1}^{M+N} e^{in\sigma_r} k_\mu^{(r)} \right) \left(\frac{1}{E} g \frac{1}{E - q_c^{\frac{n}{2}} E^T} \right)^{\mu\nu} \left(\sum_{r=M+1}^{M+N} e^{-in\sigma_r} k_\nu^{(r)} \right) \right] \\
&\quad \times \prod_{n=1}^{\infty} \int \frac{d\overline{\psi}_n d\psi_n}{(2i)^{26}} \left(\frac{\alpha'}{4n\pi q_c^{\frac{n}{2}}} \right)^{26} \frac{-\det g_{\mu\nu}}{\det^2 (g - q_c^{\frac{n}{2}} N)_{\mu\nu}} \\
&\qquad \times \exp \left[-\frac{\alpha'}{4n} \overline{\psi}_{n\mu} \left(\frac{1}{q_c^{\frac{n}{2}} g} + \frac{1}{g - q_c^{\frac{n}{2}} N^T} N^T \frac{1}{g} + \frac{1}{g} N \frac{1}{g - q_c^{\frac{n}{2}} N} \right)^{\mu\nu} \psi_{n\nu} \right] \\
&\qquad \times \exp \left[\frac{\alpha'}{n\sqrt{2}} \psi_{n\mu} \left(\frac{1}{E^T - q_c^{\frac{n}{2}} E} \right)^{\mu\nu} \left(\sum_{r=1}^{M} e^{-in\sigma_r} k_\nu^{(r)} \right) \right. \\
&\qquad\qquad\qquad\qquad \left. + \frac{\alpha'}{n\sqrt{2}} \overline{\psi}_{n\mu} \left(\frac{1}{E - q_c^{\frac{n}{2}} E^T} \right)^{\mu\nu} \left(\sum_{r=1}^{M} e^{in\sigma_r} k_\nu^{(r)} \right) \right] \\
&\qquad \times \exp \left[-\frac{\alpha'}{n\sqrt{2}} \overline{\psi}_{n\mu} \left(\frac{1}{E^T - q_c^{\frac{n}{2}} E} \right)^{\mu\nu} \left(\sum_{r=M+1}^{M+N} e^{in\sigma_r} k_\nu^{(r)} \right) \right. \\
&\qquad\qquad\qquad\qquad \left. - \frac{\alpha'}{n\sqrt{2}} \psi_{n\mu} \left(\frac{1}{E - q_c^{\frac{n}{2}} E^T} \right)^{\mu\nu} \left(\sum_{r=M+1}^{M+N} e^{-in\sigma_r} k_\nu^{(r)} \right) \right] . \qquad (6.3)
\end{aligned}
$$

The above factorization may be compared with the previous expression of the amplitudes. Terms in the first four lines can be found exactly in Eq.(3.45). The other terms describe a factorization of F in the same equation. If we integrate out $\psi_{n\mu}$ and $\overline{\psi}_{n\mu}$ they provide Eq.(3.47). These variables describe fluctuations of $P_\mu(\sigma)$. Corresponding degrees of freedom of closed-string are given by the massive modes. The factorization of F in terms of those complex variables are plausible since F is the sum of contributions of these massive modes.

6.2 Factorization by open Wilson lines

We examine the zero-slope limit of the factorization (6.3). The limit we discuss is same as that investigated previously to capture the UV behavior of the world-volume theory. It is taken by fixing parameters $s^{(c)} = \alpha'|\tau^{(c)}|$ and σ_r besides the open-string tensors. We also keep ψ_n and $\overline{\psi}_n$ intact. They do not scale under the limit. It should be noticed that the complex variables used originally to parametrize $P_\mu(\sigma)$ do scale under the limit. This is because ψ_n and $\overline{\psi}_n$ in (6.3) are the rescaled ones introduced by multiplying the original variables by $2q_c^{n/4}$.

We first focus on the integral over ψ_n and $\overline{\psi}_n$ in Eq.(6.3). Let us start by considering the first exponential in the integral. Scaling part in the exponent is expressed by means of the closed-string tensors and $q_c = e^{2\pi i\tau^{(c)}}$. It has the form :

$$\alpha'\left(\frac{1}{q_c^{\frac{n}{2}}g} + \frac{1}{g - q_c^{\frac{n}{2}}N^T}N^T\frac{1}{g} + \frac{1}{g}N\frac{1}{g - q_c^{\frac{n}{2}}N}\right). \tag{6.4}$$

Dominant contribution at the limit clearly comes from the first term :

$$\alpha'\left(\frac{1}{q_c^{\frac{n}{2}}g} + \frac{1}{g - q_c^{\frac{n}{2}}N^T}N^T\frac{1}{g} + \frac{1}{g}N\frac{1}{g - q_c^{\frac{n}{2}}N}\right) \approx q_c^{-\frac{n}{2}}\alpha' g^{-1} \approx -\frac{\theta G\theta}{4\pi^2 q_c^{\frac{n}{2}}\alpha'}, \tag{6.5}$$

where we use equalities (3.19). We turn to the determinant factors, which is also described by the closed-string tensors. Their contributions can be read as follows :

$$\left(\frac{\alpha'}{4n\pi q_c^{\frac{n}{2}}}\right)^{26}\frac{\det g}{\det^2(g - q_c^{\frac{n}{2}}N)} \approx \left(\frac{1}{4n\pi}\right)^{26}\det\left[q_c^{-\frac{n}{2}}\alpha' g^{-1}\right] \approx \left(\frac{1}{2\pi}\right)^{26}\det\left[-\frac{\theta G\theta}{8\pi^2 n q_c^{\frac{n}{2}}\alpha'}\right]. \tag{6.6}$$

The other two exponentials in the integral have similar forms. It is enough to know the behavior of $\alpha'(E^T - q_c^{\frac{n}{2}}E)^{-1}$ in their exponents. It can be read by using Eq.(3.18) as follows :

$$\frac{\alpha'}{E^T - q_c^{\frac{n}{2}}E} \approx \frac{\alpha'}{E^T} \approx -\frac{\theta}{2\pi}. \tag{6.7}$$

As we studied previously, contributions of the first four lines in the factorization (6.3) are to give rise to the straight open Wilson lines. The last pieces we need to estimate are the exponentials whose exponents are bilinear of $e^{in\sigma_r}k^{(r)}$ with the weights, $\alpha' q_c^{\frac{n}{2}}E^{-1}g(E - q_c^{\frac{n}{2}}E^T)^{-1}$. It is enough to know the behavior of these weights. Again using Eqs. (3.18) and (3.19) it can be read as

$\alpha' q_c^{\frac{n}{2}} E^{-1} g (E - q_c^{\frac{n}{2}} E^T)^{-1} \approx -\alpha' q_c^{\frac{n}{2}} G^{-1} \approx 0$. Thus we can neglect these pieces in the zero-slope limit.

Collecting all these estimations we can obtain the zero-slope limit of the factorization. It turns out to be as follows :

$$\begin{aligned}
&\int [dP_N] \left\langle B_N; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \right| q_c^{\frac{1}{4}(L_0 + \tilde{L}_0 - 2)} \left| P_N \right\rangle \\
&\qquad\qquad \times \left\langle P_N \right| q_c^{\frac{1}{4}(L_0 + \tilde{L}_0 - 2)} \left| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \\
&\approx \int dp_0 \exp\left\{ -\frac{\pi s^{(c)}}{2} \left(p_{0\mu} g^{\mu\nu} p_{0\nu} - \frac{4}{\alpha'} \right) \right\} \\
&\quad \times \prod_{n=1}^{\infty} \int \frac{d\overline{\psi}_n d\psi_n}{(4\pi i)^{26}} \left\{ -\det\left(-\frac{\theta G \theta}{8\pi^2 n q_c^{\frac{n}{2}} \alpha'} \right) \right\} \exp\left\{ \frac{\overline{\psi}_{n\mu} \left(\theta G \theta\right)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n q_c^{\frac{n}{2}} \alpha'} \right\} \\
&\qquad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{4}} \delta^{26}\left(p_0 - \sum_{r=1}^{M} k^{(r)} \right) \prod_{1 \le r < s \le M} e^{\frac{i}{2} k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu \epsilon(\sigma_r - \sigma_s)} \\
&\qquad\quad \times \prod_{r=1}^{M} \exp\left[-\frac{i}{2\pi} k^{(r)}_\mu \theta^{\mu\nu} \left\{ p_{0\nu} \sigma_r + \frac{1}{\sqrt{2}} \sum_{n=1}^{\infty} \frac{i}{n} \left(\psi_{n\nu} e^{-in\sigma_r} - \overline{\psi}_{n\nu} e^{in\sigma_r} \right) \right\} \right] \\
&\qquad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{4}} \delta^{26}\left(p_0 + \sum_{r=M+1}^{M+N} k^{(r)} \right) \prod_{M+1 \le r < s \le M+N} e^{-\frac{i}{2} k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu \epsilon(\sigma_r - \sigma_s)} \\
&\qquad\quad \times \prod_{r=M+1}^{M+N} \exp\left[-\frac{i}{2\pi} k^{(r)}_\mu \theta^{\mu\nu} \left\{ p_{0\nu} \sigma_r + \frac{1}{\sqrt{2}} \sum_{n=1}^{\infty} \frac{i}{n} \left(\psi_{n\nu} e^{-in\sigma_r} - \overline{\psi}_{n\nu} e^{in\sigma_r} \right) \right\} \right] .
\end{aligned} \tag{6.8}$$

In the above $q_c = e^{-\frac{2\pi s^{(c)}}{\alpha'}}$ and the closed-string metric is understood as $g^{\mu\nu} \approx -\frac{(\theta G \theta)^{\mu\nu}}{(2\pi\alpha')^2}$. Both of the integrals with respect to p_0 and ψ_n ($\bar{\psi}_n$) become Gaussian integrals. The Gaussian weight of p_0 is $\pi s^{(c)} g^{-1} \approx -s^{(c)} \frac{\theta G \theta}{4\pi\alpha'^2}$ while the weight of ψ_n is $-\frac{\theta G \theta}{8\pi^2 n q_c^{\frac{n}{2}} \alpha'}$, which is equal to $-s^{(c)} \frac{\theta G \theta}{8\pi\alpha'^2} \times \left(\frac{n\pi s^{(c)}}{\alpha'} \right)^{-1} e^{\frac{n\pi s^{(c)}}{\alpha'}}$. Therefore the fluctuation of ψ_n is suppressed relative to that of p_0 exponentially by $\sqrt{\left(\frac{n\pi s^{(c)}}{\alpha'} \right) e^{-\frac{n\pi s^{(c)}}{\alpha'}}}$. The mean value of ψ_n ($\overline{\psi}_n$) is $O(q_c^{\frac{n}{2}} \alpha')$. Since it is negligible the integration of ψ_n and $\overline{\psi}_n$ can be accomplished by just replacing it with the unity[7]. After these replacements the above factorization leads to Eq.(4.10).

Let us describe the zero-slope limit of the generating function by using the factorization

---

[7] The Gaussian integral gives suitable products of $e^{\frac{2\alpha' q_c^{\frac{n}{2}}}{n} (e^{\pm i n \sigma_r} k^{(r)}_\mu) G^{\mu\nu} (e^{\mp i n \sigma_r} k^{(r)}_\nu)}$. It can be neglected at the zero-slope limit. Actually speaking, these integrals cancel with the subdominant terms which we abandon at the preceding paragraph.

(6.8). We start with J_T defined by (4.8). The LHS of Eq.(6.8) is the momentum eigenstate factorization of the amplitude (3.42). Hence (6.8) leads to a factorization of J_T at the zero-slope limit. As regards the delta functions we translate them into integrals as given in (4.13). By a straightforward calculation we obtain the following factorization of J_T at the zero-slope limit :

$$
\begin{aligned}
& J_T\left((k^{(1)},\cdots,k^{(M)});(k^{(M+1)},\cdots,k^{(M+N)})\right) \\
& \approx \int dp_0 \prod_{n=1}^{\infty} \frac{d\overline{\psi}_n d\psi_n}{(4\pi i)^{26}} \\
& \quad \times \left[\frac{2\pi}{\alpha'}\int_0^{\infty} ds^{(c)} \exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)\right\}\right. \\
& \qquad\qquad \left.\times \prod_{n=1}^{\infty}\left\{-\det\left(\frac{-\theta G\theta}{8\pi^2 n q_c^{\frac{n}{2}}\alpha'}\right)\exp\left\{\frac{\overline{\psi}_{n\mu}(\theta G\theta)^{\mu\nu}\psi_{n\nu}}{16\pi^2 n q_c^{\frac{n}{2}}\alpha'}\right\}\right\}\right] \\
& \quad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}} \int \frac{dx}{(2\pi)^{26}} e^{ip_{0\mu}x^{\mu}} e^{-\frac{i}{2}p_{0\mu}\theta^{\mu\nu}\sum_{r=1}^{M}k_{\nu}^{(r)}} \\
& \qquad\qquad \times \int \prod_{r=1}^{M} d\sigma_r \prod_{1\le r<s\le M} e^{-\frac{i}{2}k_{\mu}^{(r)}\theta^{\mu\nu}k_{\nu}^{(s)}} \prod_{r=1}^{M} e^{ik_{\mu}^{(r)}x^{\mu}(\sigma_r)} \\
& \quad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}} \int \frac{dx}{(2\pi)^{26}} e^{-ip_{0\mu}x^{\mu}} e^{-\frac{i}{2}p_{0\mu}\theta^{\mu\nu}\sum_{r=M+1}^{M+N}k_{\nu}^{(r)}} \\
& \qquad\qquad \times \int \prod_{r=M+1}^{M+N-1} d\sigma_r \prod_{M+1\le r<s\le M+N} e^{\frac{i}{2}k_{\mu}^{(r)}\theta^{\mu\nu}k_{\nu}^{(s)}} \prod_{r=M+1}^{M+N} e^{ik_{\mu}^{(r)}x^{\mu}(\sigma_r)}. \qquad (6.9)
\end{aligned}
$$

Here we introduce a curve $x(\sigma)$ $(0 \le \sigma \le 2\pi)$ parametrized by p_0 and $(\psi_n, \overline{\psi}_n)$ as follows :

$$
x^{\mu}(\sigma) \equiv x^{\mu} + \frac{\theta^{\mu\nu}}{2\pi}\left\{p_{0\nu}\sigma + \frac{1}{\sqrt{2}}\sum_{n=1}^{\infty}\frac{i}{n}\left(\psi_{n\nu}e^{-in\sigma} - \overline{\psi}_{n\nu}e^{in\sigma}\right)\right\}. \qquad (6.10)
$$

The integrations of σ_r are performed over the region :

$$
\begin{aligned}
& 0 \le \sigma_1 \le \cdots \le \sigma_M \le 2\pi, \\
& 0 \le \sigma_{M+1} \le \cdots \le \sigma_{M+N-1} \le 2\pi. \qquad (6.11)
\end{aligned}
$$

The generating function is defined by (4.12). The zero-slope limit can be obtained from (6.9) by multiplying $\int \prod_r \frac{dk^{(r)}}{(2\pi)^{13}} \prod_r \varphi(k^{(r)})$ and summing up with respect to M and N. These are the same steps as taken previously to obtain the factorization by the straight open Wilson lines. At the present factorization they give us the following zero-slope limit :

$$
\sum_{M=0}^{\infty}\sum_{N=1}^{\infty}\int\prod_{r=1}^{M+N}\frac{dk^{(r)}}{(2\pi)^{13}} J_T\left((k^{(1)},\cdots,k^{(M)});(-k^{(M+1)},\cdots,-k^{(M+N)})\right)
$$

$$
\begin{aligned}
&\times \prod_{r=1}^{M} \varphi(k^{(r)}) \prod_{r=M+1}^{M+N} \varphi(-k^{(r)}) \\
\approx & \int dp_0 \prod_{n=1}^{\infty} \frac{d\overline{\psi}_n d\psi_n}{(4\pi i)^{26}} \\
& \times \left[\frac{2\pi}{\alpha'} \int_0^{\infty} ds^{(c)} \exp\left\{ -\frac{\pi s^{(c)}}{2} \left(p_{0\mu} g^{\mu\nu} p_{0\nu} - \frac{4}{\alpha'} \right) \right\} \right. \\
& \qquad \left. \times \prod_{n=1}^{\infty} \left\{ -\det \left(\frac{-\theta G \theta}{8\pi^2 n q_c^{\frac{n}{2}} \alpha'} \right) \exp\left\{ \frac{\overline{\psi}_{n\mu} (\theta G \theta)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n q_c^{\frac{n}{2}} \alpha'} \right\} \right\} \right] \\
& \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{4}} \int \frac{dx}{(2\pi)^{26}} e^{ip_{0\mu} x^{\mu}} \star \mathcal{P}_{\star} \left[\exp\left\{ \int_0^{2\pi} d\sigma \phi(x(\sigma)) \right\} \right] \\
& \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}} \right)^{\frac{1}{4}} \overline{\int \frac{dx}{(2\pi)^{26}} e^{ip_{0\mu} x^{\mu}} \star \mathcal{P}_{\star} \left[\exp\left\{ \int_0^{2\pi} d\sigma \phi(x(\sigma)) \right\} \right] \star \phi(x(2\pi))}. \quad (6.12)
\end{aligned}
$$

Modulo fixing the $U(1)$ symmetry the generating function is factorized into a sum or an integral of products of two open Wilson lines taken along the curves. The first term in the above integral can be thought of as a measure for the curves. Actually we can decompose $x(\sigma)$ into $x_0(\sigma) + \delta x(\sigma)$. Here $x_0(\sigma)$ is a straight line which connects two ends of the curve. $\delta x(\sigma)$ describes a fluctuation from the straight line and becomes 2π-periodic. x and p_0 in Eq.(6.10) are determined by $x_0(\sigma)$ while ψ_n are determined by $\delta x(\sigma)$. It may be also interesting to describe the curves in terms of loops in the momentum space. Let $p(\sigma)$ be a loop in the momentum space. This gives a curve $x(\sigma)$ by $\dot{x}^{\mu}(\sigma) = \theta^{\mu\nu} p_{\nu}(\sigma)$. p_0 and ψ_n in Eq.(6.10) are determined by $p = \int_0^{2\pi} d\sigma p(\sigma)$ and $\psi_n = \sqrt{2} \int_0^{2\pi} d\sigma e^{in\sigma} p(\sigma)$.

The integration of $\delta x(\sigma)$ are suppressed exponentially compared with $x_0(\sigma)$. It can be accomplished by replacing the integration variables with their mean values and becomes unity at the zero-slope limit. Then the above factorization reduces to that by the straight open Wilson lines.

7 Gluons in Closed-String Theory

Our study of gauge theory starts from this section. An analogue of gluon vertex operator of open-string is introduced in closed-string theory. Investigation of its Bogolubov transformation leads a renormalization of this operator. It will be shown that the renormalized operator enjoys the standard properties of (open-string) gluon vertex operator, including the action of the Virasoro algebra. These operators, acting on the Neumann boundary state, give rise to boundary states which turn out to be identified with the boundary states of (open-string) off-shell gluons. In particular we will show that closed-string tree amplitudes between these states coincide with the corresponding gluon one-loop amplitudes of open-string. Our discussion in this section goes almost parallel to Section 3 where the boundary states of off-shell open-string tachyons are constructed.

7.1 Bogolubov transformation and renormalization

In closed-string theory an analogue of gluon-vertex operator may be taken as

$$\begin{aligned} V_A\left(\sigma,\tau;k\right) &\equiv \,:A_\mu(k)\partial_\sigma X^\mu(\sigma,\tau)\cdot e^{ik_\nu X^\nu(\sigma,\tau)}: \\ &= i:A_\mu(k)\left(z\partial-\bar{z}\bar{\partial}\right)X^\mu(z,\bar{z})\cdot e^{ik_\nu X^\nu(z,\bar{z})}: \ , \end{aligned} \tag{7.1}$$

where $(p+1)$-vectors k_μ and $A_\mu(k)$ are the momentum and the polarization vectors. The polarization vector is the Fourier transform of $U(1)$ gauge field $\mathcal{A}_\mu(x)$:

$$\mathcal{A}_\mu(x)=\int\frac{d^{p+1}k}{(2\pi)^{\frac{p+1}{2}}}A_\mu(k)e^{ik_\nu x^\nu}\ . \tag{7.2}$$

The gauge field takes value in $\mathbb{R}$. This yields $\overline{A_\mu(k)}=A_\mu(-k)$. While we concentrate on $U(1)$ gauge group in this paper, it can be straightforwardly generalized to $U(N)$ by assigning the Chan-Paton indices.

Let us express the above operator in an auxiliary form. This often makes subsequent calculation facile. Let $a\in\mathbb{R}$ be an auxiliary parameter. We write the operator in an exponential form :

$$V_A(\sigma,\tau;k)=i\frac{\partial}{\partial a}\hat{V}_A(\sigma,\tau;k;a)\Big|_{a=0}\ , \tag{7.3}$$

with

$$\begin{aligned} &\hat{V}_A(\sigma,\tau;k;a)\equiv\,:e^{i\left\{k_\mu-aA_\mu(k)\partial_\sigma\right\}X^\mu(\sigma,\tau)}: \\ &\quad=\exp\left(ik_\mu\hat{x}_0^\mu\right)\,|z|^{\alpha' k_\mu\hat{p}_0^\mu} \\ &\quad\times\prod_{n=1}^{\infty}\exp\left[\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\left\{\Big(k_\mu-inaA_\mu(k)\Big)\alpha_{-n}^\mu z^n+\Big(k_\mu+inaA_\mu(k)\Big)\tilde{\alpha}_{-n}^\mu\bar{z}^n\right\}\right] \\ &\quad\times\prod_{n=1}^{\infty}\exp\left[-\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\left\{\Big(k_\mu+inaA_\mu(k)\Big)\alpha_{n}^\mu z^{-n}+\Big(k_\mu-inaA_\mu(k)\Big)\tilde{\alpha}_{n}^\mu\bar{z}^{-n}\right\}\right]\ . \end{aligned} \tag{7.4}$$

The relation (7.3) implies that the terms proportional to higher powers of a become irrelevant to the amplitudes.

We consider the Bogolubov transformation for $\hat{V}_A$ generated by g_N given in Eq.(3.2). Using (3.4) we can write down the transform as follows :

$$g_N^{-1}\hat{V}_A\left(\sigma,\tau;k;a\right)g_N = R(\tau;a)\ \mathrm{ad}_{g_N^{-1}}\hat{V}_A(\sigma,\tau;k;a)\ , \tag{7.5}$$

where

$$\begin{aligned} R(\tau;a) \ \equiv\ & \left(\frac{|z|^2}{|z|^2-1}\right)^{-\frac{\alpha'}{2}k_\mu\left(g^{-1}Ng^{-1}\right)^{\mu\nu}k_\nu} \\ & \times\exp\left[-ia\frac{\alpha'}{2}A_\mu(k)\left(g^{-1}\left(N-N^T\right)g^{-1}\right)^{\mu\nu}k_\nu\frac{1}{|z|^2-1}\right. \\ & \qquad\left.-a^2\frac{\alpha'}{2}A_\mu(k)\left(g^{-1}Ng^{-1}\right)^{\mu\nu}A_\nu(k)\left(\frac{|z|}{|z|^2-1}\right)^2\right] \ . \end{aligned} \tag{7.6}$$

The operator $\mathrm{ad}_{g_N^{-1}}\hat{V}_A(\sigma,\tau;k;a)$ is defined to be

$$\mathrm{ad}_{g_N^{-1}}\hat{V}_A(\sigma,\tau;k;a) = \mathcal{M}_A(\bar{z};k;a)\tilde{\mathcal{M}}_A(z;k;a)\hat{V}_A(\sigma,\tau;k;a)\ , \tag{7.7}$$

where $\mathcal{M}_A(\bar{z};k;a)$ and $\tilde{\mathcal{M}}_A(z;k;a)$ consist of the creation modes alone and take the forms of

$$\begin{aligned} \mathcal{M}_A(\bar{z};k;a) &= \prod_{n=1}^{\infty}\exp\left\{\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\Big(k_\mu - inaA_\mu(k)\Big)\left(g^{-1}N^T\right)^\mu{}_\nu\alpha^\nu_{-n}\bar{z}^{-n}\right\}\ , \\ \tilde{\mathcal{M}}_A(z;k;a) &= \prod_{n=1}^{\infty}\exp\left\{\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\Big(k_\mu + inaA_\mu(k)\Big)\left(g^{-1}N\right)^\mu{}_\nu\tilde{\alpha}^\nu_{-n}z^{-n}\right\}\ . \end{aligned} \tag{7.8}$$

In the transform (7.5) we can find the same singularity structure as that of the tachyon. It is due to the factor $R(\tau;a)$, which is responsible for the singularity of the transform (7.5) at the world-sheet boundary, $\tau=0(\Leftrightarrow|z|=1)$. This factor contains an exponential in addition to the same factor which appeared in the transform $g_N^{-1}V_Tg_N$. Let us make a few comments about this extra factor. The first term of the exponent is proportional to a. The closed-string tensor $g^{-1}(N-N^T)g^{-1}$ used there is translated into $\frac{2}{\pi\alpha'}\theta^{\mu\nu}$. The second term becomes irrelevant since it is proportional to a^2. Therefore the net effect of this extra factor is to give a term proportional to θ. Particularly in the absence of a constant B field, $R(\tau;a)$ reduces to that of the tachyon.

The singular factor which appears in the transform of the tachyon vertex operator has been expressed in Eq.(3.12) as the self-contraction between the chiral and the anti-chiral parts of V_T. In the present case as well, we can think of the singular factor as such a self-contraction of $\hat{V}_A(\sigma,\tau;k;a)$. In fact, we can express $R(\tau;a)$ by using the chiral–anti-chiral correlation

$\langle X^\mu(z_1)\tilde{X}^\nu(\bar{z}_2)\rangle = -\frac{\alpha'}{2}\left(g^{-1}Ng^{-1}\right)^{\mu\nu}\ln\left(1-\frac{1}{z_1\bar{z}_2}\right)$ of Green's function $\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)$ as follows :

$$R(\tau;a) = \exp\left[-\left(k_\mu - iaA_\mu(k)z\partial\right)\left(k_\nu + iaA_\nu(k)\bar{z}\bar{\partial}\right)\left\langle X^\mu(z)\tilde{X}^\nu(\bar{z})\right\rangle\right] \; . \tag{7.9}$$

As mentioned in the analysis of tachyon, boundary states induce correlations between the chiral and the anti-chiral sectors, and this feature is characteristic of systems of interacting closed- and open-strings.

We wish to find out a relation between $\mathrm{ad}_{g_N^{-1}}\hat{V}_A$ and open-string gluon vertex operators. Previous discussions to establish the relation between $\mathrm{ad}_{g_N^{-1}}V_T$ and open-string tachyon vertex operators consist of : i) Description of the OPE between $\mathrm{ad}_{g_N^{-1}}V_T$, which turns out to be given in (3.9) by using Green's function $\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)$ and allows us to introduce the renormalized open-string tachyon vertex operators V_T^{ren}, ii) description of the diffS^1 action on V_T^{ren}, which turns out to reduce at the boundary to the standard Virasoro action on open-string tachyon vertex operators, and iii) reproduction of open-string one-loop tachyon amplitudes by the boundary state formalism. We repeat these analyses as for gluons in the following.

Let us examine the OPE between $\mathrm{ad}_{g_N^{-1}}\hat{V}_A$. It is convenient to start with the OPE between the auxiliary operators. The standard calculation leads to

$$\begin{aligned}
&\hat{V}_A\left(\sigma_1,\tau_1;k^{(1)};a_1\right)\hat{V}_A\left(\sigma_2,\tau_2;k^{(2)};a_2\right)\\
&\quad = \exp\Big[i\left\{k_{(1)} - ia_1A_\mu(k^{(1)})\left(z_1\partial_{z_1} - \bar{z}_1\bar{\partial}_{\bar{z}_1}\right)\right\}\\
&\qquad\qquad \times i\left\{k_\mu^{(2)} - ia_2A_\mu(k^{(2)})\left(z_2\partial_{z_2} - \bar{z}_2\bar{\partial}_{\bar{z}_2}\right)\right\}\left(-\frac{\alpha'}{2}g^{\mu\nu}\ln|z_1-z_2|^2\right)\Big]\\
&\qquad \times :\hat{V}_A\left(\sigma_1,\tau_1;k^{(1)};a_1\right)\hat{V}_A\left(\sigma_2,\tau_2;k^{(2)};a_2\right): \; ,
\end{aligned} \tag{7.10}$$

for $|z_1| \geq |z_2|$ and $z_1 \neq z_2$. Since $\mathrm{ad}_{g_N^{-1}}\hat{V}_A$ have the form given in (7.7), we can obtain their OPE from the above by taking account of the OPEs between $\mathcal{M}_A, \tilde{\mathcal{M}}_A$ and $\hat{V}_A$. We find that

$$\begin{aligned}
&\mathrm{ad}_{g_N^{-1}}\hat{V}_A\left(\sigma_1,\tau_1;k^{(1)};a_1\right)\;\mathrm{ad}_{g_N^{-1}}\hat{V}_A\left(\sigma_2,\tau_2;k^{(2)};a_2\right)\\
&\quad = \exp\Big[i\Big\{k_\mu^{(1)} - ia_1A_\mu(k^{(1)})\left(z_1\partial_{z_1} - \bar{z}_1\bar{\partial}_{\bar{z}_1}\right)\Big\}\\
&\qquad\qquad \times i\Big\{k_\mu^{(2)} - ia_2A_\mu(k^{(2)})\left(z_2\partial_{z_2} - \bar{z}_2\bar{\partial}_{\bar{z}_2}\right)\Big\}\mathbb{G}^{\mu\nu}(z_1,\bar{z}_1|z_2,\bar{z}_2)\Big]\\
&\qquad \times :\mathrm{ad}_{g_N^{-1}}\hat{V}_A\left(\sigma_1,\tau_1;k^{(1)};a_1\right)\;\mathrm{ad}_{g_N^{-1}}\hat{V}_A\left(\sigma_2,\tau_2;k^{(2)};a_2\right): \; ,
\end{aligned} \tag{7.11}$$

for $|z_1| \geq |z_2|$, $|z_1z_2| \geq 1$ and $z_1 \neq z_2$. This is a direct extension of the tachyon case (3.9) and suggests an interpretation of $\mathrm{ad}_{g_N^{-1}}\hat{V}_A$ in terms of the gluon vertex operator.

Let us introduce the renormalized operator $\hat{V}_A^{ren}$ by the following subtraction :

$$\hat{V}_A^{ren}\left(\sigma,\tau;k;a\right) \equiv e^{\frac{i}{2\pi}aA_\mu(k)\theta^{\mu\nu}k_\nu} R(\tau;a)^{-1}\hat{V}_A\left(\sigma,\tau;k;a\right) \ . \tag{7.12}$$

In other words,

$$g_N^{-1}\hat{V}_A^{ren}\left(\sigma,\tau;k;a\right)g_N = e^{\frac{i}{2\pi}aA_\mu(k)\theta^{\mu\nu}k_\nu}\ \mathrm{ad}_{g_N^{-1}}\hat{V}_A(\sigma,\tau;k;a) \ . \tag{7.13}$$

In addition to the subtraction of the singular factor $R(\tau;a)$, a finite subtraction is made in the above by the multiplication of $e^{\frac{i}{2\pi}aA_\mu(k)\theta^{\mu\nu}k_\nu}$. Although it vanishes in the absence of B field, this factor becomes surely necessary in general. Without this subtraction we cannot reproduce even the standard Virasoro action which we present below. The renormalized operator (7.12) gives rise to V_A^{ren} by the relation (7.3). We refer to $V_A^{ren}(\sigma,\tau;k)$ as renormalized gluon vertex operator with momentum k_μ and polarization $A_\mu(k)$. It can be written as follows :

$$\begin{aligned}
V_A^{ren}\left(\sigma,\tau;k\right) &\equiv i\frac{\partial}{\partial a}\hat{V}_A^{ren}\left(\sigma,\tau;k;a\right)\Big|_{a=0} \\
&= \left(\frac{|z|^2}{|z|^2-1}\right)^{\frac{\alpha'}{2}k_\mu\left(g^{-1}Ng^{-1}\right)^{\mu\nu}k_\nu} \\
&\qquad \times\left\{V_A(\sigma,\tau;k) - \frac{1}{\pi}A_\mu(k)\theta^{\mu\nu}k_\nu\left(\frac{1}{|z|^2-1}+\frac{1}{2}\right)V_T(\sigma,\tau;k)\right\} \ ,
\end{aligned} \tag{7.14}$$

where $V_T(\sigma,\tau;k)$ $\left(=\hat{V}_A(\sigma,\tau;k;a=0)\right)$ is the closed-string tachyon vertex operator.

Next we will examine the action of diffS^1 on the above renormalized operators. As a shortcut we use the description in terms of boundary states. Relevant states are introduced as $\left|\hat{B}_N[A];a;(\sigma,k)\right\rangle$ and $\left|B_N[A];(\sigma,k)\right\rangle$. The first states are defined by using $\hat{V}_A^{ren}$ as follows :

$$\begin{aligned}
\left|\hat{B}_N[A];a;(\sigma,k)\right\rangle &\equiv \lim_{\tau\to0+}\hat{V}_A^{ren}(\sigma,\tau;k;a)\left|B_N\right\rangle \\
&= \left(\frac{\left(\det E_{\mu\nu}\right)^2}{\left(2\alpha'\right)^{p+1}\left(-\det g_{\mu\nu}\right)}\right)^{\frac{1}{4}} \lim_{\tau\to0+} g_N\left(g_N^{-1}\hat{V}_A^{ren}(\sigma,\tau;k;a)g_N\right)|\mathbf{0}\rangle \\
&= \left(\frac{-\det G_{\mu\nu}}{\left(2\alpha'\right)^{p+1}}\right)^{\frac{1}{4}} e^{\frac{i}{2\pi}aA_\mu(k)\theta^{\mu\nu}k_\nu}\, g_N \lim_{\tau\to0+}\mathrm{ad}_{g_N^{-1}}\hat{V}_A(\sigma,\tau;k;a)|\mathbf{0}\rangle \\
&= \left(\frac{-\det G_{\mu\nu}}{\left(2\alpha'\right)^{p+1}}\right)^{\frac{1}{4}} e^{\frac{i}{2\pi}aA_\mu(k)\theta^{\mu\nu}k_\nu} \\
&\qquad \times\prod_{n=1}^{\infty}\exp\left[\frac{\sqrt{2\alpha'}}{n}\left\{\left(k_\mu - inaA_\mu(k)\right)\left(\frac{1}{E^T}g\right)^\mu{}_\nu\alpha_{-n}^\nu e^{in\sigma}\right.\right. \\
&\qquad\qquad \left.\left. +\left(k_\mu + inaA_\mu(k)\right)\left(\frac{1}{E}g\right)^\mu{}_\nu\tilde{\alpha}_{-n}^\nu e^{-in\sigma}\right\}\right] g_N|k_N\rangle \ ,
\end{aligned} \tag{7.15}$$

where Eqs.(7.7) and (7.13) are utilized to obtain the oscillator representation. The second states are defined similarly by using V_A^{ren} as

$$\Big|B_N[A];(\sigma,k)\Big\rangle \equiv \lim_{\tau\to 0+} V_A^{ren}(\sigma,\tau;k)\,|B_N\rangle = i\frac{\partial}{\partial a}\Big|\hat{B}_N[A];a;(\sigma,k)\Big\rangle\Big|_{a=0} \ . \qquad (7.16)$$

The oscillator representation can be obtained from (7.15). This yields

$$\begin{aligned}\Big|B_N[A];(\sigma,k)\Big\rangle &= i\frac{\partial}{\partial a}\Big|\hat{B}_N[A];a;(\sigma,k)\Big\rangle\Big|_{a=0}\\ &= \left[-\frac{1}{2\pi}A_\mu(k)\theta^{\mu\nu}k_\nu + \sum_{n=1}^{\infty}\sqrt{2\alpha'}A_\mu(k)\left\{\left(\frac{1}{E^T}g\right)^\mu{}_\nu \alpha^\nu_{-n}e^{in\sigma} - \left(\frac{1}{E}g\right)^\mu{}_\nu \tilde{\alpha}^\nu_{-n}e^{-in\sigma}\right\}\right]\\ &\quad\times|B_N;(\sigma,k)\rangle \ , \end{aligned} \qquad (7.17)$$

where $|B_N;(\sigma,k)\rangle$ is the boundary state with a single open-string tachyon given in (3.20).

Let us recall that generators of diffS^1 are $L_m - \tilde{L}_{-m}$ ($m \in \mathbb{Z}$). Their actions on the states $\Big|\hat{B}_N[A];a;(\sigma,k)\Big\rangle$ turn out to be as follows :

$$\begin{aligned}&\Big(L_m - \tilde{L}_{-m}\Big)\Big|\hat{B}_N[A];a;(\sigma,k)\Big\rangle\\ &\quad = e^{im\sigma}\Bigg[\sqrt{2\alpha'}\sum_{n=1}^{\infty}\{k_\mu - i(n+m)aA_\mu(k)\}\left(\frac{1}{E^T}g\right)^\mu{}_\nu \alpha^\nu_{-n}e^{in\sigma}\\ &\qquad -\sqrt{2\alpha'}\sum_{n=1}\{k_\mu + i(n-m)aA_\mu(k)\}\left(\frac{1}{E}g\right)^\mu{}_\nu \tilde{\alpha}^\nu_{-n}e^{-in\sigma}\\ &\qquad +\alpha' m k_\mu G^{\mu\nu}k_\nu + iaA_\mu(k)\left(\frac{m}{2\pi}\theta^{\mu\nu} - \alpha' m^2 G^{\mu\nu}\right)k_\nu\\ &\qquad -a^2\alpha'\frac{1}{6}m(m-1)(m+1)A_\mu(k)\,G^{\mu\nu}A_\nu(k)\Bigg]\Big|\hat{B}_N[A];a;(\sigma,k)\Big\rangle \ , \end{aligned} \qquad (7.18)$$

for $\forall m \in \mathbb{Z}$. As regards the states $\Big|B_N[A];(\sigma,k)\Big\rangle$, the actions of $L_n - \tilde{L}_{-n}$ can be read from the above by differentiating it with respect to a and then setting $a = 0$. We obtain

$$\begin{aligned}\Big(L_m - \tilde{L}_{-m}\Big)\Big|B_N[A];(\sigma,k)\Big\rangle &= e^{im\sigma}\left\{-i\frac{\partial}{\partial\sigma} + m\left(\alpha' k_\mu G^{\mu\nu}k_\nu + 1\right)\right\}\Big|B_N[A];(\sigma,k)\Big\rangle\\ &\quad + e^{im\sigma}\alpha' m^2 k_\mu G^{\mu\nu}A_\nu(k)\,|B_N;(\sigma,k)\rangle \ . \end{aligned} \qquad (7.19)$$

The RHS can be compared with the standard Virasoro action on (open-string) gluon vertex operators of the same momenta and polarizations. We find that they are identical with each other.

The Ishibashi condition imposed on $\Big|B_N[A];(\sigma,k)\Big\rangle$, namely the vanishing of the RHS of Eq.(7.19), modulo total derivative with respect to σ, requires the well-known physical state condition of the gauge field :

$$\alpha' k_\mu G^{\mu\nu} k_\nu = 0 \ , \quad k_\mu G^{\mu\nu} A_\nu(k) = 0 \ . \tag{7.20}$$

By the same argument presented for the tachyon the above action is translated in closed-string field theory to the action of the BRST charge Q_c. The BRST invariance of these boundary states becomes precisely the on-shell condition of the gauge field.

7.2 Boundary states with M gluons

7.2.1 Construction of boundary states with M gluons

Previous constructions (7.15) and (7.17) are generalized to the cases of M off-shell gluons. Let us start with the states

$$\begin{aligned}
&\hat{V}_A^{ren}\left(\sigma_1,\tau_1;k^{(1)};a_1\right)\cdots\hat{V}_A^{ren}\left(\sigma_M,\tau_M;k^{(M)};a_M\right)|B_N\rangle \\
&\quad = \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} g_N \left(g_N^{-1}\hat{V}_A^{ren}(\sigma_1,\tau_1;k^{(1)};a_1)g_N\right)\cdots \\
&\qquad\qquad \cdots\left(g_N^{-1}\hat{V}_A^{ren}(\sigma_M,\tau_M;k^{(M)};a_M)g_N\right)|\mathbf{0}\rangle \ ,
\end{aligned} \tag{7.21}$$

where the insertion points $z_r = e^{\tau_r + i\sigma_r}$ $(r = 1, 2, \ldots, M)$ of the auxiliary renormalized operators $\hat{V}_A^{ren}$ are distinct with each other, satisfying the condition $|z_1| \geq |z_2| \geq \cdots |z_M| \geq 1$. The oscillator representation can be obtained by taking account of the relation (7.13) and then using the OPE (7.11). It turns out to be

$$\begin{aligned}
&\hat{V}_A^{ren}\left(\sigma_1,\tau_1;k^{(1)};a_1\right)\cdots\hat{V}_A^{ren}\left(\sigma_M,\tau_M;k^{(M)};a_M\right)|B_N\rangle \\
&\quad = \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \prod_{r=1}^{M} e^{\frac{i}{2\pi}a_r A_\mu(k^{(r)})\theta^{\mu\nu}k^{(r)}} \\
&\qquad \times \prod_{r<s}^{M}\exp\Big[i\Big\{k_\mu^{(r)} - ia_r A_\mu(k^{(r)})\left(z_r\partial_{z_r} - \bar{z}_r\bar{\partial}_{\bar{z}_r}\right)\Big\} \\
&\qquad\qquad\qquad \times i\Big\{k_\mu^{(s)} - ia_s A_\mu(k^{(s)})\left(z_s\partial_{z_s} - \bar{z}_s\bar{\partial}_{\bar{z}_s}\right)\Big\}\mathbb{G}^{\mu\nu}(z_r,\bar{z}_r|z_s,\bar{z}_s)\Big] \\
&\qquad \times \prod_{n=1}^{\infty}\exp\Big[\sqrt{\frac{\alpha'}{2}}\frac{1}{n}\sum_{r=1}^{M}\Big\{\left(k_\mu^{(r)} - ina_r A_\mu(k^{(r)})\right)\left(z_r^n\delta_\nu^\mu + \bar{z}_r^{-n}\left(g^{-1}N^T\right)^\mu{}_\nu\right)\alpha_{-n}^\nu \\
&\qquad\qquad\qquad\qquad + \left(k_\mu^{(r)} + ina_r A_\mu(k^{(r)})\right)\left(\bar{z}_r\delta_\nu^\mu + z_r^{-n}\left(g^{-1}N\right)^\mu{}_\nu\right)\tilde{\alpha}_{-n}^\nu\Big\}\Big]
\end{aligned}$$

$$\times g_N \left| \sum_{r=1}^{M} k_N^{(r)} \right\rangle \ . \tag{7.22}$$

Limits $\forall \tau_r \to 0+$ of the above states may be interpreted as auxiliary boundary states of M off-shell gluons. They are auxiliary since the parameters a_r are still included. This limiting process corresponds to sending the operators onto the world-sheet boundary. We then obtain the following states :

$$\begin{aligned}
&\left| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), (\sigma_2, k^{(2)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \\
&\quad \equiv \lim_{\forall \tau_r \to 0+} \hat{V}_A^{ren}\left(\sigma_1, \tau_1; k^{(1)}; a_1\right) \cdots \hat{V}_A^{ren}\left(\sigma_M, \tau_M; k^{(M)}; a_M\right) |B_N\rangle \\
&\quad = \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} e^{\frac{i}{2\pi} \sum_{r=1}^{M} \left(a_r A_\mu(k^{(r)}) - \sigma_r k_\mu^{(r)} \right) \theta^{\mu\nu} \sum_{s=1}^{M} k_\nu^{(s)}} \prod_{r<s}^{M} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \\
&\qquad \times \prod_{r<s}^{M} \exp\left[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\nu^{(s)} - a_s A_\nu(k^{(s)}) \partial_{\sigma_s} \right) \ln \left| e^{i\sigma_r} - e^{i\sigma_s} \right| \right] \\
&\qquad \times \prod_{n=1}^{\infty} \exp\left[\frac{\sqrt{2\alpha'}}{n} \sum_{r=1}^{M} \left\{ \left(k_\mu^{(r)} - ina_r A_\mu(k^{(r)}) \right) \left(\frac{1}{E^T} g \right)^{\mu}{}_{\nu} \alpha_{-n}^{\nu} e^{in\sigma_r} \right. \right. \\
&\qquad\qquad \left. \left. + \left(k_\nu^{(r)} + ina_r A_\mu(k^{(r)}) \right) \left(\frac{1}{E} g \right)^{\mu}{}_{\nu} \tilde{\alpha}_{-n}^{\nu} e^{-in\sigma_r} \right\} \right] g_N \left| \sum_{r=1}^{M} k_N^{(r)} \right\rangle \ .
\end{aligned} \tag{7.23}$$

We can repeat the above argument with $\hat{V}_A^{ren}$ replaced by V_A^{ren}. This gives rise to the states

$$\left| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \equiv \lim_{\forall \tau_r \to 0+} V_A^{ren}(\sigma_1, \tau_1; k^{(1)}) \cdots V_A^{ren}(\sigma_M, \tau_M; k^{(M)}) |B_N\rangle . \tag{7.24}$$

These states are expected to be identified with boundary states of M off-shell gluons. In order to justify this identification, we should evaluate closed-string free propagations between these boundary states and compare them with the corresponding open-string one-loop amplitudes. Nevertheless, the oscillator representation of the states (7.24) becomes too complicated to evaluate the amplitudes. Taking account of the relation (7.14) it becomes convenient to use instead the following equivalent realization :

$$\begin{aligned}
&\left| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \\
&\qquad = i^M \prod_{r=1}^{M} \left. \frac{\partial}{\partial a_r} \right|_{a_r=0} \left| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle ,
\end{aligned} \tag{7.25}$$

where the operation $\prod_{r=1}^{M} \frac{\partial}{\partial a_r} \Big|_{a_r=0}$ means putting $a_r = 0$ $(\forall r)$ after the differentiations.

We note that the boundary states (7.25) become available in the absence of B field by putting $B_{\mu\nu} = 0$.

7.2.2 Construction of dual boundary states

The dual boundary states are required in order to obtain the closed-string tree amplitudes. We construct these states by taking the same route as the tachyon case.

We begin by considering the Bogolubov transform of $\hat{V}_A$ generated by $g_N^\dagger$ which is the BPZ dual of g_N given in Eq.(3.32). We have

$$g_N^\dagger \hat{V}_A(\sigma, \tau; k; a) \left(g_N^\dagger\right)^{-1} = R_\infty(\tau, a) \ \mathrm{ad}_{g_N^\dagger} \hat{V}_A(\sigma, \tau; k; a) \ . \tag{7.26}$$

Here we introduce a regular operator $\mathrm{ad}_{g_N^\dagger} \hat{V}_A(\sigma, \tau; k; a)$ defined as

$$\mathrm{ad}_{g_N^\dagger} \hat{V}_A(\sigma, \tau; k; a) = \hat{V}_A(\sigma, \tau; k; a) \mathcal{M}_{A\infty}(\bar{z}; k; a) \tilde{\mathcal{M}}_{A\infty}(z; k; a) \ , \tag{7.27}$$

with

$$\begin{aligned} \mathcal{M}_{A\infty}(\bar{z}; k; a) &= \prod_{n=1}^{\infty} \exp\left[-\sqrt{\frac{\alpha'}{2}} \frac{1}{n} \left(k_\mu + inaA_\mu(k)\right) \left(g^{-1} N^T\right)^\mu{}_\nu \alpha_n^\nu \bar{z}^n \right] \ , \\ \tilde{\mathcal{M}}_{A\infty}(z; k; a) &= \prod_{n=1}^{\infty} \exp\left[-\sqrt{\frac{\alpha'}{2}} \frac{1}{n} \left(k_\mu - inaA_\mu(k)\right) \left(g^{-1} N\right)^\mu{}_\nu \tilde{\alpha}_n^\nu z^n \right] \ . \end{aligned} \tag{7.28}$$

$R_\infty(\tau; a)$ in the above denotes the singular factor. It takes the same form as $R(\tau; a)$ with z and $\theta^{\mu\nu}$ replaced by $1/z$ and $-\theta^{\mu\nu}$ respectively :

$$\begin{aligned} R_\infty(\tau, a) &= \left(1 - |z|^2\right)^{\frac{\alpha'}{2} k_\mu (g^{-1} N g^{-1})^{\mu\nu} k_\nu} \\ &\quad \times \exp\left[-a^2 \frac{\alpha'}{2} A_\mu(k) \left(g^{-1} N g^{-1}\right)^{\mu\nu} A_\nu(k) \left(\frac{|z|}{1 - |z|^2}\right)^2 \right. \\ &\qquad \left. + ia \frac{\alpha'}{2} A_\mu(k) \left(g^{-1} \left(N - N^T\right) g^{-1}\right)^{\mu\nu} k_\nu \frac{|z|^2}{1 - |z|^2} \right] \ . \end{aligned} \tag{7.29}$$

This factor represents the self-contraction between the chiral and the anti-chiral pieces of $\hat{V}_A$. It can be written in terms of the chiral–anti-chiral correlation $\langle X^\mu(z_1) \tilde{X}^\nu(\bar{z}_2) \rangle_\infty$ of Green's function $\mathbb{G}_\infty^{\mu\nu}(z_1, \bar{z}_1 | z_2, \bar{z}_2)$:

$$R_\infty(\tau; a) = \exp\left[-\left(k_\mu - iaA_\mu(k) z\partial\right) \left(k_\nu + iaA_\nu(k) \bar{z}\bar{\partial}\right) \left\langle X^\mu(z) \tilde{X}^\nu(\bar{z}) \right\rangle_\infty \right] \ . \tag{7.30}$$

Dual of the auxiliary renormalized gluon vertex operator, $\hat{V}_{A\infty}^{ren}(\sigma, \tau; k; a)$ is introduced by the following subtractions :

$$\hat{V}_{A\infty}^{ren}(\sigma, \tau; k; a) = e^{-\frac{i}{2\pi} a A_\mu(k) \theta^{\mu\nu} k_\nu} R_\infty(\tau; a)^{-1} \hat{V}_A(\sigma, \tau; k; a) \ , \tag{7.31}$$

where the finite subtraction has been made as required previously. Dual of the renormalized gluon operator, which we call $V^{ren}_{A\infty}(\sigma,\tau;k)$, is obtained from the above by using the relation (7.3) :

$$\begin{aligned}
V^{ren}_{A\infty}(\sigma,\tau;k;a) &\equiv i\frac{\partial}{\partial a}\hat{V}^{ren}_{A\infty}(\sigma,\tau;k;a)\bigg|_{a=0} \\
&= \left(1-|z|^2\right)^{-\frac{\alpha'}{2}k_\mu\left(g^{-1}Ng^{-1}\right)^{\mu\nu}k_\nu} \\
&\quad\times\left\{V_A(\sigma,\tau;k)+\frac{1}{\pi}\left(A_\mu(k)\theta^{\mu\nu}k_\nu\right)\left(\frac{|z|^2}{1-|z|^2}+\frac{1}{2}\right)V_T(\sigma,\tau;k)\right\} \ . \qquad (7.32)
\end{aligned}$$

Let us describe the dual boundary states. We first consider the states

$$\langle B_N|\hat{V}^{ren}_{A\infty}(\sigma_1,\tau_1;k^{(1)};a_1)\cdots\hat{V}^{ren}_{A\infty}(\sigma_M,\tau_M;k^{(M)};a_M)$$

where the insertion points are all distinct satisfying the condition, $1\geq|z_1|\geq|z_2|\geq\cdots\geq|z_M|$. We then take the limit $\forall\tau_r\rightarrow 0-$ of these states. Thus we obtain the duals. They take the forms of

$$\begin{aligned}
&\left\langle\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),(\sigma_2,k^{(2)}),\cdots,(\sigma_M,k^{(M)})\right| \\
&\quad\equiv\lim_{\forall\tau_r\rightarrow 0-}\langle B_N|\hat{V}^{ren}_{A\infty}(\sigma_1,\tau_1;k^{(1)};a_1)\cdots\hat{V}^{ren}_{A\infty}(\sigma_M,\tau_M;k^{(M)};a_M) \\
&\quad=\left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}}e^{-\frac{i}{2\pi}\sum_{r,s=1}^{M}\left(a_rA_\mu(k^{(r)})-\sigma_rk^{(r)}_\mu\right)\theta^{\mu\nu}k^{(s)}_\nu}\prod_{r<s}^{M}e^{-\frac{i}{2}k^{(r)}_\mu\theta^{\mu\nu}k^{(s)}_\nu\epsilon(\sigma_r-\sigma_s)} \\
&\qquad\times\prod_{r<s}^{M}\exp\Big[2\alpha'G^{\mu\nu}\left(k^{(r)}_\mu-a_rA_\mu(k^{(r)})\partial_{\sigma_r}\right)\left(k^{(s)}_\mu-a_sA_\mu(k^{(s)})\partial_{\sigma_s}\right)\ln\left|e^{i\sigma_r}-e^{i\sigma_s}\right|\Big] \\
&\qquad\times\left\langle-\sum_{r=1}^{M}k^{(r)}_N\right|g^\dagger_N \\
&\qquad\quad\times\prod_{n=1}^{\infty}\exp\left[-\frac{\sqrt{2\alpha'}}{n}\sum_{r=1}^{M}\left\{\left(k^{(r)}_\mu+ina_rA_\mu(k^{(r)})\right)\left(\frac{1}{E^T}g\right)^{\mu}{}_{\nu}\alpha^\nu_ne^{-in\sigma_r}\right.\right. \\
&\qquad\qquad\left.\left.+\left(k^{(r)}_\mu-ina_rA_\mu(k^{(r)})\right)\left(\frac{1}{E}g\right)^{\mu}{}_{\nu}\tilde{\alpha}^\nu_ne^{in\sigma_r}\right\}\right] \ . \qquad (7.33)
\end{aligned}$$

The oscillator representation in the above is derived from Eq.(7.31) and the following OPE between $\mathrm{ad}_{g^\dagger_N}\hat{V}_A$:

$$\begin{aligned}
&\mathrm{ad}_{g^\dagger_N}\hat{V}_A(\sigma_1,\tau_1;k^{(1)};a_1)\ \ \mathrm{ad}_{g^\dagger_N}\hat{V}_A(\sigma_2,\tau_2;k^{(2)};a_2) \\
&\quad=\exp\Big[-\Big\{k^{(1)}_\mu-ia_1A_\mu(k^{(1)})\left(z_1\partial_1-\bar{z}_1\bar{\partial}_1\right)\Big\} \\
&\qquad\qquad\times\Big\{k^{(2)}_\nu-ia_2A_\nu(k^{(2)})\left(z_2\partial_2-\bar{z}_2\bar{\partial}_2\right)\Big\}\mathbb{G}^{\mu\nu}_\infty(z_1,\bar{z}_1|z_2,\bar{z}_2)\Big] \\
&\qquad\times:\mathrm{ad}_{g^\dagger_N}\hat{V}_A(\sigma_1,\tau_1;k^{(1)};a_1)\ \ \mathrm{ad}_{g^\dagger_N}\hat{V}_A(\sigma_2,\tau_2;k^{(2)};a_2): \ , \qquad (7.34)
\end{aligned}$$

for $|z_1| \geq |z_2|$, $|z_1 z_2| \leq 1$ and $z_1 \neq z_2$. Duals to the states (7.24) are obtained from (7.33) as follows :

$$\begin{aligned} &\Big\langle B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \Big| \\ &\qquad = i^M \prod_{r=1}^{M} \frac{\partial}{\partial a_r}\bigg|_{a_r=0} \Big\langle \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \Big| \ . \end{aligned} \tag{7.35}$$

7.2.3 Closed-string propagation

We now come to computations of closed-string tree amplitudes between the boundary states (7.24). It is a straightforward generalization of what we did for the boundary states of off-shell open-string tachyons but becomes much complicated.

To avoid an unnecessary complication and make the result transparent let us calculate the following amplitudes :

$$\begin{aligned} &\Big\langle \hat{B}_N[A]; \{a_r\}; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \Big| \\ &\qquad \times\, q_c^{\frac{1}{2}\left(L_0 + \tilde{L}_0 - 2\right)} \Big| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \Big\rangle, \end{aligned} \tag{7.36}$$

where $q_c = e^{2\pi i \tau^{(c)}}$. These amplitudes reduce to the amplitudes in question by using the relations (7.25) and (7.35).

We can factorize the amplitudes into products of two kinds of contributions from the zero-modes and the massive modes of closed-string. By using the oscillator representations (7.23) and (7.33) these become as follows :

$$\begin{aligned} &\Big\langle \hat{B}_N[A]; \{a_r\}; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \Big| \\ &\qquad \times q_c^{\frac{1}{2}\left(L_0 + \tilde{L}_0 - 2\right)} \Big| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \Big\rangle \\ &= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{2}} \delta^{(p+1)}\left(\sum_{r=1}^{M+N} k^{(r)} \right) \ q_c^{-1-\frac{\alpha'}{4}\sum_{r=1}^{M}\sum_{s=M+1}^{M+N} k_\mu^{(r)} g^{\mu\nu} k_\nu^{(s)}} \\ &\times \prod_{1\leq r<s\leq M} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \prod_{M+1\leq r<s\leq M+N} e^{-\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \\ &\times e^{\frac{i}{2\pi}\sum_{r,s=1}^{M}\left(a_r A_\mu(k^{(r)}) - \sigma_r k_\mu^{(r)} \right)\theta^{\mu\nu} k_\nu^{(s)}} \ e^{-\frac{i}{2\pi}\sum_{r,s=M+1}^{M+N}\left(a_r A_\mu(k^{(r)}) - \sigma_r k_\mu^{(r)} \right)\theta^{\mu\nu} k_\nu^{(s)}} \\ &\times \prod_{1\leq r<s\leq M} \exp\left[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\mu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s} \right) \ln\left| e^{i\sigma_r} - e^{i\sigma_s} \right| \right] \\ &\times \prod_{M+1\leq r<s\leq M+N} \exp\left[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\mu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s} \right) \ln\left| e^{i\sigma_r} - e^{i\sigma_s} \right| \right] \\ &\times F_A\left(q_c, \{\sigma_r\}, \left\{k^{(r)}\right\}; \{a_r\} \right) \ . \end{aligned} \tag{7.37}$$

Here F_A represents the sum of contributions from the massive modes of closed-string. It is given by the following infinite products :

$$
\begin{aligned}
&F_A\left(q_c, \{\sigma_r\}, \left\{k^{(r)}\right\}; \{a_r\}\right) \\
&\quad = \prod_{n=1}^{\infty} \langle 0| \exp\Bigg[-\frac{q_c^n}{n} \alpha_n^\mu N_{\mu\nu} \tilde{\alpha}_n^\nu \\
&\qquad\qquad - \frac{\sqrt{2\alpha'} q_c^{\frac{n}{2}}}{n} \sum_{r=M+1}^{M+N} \Bigg\{ \left(k_\mu^{(r)} + ina_r A_\mu(k^{(r)})\right) \left(\frac{1}{E^T} g\right)^\mu{}_\nu \alpha_n^\nu e^{-in\sigma_r} \\
&\qquad\qquad\qquad + \left(k_\mu^{(r)} - ina_r A_\mu(k^{(r)})\right) \left(\frac{1}{E} g\right)^\mu{}_\nu \tilde{\alpha}_n^\nu e^{in\sigma_r} \Bigg\} \Bigg] \\
&\qquad \times \exp\Bigg[-\frac{1}{n} \alpha_{-n}^\mu N_{\mu\nu} \tilde{\alpha}_{-n}^\nu \\
&\qquad\qquad + \frac{\sqrt{2\alpha'}}{n} \sum_{r=1}^{M} \Bigg\{ \left(k_\mu^{(r)} - ina_r A_\mu(k^{(r)})\right) \left(\frac{1}{E^T} g\right)^\mu{}_\nu \alpha_{-n}^\nu e^{in\sigma_r} \\
&\qquad\qquad\qquad + \left(k_\mu^{(r)} + ina_r A_\mu(k^{(r)})\right) \left(\frac{1}{E} g\right)^\mu{}_\nu \tilde{\alpha}_{-n}^\nu e^{-in\sigma_r} \Bigg\} \Bigg] |0\rangle \ .
\end{aligned}
\tag{7.38}
$$

We need to evaluate the above infinite products. These are carried out in Appendix B. We just quote the result obtained there. These turn out to be as follows :

$$
\begin{aligned}
&F_A\left(q_c, \{\sigma_r\}, \left\{k^{(r)}\right\}; \{a_r\}\right) \\
&= \prod_{n=1} \left(1 - q_c^n\right)^{-p-1} \\
&\quad \times \prod_{1 \le r < s \le M} \exp\Bigg[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r}\right) \left(k_\nu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s}\right) \\
&\qquad\qquad \times \ln \left\{ \frac{\prod_{n=1}^\infty \left(1 - e^{i(\sigma_r - \sigma_s)} q_c^n\right) \left(1 - e^{-i(\sigma_r - \sigma_s)} q_c^n\right)}{\prod_{n=1}^\infty \left(1 - q_c^n\right)^2} \right\} \Bigg] \\
&\quad \times \prod_{M+1 \le r < s \le M+N} \exp\Bigg[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r}\right) \left(k_\nu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s}\right) \\
&\qquad\qquad \times \ln \left\{ \frac{\prod_{n=1}^\infty \left(1 - e^{i(\sigma_r - \sigma_s)} q_c^n\right) \left(1 - e^{-i(\sigma_r - \sigma_s)} q_c^n\right)}{\prod_{n=1}^\infty \left(1 - q_c^n\right)^2} \right\} \Bigg] \\
&\quad \times \prod_{r=1}^{M} \prod_{s=M+1}^{M+N} \exp\Bigg[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r}\right) \left(k_\nu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s}\right)
\end{aligned}
$$

$$\times \ln \left\{ \frac{\prod_{m=0}^{\infty} \left(1 - e^{i(\sigma_r - \sigma_s)} q_c^{m+\frac{1}{2}}\right) \left(1 - e^{-i(\sigma_r - \sigma_s)} q_c^{m+\frac{1}{2}}\right)}{\prod_{n=1}^{\infty} (1 - q_c^n)^2} \right\} \Bigg]$$

$$\times \exp \left[2\alpha' G^{\mu\nu} \sum_{r=1}^{M+N} (a_r)^2 A_\mu(k^{(r)}) A_\nu(k^{(r)}) \ \ln \left\{ \prod_{n=1}^{\infty} (1 - q_c^n) \right\} \right] \ . \tag{7.39}$$

We have used the total momentum conservation, $\delta^{(p+1)} \left(\sum_{r=1}^{M+N} k^{(r)} \right)$ to obtain the above expression. The last exponential will be ignored since the exponent is proportional to $(a_r)^2$ and this term brings about nothing when $\prod_{r=1}^{M+N} \frac{\partial}{\partial a_r} \Big|_{a_r=0}$ operate on the amplitudes.

The amplitudes which are obtained by plugging Eq.(7.39) into Eq.(7.37) may be written down by using the elliptic θ-functions. With a similar manipulation which leads Eq.(3.50) we can recast the amplitudes into the following forms :

$$\begin{aligned}
&\Big\langle \hat{B}_N[A]; \{a_r\}; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \Big| \\
&\qquad\qquad \times q_c^{\frac{1}{2}\left(L_0 + \tilde{L}_0 - 2\right)} \Big| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \Big\rangle \\
&= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{2}} \ q_c^{\frac{p-25}{24}} \eta\left(\tau^{(c)}\right)^{-p+1} \ \delta^{(p+1)} \left(\sum_{r=1}^{M+N} k^{(r)} \right) \\
&\quad \times \prod_{1 \le r < s \le M} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \prod_{M+1 \le r < s \le M+N} e^{-\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)} \epsilon(\sigma_r - \sigma_s)} \\
&\quad \times q_c^{-\frac{1}{16\pi^2 \alpha'} K_\mu (\theta G \theta)^{\mu\nu} K_\nu} \prod_{r=1}^{M+N} e^{\frac{i}{2\pi} \left(a_r A_\mu(k^{(r)}) - \sigma_r k_\mu^{(r)} \right) \theta^{\mu\nu} K_\nu} \\
&\quad \times \prod_{1 \le r < s \le M} \exp \Bigg[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \\
&\qquad\qquad\qquad \times \left(k_\mu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s} \right) \ln \left\{ \frac{\theta_1 \left(\frac{|\sigma_r - \sigma_s|}{2\pi} \middle| \tau^{(c)} \right)}{\eta\left(\tau^{(c)}\right)^3} \right\} \Bigg] \\
&\quad \times \prod_{M+1 \le r < s \le M+N} \exp \Bigg[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \\
&\qquad\qquad\qquad \times \left(k_\mu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s} \right) \ln \left\{ \frac{\theta_1 \left(\frac{|\sigma_r - \sigma_s|}{2\pi} \middle| \tau^{(c)} \right)}{\eta\left(\tau^{(c)}\right)^3} \right\} \Bigg] \\
&\quad \times \prod_{r=1}^{M} \prod_{s=M+1}^{M+N} \exp \Bigg[2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \\
&\qquad\qquad\qquad \times \left(k_\mu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s} \right) \ln \left\{ \frac{\theta_4 \left(\frac{\sigma_r - \sigma_s}{2\pi} \middle| \tau^{(c)} \right)}{\eta\left(\tau^{(c)}\right)^3} \right\} \Bigg] \ ,
\end{aligned}$$

$$\tag{7.40}$$

where we put $K_\mu \equiv \sum_{r=1}^{M} k_\mu^{(r)}$. In the above expression we have included the contribution of the ghosts, $\prod_{n=1}^{\infty}(1-q_c^n)^2$.

7.3 Comparison with open-string one-loop calculation

We compare the amplitudes (7.40), taking account of closed-string propagations along the Dirichlet directions, with open-string gluon one-loop amplitudes. The discussion goes parallel to the case of the tachyon. We first describe the open-string calculation. We use the same conventions as in subsection 3.4.

7.3.1 Open-string one-loop calculation

Open-string gluon vertex operator of momentum k is given by

$$V_A^{open}(\xi,\sigma;k) \equiv iA_\mu(k) : (\partial_\rho + \overline{\partial}_\rho)X^\mu(\rho,\bar\rho)e^{ik_\nu X^\nu(\rho,\bar\rho)} :\Big|_{\rho=\xi e^{i\sigma}} , \tag{7.41}$$

where ρ and (ξ,σ) are respectively the complex and the radial coordinates of the upper half-plane (open-string world sheet). $A_\mu(k)$ are the polarization vectors or the Fourier modes of the $U(1)$ gauge field $\mathcal{A}_\mu(x)$. We consider the scattering process of $M+N$ gluons with momenta $k^{(r)}$. Diagram relevant to the one-loop scattering process can be drawn on the upper half-plane as depicted in Figure 3. The corresponding gluon amplitude, which we call $I_A\left((k^{(1)},\cdots,k^{(M)});(k^{(M+1)},\cdots,k^{(M+N)})\right)$, is given by a sum of traces of their products arranged in cyclically distinct orders with keeping their partial orderings at the each end :

$$\begin{aligned} & I_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\ &\quad \equiv \mathrm{Tr}\left\{\Delta V_A^{open}(1,0;k^{(1)})\cdots\Delta V_A^{open}(1,0;k^{(M)})\right. \\ &\qquad\qquad \left.\times\ \Delta V_A^{open}(1,\pi;k^{(M+1)})\cdots\Delta V_A^{open}(1,\pi;k^{(M+N)})\right\} \\ &\qquad + \quad \cdots\cdots . \end{aligned} \tag{7.42}$$

In order to obtain the above amplitude it becomes convenient to introduce the gluon vertex operator in an exponential form. Let $a \in \mathbb{R}$ be an auxiliary parameter. We put [41]

$$\hat V_A^{open}(\xi,\sigma;k;a) \equiv\ : \exp\left\{ik_\mu X^\mu(\rho,\bar\rho) + iaA_\mu(k)(\partial_\rho+\overline{\partial}_\rho)X^\mu(\rho,\bar\rho)\right\} :\Big|_{\rho=\xi e^{i\sigma}} . \tag{7.43}$$

This operator is related with the vertex operator (7.41) by

$$V_A^{open}(\xi,\sigma;k) = \frac{\partial}{\partial a}\hat V_A^{open}(\xi,\sigma;k;a)\bigg|_{a=0} . \tag{7.44}$$

The corresponding amplitude constructed from $\hat{V}_A^{open}$ instead of V_A^{open} will be called $\hat{I}_A$. It is given by

$$\begin{aligned} &\hat{I}_A\left(\left(\left(k^{(1)},a_1\right),\cdots,\left(k^{(M)},a_M\right)\right);\left(\left(k^{(M+1)},a_{M+1}\right),\cdots,\left(k^{(M+N)},a_{M+N}\right)\right)\right) \\ &\quad\equiv \mathrm{Tr}\left\{\Delta\hat{V}_A^{open}\left(1,0;k^{(1)};a_1\right)\cdots\Delta\hat{V}_A^{open}\left(1,0;k^{(M)};a_M\right)\right. \\ &\qquad\qquad\left.\times\,\Delta\hat{V}_A^{open}\left(1,\pi;k^{(M+1)};a_{M+1}\right)\cdots\Delta\hat{V}_A^{open}\left(1,\pi;k^{(M+N)};a_{M+N}\right)\right\} \\ &\qquad+\quad\cdots\cdots\,. \end{aligned} \tag{7.45}$$

The gluon scattering amplitude I_A can be obtained from $\hat{I}_A$ by differentiating it with respect to a_r and making them vanish:

$$\begin{aligned} &I_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\ &=\prod_{r=1}^{M+N}\left.\frac{\partial}{\partial a_r}\right|_{a_r=0}\hat{I}_A\left(\left(\left(k^{(1)},a_1\right),\cdots,\left(k^{(M)},a_M\right)\right);\left(\left(k^{(M+1)},a_{M+1}\right),\cdots,\left(k^{(M+N)},a_{M+N}\right)\right)\right)\,. \end{aligned} \tag{7.46}$$

Due to the existence of the auxiliary parameter the Virasoro algebra acts on the auxiliary gluon vertex operator $\hat{V}_A^{open}$ in an unfamiliar form. The generators have been given in (3.56). Their action turns out to be as follows : For $n\in\mathbb{Z}$

$$\begin{aligned} &[L_n,\hat{V}_A^{open}(\xi,0;k;a)] \\ &=\left\{\xi^{n+1}\frac{\partial}{\partial\xi}+(n+1)\alpha' k_\mu G^{\mu\nu}k_\nu\xi^n+(n+1)\xi^n a\frac{\partial}{\partial a}\right. \\ &\qquad+n(n+1)a\alpha' k_\mu G^{\mu\nu}A_\nu(k)\xi^{n-1} \\ &\qquad\qquad\left.+\frac{n(n^2-1)}{6}a^2\alpha' A_\mu(k)G^{\mu\nu}A_\nu(k)\xi^{n-2}\right\}\hat{V}_A^{open}(\xi,0;k;a), \end{aligned} \tag{7.47}$$

and the same form on $\hat{V}_A^{open}(\xi,\pi;k;a)$ except replacing ξ by $-\xi$ in the RHS. By applying the relation (7.44) to the above equation we obtain the standard Virasoro action on the gluon vertex operators. The auxiliary operators become the tachyon vertex operators simply by letting a vanish in (7.43), and the above action reduces to the standard action on the tachyon vertex operators.

We can evaluate the amplitude $\hat{I}_A$ in the standard manner. The open-string propagators in (7.45) may be replaced by the integral forms. The above Virasoro action implies :

$$\begin{aligned} \xi^{L_0}\hat{V}_A^{open}(1,0;k;a)\xi^{-L_0} &= \xi^{\alpha' k_\mu G^{\mu\nu}k_\nu}\hat{V}_A^{open}(\xi,0;k;\xi a), \\ \xi^{L_0}\hat{V}_A^{open}(1,\pi;k)\xi^{-L_0} &= \xi^{\alpha' k_\mu G^{\mu\nu}k_\nu}\hat{V}_A^{open}(\xi,\pi;k;\xi a). \end{aligned} \tag{7.48}$$

Utilizing these properties we can write the amplitude in the following form :

$$
\begin{aligned}
&\hat{I}_A\left(\left(\left(k^{(1)}, a_1\right), \cdots, \left(k^{(M)}, a_M\right)\right); \left(\left(k^{(M+1)}, a_{M+1}\right), \cdots, \left(k^{(M+N)}, a_{M+N}\right)\right)\right) \\
&= \int_{0 \le \xi_{M+N} \le \xi_{M+N-1} \le \cdots \le \xi_1 \le 1} \prod_{r=1}^{M+N} \frac{d\xi_r}{\xi_r} \times \prod_{r=1}^{M+N} \xi_r^{\alpha' k_\mu^{(r)} G_{\mu\nu} k_\nu^{(r)}} \\
&\quad \times \mathrm{Tr}\left\{ \hat{V}_A^{open}\left(\xi_1, 0; k^{(1)}; \xi_1 a_1\right) \cdots \hat{V}_A^{open}\left(\xi_M, 0; k^{(M)}; \xi_M a_M\right) \right. \\
&\qquad \left. \times \hat{V}_A^{open}\left(\xi_{M+1}, \pi; k^{(M+1)}; \xi_{M+1} a_{M+1}\right) \cdots \hat{V}_A^{open}\left(\xi_{M+N}, \pi; k^{(M+N)}; \xi_{M+N} a_{M+N}\right) \xi_{M+N}^{L_0 - 1} \right\} \\
&\quad + \quad \cdots\cdots .
\end{aligned}
\tag{7.49}
$$

The coordinates ξ_r in the RHS are insertion points of the auxiliary gluon vertices and provide a parametrization of the diagram. Another parametrization can be obtained by mapping the diagram to the cylinder with width π as depicted in Figure 4. Correspondingly ξ_r are mapped to ν_r by $\nu_r = -\ln \xi_r$. We put $\tau^{(o)} \equiv -\ln \xi_{M+N}$. These are the open-string parameters.

To evaluate the RHS of Eq.(7.49), we first normal-order the products of the auxiliary operators within the traces by taking account of their OPEs. For instance we use

$$
\begin{aligned}
&\hat{V}_A^{open}\left(\xi_1, 0; k^{(1)}; a_1\right) \hat{V}_A^{open}\left(\xi_2, 0; k^{(2)}; a_2\right) \\
&\quad = e^{-\frac{i}{2} k_\mu^{(1)} \theta^{\mu\nu} k_\nu^{(2)}} \times e^{2\alpha' G^{\mu\nu} \left(k_\mu^{(1)} + a_1 A_\mu(k^{(1)}) \frac{\partial}{\partial \xi_1}\right)\left(k_\nu^{(2)} + a_2 A_\nu(k^{(2)}) \frac{\partial}{\partial \xi_2}\right) \ln(\xi_1 - \xi_2)} \\
&\qquad\qquad \times : \hat{V}_A^{open}\left(\xi_1, 0; k^{(1)}; a_1\right) \hat{V}_A^{open}\left(\xi_2, 0; k^{(2)}; a_2\right) : \quad , \\
&\hat{V}_A^{open}\left(\xi_1, 0; k^{(1)}; a_1\right) \hat{V}_A^{open}\left(\xi_2, \pi; k^{(2)}; a_2\right) \\
&\quad = e^{-\frac{i}{2} k_\mu^{(1)} \theta^{\mu\nu} k_\nu^{(2)}} \times e^{2\alpha' G^{\mu\nu} \left(k_\mu^{(1)} + a_1 A_\mu(k^{(1)}) \frac{\partial}{\partial \xi_1}\right)\left(k_\nu^{(2)} - a_2 A_\nu(k^{(2)}) \frac{\partial}{\partial \xi_2}\right) \ln(\xi_1 + \xi_2)} \\
&\qquad\qquad \times : \hat{V}_A^{open}\left(\xi_1, 0; k^{(1)}; a_1\right) \hat{V}_A^{open}\left(\xi_2, \pi; k^{(2)}; a_2\right) : \quad ,
\end{aligned}
\tag{7.50}
$$

for $\xi_1 > \xi_2$. We still need to take traces of these normal-ordered operators. After straightforward but wearisome calculations the amplitude can be written down in terms of the open-string parameters. It turns out to be the following integral :

$$
\begin{aligned}
&\hat{I}_A\left(\left(\left(k^{(1)}, a_1\right), \cdots, \left(k^{(M)}, a_M\right)\right); \left(\left(k^{(M+1)}, a_{M+1}\right), \cdots, \left(k^{(M+N)}, a_{M+N}\right)\right)\right) \\
&= \sqrt{-\mathrm{det} G_{\mu\nu}}\, \delta^{(p+1)}\left(\sum_{r=1}^{M+N} k^{(r)}\right) \\
&\quad \times \prod_{1 \le r < s \le M} e^{-\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)}} \prod_{M+1 \le r < s \le M+N} e^{\frac{i}{2} k_\mu^{(r)} \theta^{\mu\nu} k_\nu^{(s)}} \\
&\quad \times \int d\tau^{(o)} \prod_{r=1}^{M+N-1} d\nu_r \left(\frac{\pi}{\alpha' \tau^{(o)}}\right)^{\frac{p+1}{2}} e^{\frac{26-D}{24} \tau^{(o)}} \eta\left(-\frac{\tau^{(o)}}{2\pi i}\right)^{-D+2} \times e^{\frac{K_\mu (\theta G \theta)^{\mu\nu} K_\nu}{4\alpha' \tau^{(o)}}}
\end{aligned}
$$

$$
\begin{aligned}
&\times \prod_{r=1}^{M} \prod_{s=M+1}^{M+N} \exp\left[2\alpha' G^{\mu\nu}\left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\frac{\partial}{\partial\nu_r}\right)\left(k_\nu^{(s)} + a_s A_\nu(k^{(s)})\frac{\partial}{\partial\nu_s}\right)\right.\\
&\qquad\qquad \left.\times \ln\left\{e^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\left.\frac{(\nu_s+i\pi)-\nu_r}{2\pi i}\right|\frac{-\tau^{(o)}}{2\pi i}\right)\right\}\right]\\
&\times \prod_{r=1}^{M}\exp\left\{-a_r\alpha' k_\mu^{(r)} G^{\mu\nu} A_\nu(k^{(r)})\right\}\prod_{r=1}^{M}\exp\left\{\frac{i}{\tau^{(o)}}K_\mu\theta^{\mu\nu}\left(k_\nu^{(r)}\nu_r - a_r A_\nu(k^{(r)})\right)\right\}\\
&\times \prod_{1\le r<s\le M}\exp\left[2\alpha' G^{\mu\nu}\left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\frac{\partial}{\partial\nu_r}\right)\left(k_\nu^{(s)} - a_s A_\nu(k^{(s)})\frac{\partial}{\partial\nu_s}\right)\right.\\
&\qquad\qquad \left.\times \ln\left\{ie^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\left.\frac{\nu_s-\nu_r}{2\pi i}\right|\frac{-\tau^{(o)}}{2\pi i}\right)\right\}\right]\\
&\times \prod_{r=M+1}^{M+N}\exp\left\{a_r\alpha' k_\mu^{(r)} G^{\mu\nu} A_\nu(k^{(r)})\right\}\prod_{r=M+1}^{M+N}\exp\left\{\frac{i}{\tau^{(o)}}K_\mu\theta^{\mu\nu}\left(k_\nu^{(r)}\nu_r + a_r A_\nu(k^{(r)})\right)\right\}\\
&\times \prod_{M+1\le r<s\le M+N}\exp\left[2\alpha' G^{\mu\nu}\left(k_\mu^{(r)} + a_r A_\mu(k^{(r)})\frac{\partial}{\partial\nu_r}\right)\left(k_\nu^{(s)} + a_s A_\nu(k^{(s)})\frac{\partial}{\partial\nu_s}\right)\right.\\
&\qquad\qquad \left.\times \ln\left\{ie^{-\frac{(\nu_s-\nu_r)^2}{2\tau^{(0)}}}\eta\left(\frac{-\tau^{(o)}}{2\pi i}\right)^{-3}\theta_1\left(\left.\frac{\nu_s-\nu_r}{2\pi i}\right|\frac{-\tau^{(o)}}{2\pi i}\right)\right\}\right]\ ,
\end{aligned}
\tag{7.51}
$$

where the integration of the open-string parameters are performed over the region (3.63), and K_μ is given by $K_\mu = \sum_{r=1}^{M} k_\mu^{(r)}$. Contribution of the world-sheet reparametrization ghosts has been included in the above.

In order to compare the amplitude (7.51) with Eq.(7.40) we map the cylinder drawn on the u-plane (Figure 4) to the cylinder on the v-plane (Figure 5) by conformal transformation $v = 2\pi u/\tau^{(o)}$. ν_r are mapped to σ_r by $\sigma_r = 2\pi\nu_r/\tau^{(o)}$. We put $\tau^{(c)} \equiv 2\pi i/\tau^{(o)}$. These are the closed-string parameters. We note that an insertion point of the $(M+N)$-th auxiliary gluon vertex operator is fixed at $\sigma_{M+N} = 2\pi$. By making use of the modular transformations (3.64) the above one-loop amplitude can be described as an integral with respect to $\tau^{(c)}$ and σ_r. Subsequently we use the relation (7.46). An integral form of the $M+N$ gluon scattering amplitude is obtained in terms of the closed-string parameters. This turns out to be as follows :

$$
\begin{aligned}
&I_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right)\\
&=\left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{1/2}\delta^{(p+1)}\left(\sum_{r=1}^{M+N}k^{(r)}\right)
\end{aligned}
$$

$$
\begin{aligned}
&\times \prod_{1\le r<s\le M} e^{-\frac{i}{2}k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu} \prod_{M+1\le r<s\le M+N} e^{\frac{i}{2}k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu} \\
&\times (-2\pi i) \prod_{r=1}^{M+N} \left.\frac{\partial}{\partial a_r}\right|_{a_r=0} \int_{i0}^{i\infty} d\tau^{(c)} \int \prod_{r=1}^{M+N-1} d\sigma_r \; e^{\frac{26-D}{24}\frac{2\pi i}{\tau^{(c)}}} \left(-i\tau^{(c)}\right)^{-\frac{d}{2}} \eta(\tau^{(c)})^{-D+2} \\
&\times e^{\frac{\tau^{(c)}}{8\pi i\alpha'} K_\mu (\theta G\theta)^{\mu\nu} K_\nu} \times \prod_{r=1}^{M+N} (-i\tau^{(c)})^{\alpha' k^{(r)}_\mu G^{\mu\nu} k^{(r)}_\nu} \\
&\times \prod_{r=1}^{M} \prod_{s=M+1}^{M+N} \exp\Bigg[2\alpha' G^{\mu\nu} \left(k^{(r)}_\mu - a_r A_\mu(k^{(r)}) \frac{\partial}{\partial\sigma_r} \right) \left(k^{(s)}_\nu + a_s A_\nu(k^{(s)}) \frac{\partial}{\partial\sigma_s} \right) \\
&\qquad\qquad\qquad \times \ln\left\{ \eta\left(\tau^{(c)}\right)^{-3} \theta_4\left(\left.\frac{\sigma_s-\sigma_r}{2\pi i}\right| \tau^{(c)} \right) \right\} \Bigg] \\
&\times \prod_{r=1}^{M} \exp\left\{ -\frac{a_r}{(-i\tau^{(c)})} \alpha' k^{(r)}_\mu G^{\mu\nu} A_\nu(k^{(r)}) \right\} \prod_{r=1}^{M} \exp\left\{ \frac{i}{2\pi} K_\mu \theta^{\mu\nu} \left(k^{(r)}_\nu \sigma_r - a_r A_\nu(k^{(r)}) \right) \right\} \\
&\times \prod_{1\le r<s\le M} \exp\Bigg[2\alpha' G^{\mu\nu} \left(k^{(r)}_\mu - a_r A_\mu(k^{(r)}) \frac{\partial}{\partial\sigma_r} \right) \left(k^{(s)}_\nu - a_s A_\nu(k^{(s)}) \frac{\partial}{\partial\sigma_s} \right) \\
&\qquad\qquad\qquad \times \ln\left\{ \eta\left(\tau^{(c)}\right)^{-3} \theta_1\left(\left.\frac{\sigma_s-\sigma_r}{2\pi}\right| \tau^{(c)} \right) \right\} \Bigg] \\
&\times \prod_{r=M+1}^{M+N} \exp\left\{ \frac{a_r}{(-i\tau^{(c)})} \alpha' k^{(r)}_\mu G^{\mu\nu} A_\nu(k^{(r)}) \right\} \prod_{r=M+1}^{M+N} \exp\left\{ \frac{i}{2\pi} K_\mu \theta^{\mu\nu} \left(k^{(r)}_\nu \sigma_r + a_r A_\nu(k^{(r)}) \right) \right\} \\
&\times \prod_{M+1\le r<s\le M+N} \exp\Bigg[2\alpha' G^{\mu\nu} \left(k^{(r)}_\mu + a_r A_\mu(k^{(r)}) \frac{\partial}{\partial\sigma_r} \right) \left(k^{(s)}_\nu + a_s A_\nu(k^{(s)}) \frac{\partial}{\partial\sigma_s} \right) \\
&\qquad\qquad\qquad \times \ln\left\{ \eta\left(\tau^{(c)}\right)^{-3} \theta_1\left(\left.\frac{\sigma_s-\sigma_r}{2\pi}\right| \tau^{(c)} \right) \right\} \Bigg],
\end{aligned}
\tag{7.52}
$$

where the integration of the closed-string parameters are performed over the region (3.66). And we have rescaled the auxiliary parameters a_r to $(-i\tau^{(c)})a_r$.

The above integral form allows us to express the one-loop amplitude of $M+N$ gluons by using the closed-string amplitudes given in (7.40). Including the contribution of closed-string propagations along the Dirichlet directions we finally obtain the following equality :

$$
\begin{aligned}
&I_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\
&= 2\pi i(-)^{N+1} \prod_{r=1}^{M+N} \left.\frac{\partial}{\partial a_r}\right|_{a_r=0} \int_{i0}^{i\infty} d\tau^{(c)} \; e^{2\pi i\left(\tau^{(c)}+\frac{1}{\tau^{(c)}}\right)\frac{26-D}{24}} \prod_{r=1}^{M+N} (-i\tau^{(c)})^{\alpha' k^{(r)}_\mu G^{\mu\nu} k^{(r)}_\nu} \\
&\qquad \times \prod_{r=1}^{M} \exp\left\{ -\frac{a_r}{(-i\tau^{(c)})} \alpha' k^{(r)}_\mu G^{\mu\nu} A_\nu(k^{(r)}) \right\} \prod_{r=M+1}^{M+N} \exp\left\{ \frac{a_r}{(-i\tau^{(c)})} \alpha' k^{(r)}_\mu G^{\mu\nu} A_\nu(k^{(r)}) \right\}
\end{aligned}
$$

$$\times\left[\int\prod_{r=M+1}^{M+N-1}d\sigma_r\,\langle B_D|\otimes\left\langle\hat{B}_N[A];\{a_r\};(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right|\right]$$
$$\times\, q_c^{\frac{L_0+\tilde{L}_0-2}{2}}\left[\int\prod_{r=1}^{M}d\sigma_r\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle\otimes|B_D\rangle\right]\;, \tag{7.53}$$

where the integration is performed over the region (3.66). All the additional terms appearing in the RHS become completely vanishing when the on-shell conditions (7.20) are imposed on the gluon vertices at the critical dimensions.

8 UV limit of Non-commutative Gauge Theory

Low-energy world-volume theory of open-string gluons in the presence of a constant B field is a $U(1)$ gauge theory on the non-commutative world-volume. In this section, taking the same route as the previous study of the non-commutative scalar field theory, we investigate the UV behavior of the non-commutative gauge theory. Our study in this section is restricted to the case of 25-brane in the critical dimensions. We examine two zero-slope limits of the one-loop amplitudes of gluons. One is based on the open-string parameters and the other is on the closed-string parameters. These two limits, as we explained in the study of the non-commutative scalar field theory, are complementary.

We first take a zero-slope limit based on the open-string parameters $(\tau^{(o)},\nu_r)$. Eq.(7.51) may be used as an integral form of the amplitude. Strictly speaking, the amplitude is obtained from this integral by using the relation (7.46). The zero-slope limit will be a gauge theory one-loop amplitude, particularly written in terms of the Schwinger parameters, $s^{(o)}$ and T_r. These parameters are related with $\tau^{(o)}$ and ν_r by the relations (4.1). The auxiliary parameters a_r are also used in the amplitude (7.51). They are introduced in order to describe the gluon vertex operators (7.41) in the auxiliary forms (7.43) and to make the loop calculation tractable. At the zero-slope limit the gluon vertex operator become $iA_\mu(k)\frac{dx^\mu(s)}{ds}e^{ik_\nu x^\nu(s)}$, where $x(s)$ is a world-line parametrized by the Schwinger parameter s. The auxiliary vertex operator is expected to be $e^{i\left(k_\mu x^\mu(s)+\kappa A_\mu(k)\frac{dx^\mu(s)}{ds}\right)}$ in the world-line description. Here κ is an auxiliary parameter (a counterpart of a in the field theory). A simple dimensional analysis shows that κ is dimensionful and proportional to $\alpha' a$. Therefore the zero-slope limit must be taken by fixing the following field theory parameters in the amplitude :

$$s^{(o)}=\alpha'\tau^{(o)},\quad T_r=\alpha'\nu_r,\quad \kappa_r=\alpha' a_r. \tag{8.1}$$

Simultaneously we also need to fix open-string tensors $G_{\mu\nu}$ and $\theta^{\mu\nu}$. We rewrite the amplitude (7.51) in terms of the above parameters and then pick up the dominant contribution of the α'-expansion. These are parallel to what we did in the previous section to obtain the zero-slope limit (4.3) of the tachyon amplitude. By using the relation (7.46), we find that the following integral turns out to be the zero-slope limit :

$$
\begin{aligned}
& I_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\
& \approx \pi^{13}\sqrt{-\det G_{\mu\nu}}\,\delta^{26}\left(\sum_{r=1}^{M+N}k^{(r)}\right)\prod_{1\le r<s\le M}e^{-\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}}\prod_{M+1\le r<s\le M+N}e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}} \\
& \quad \times\prod_{r=1}^{M+N}\left.\frac{\partial}{\partial\kappa_r}\right|_{\kappa_r=0}\int ds^{(o)}(s^{(o)})^{-13}\exp\left\{\frac{s^{(o)}}{\alpha'}+\frac{K_\mu(\theta G\theta)^{\mu\nu}K_\nu}{4s^{(o)}}\right\}\int\prod_{r=1}^{M+N-1}dT_r \\
& \qquad \times\prod_{r=1}^{M+N}\exp\left\{-\epsilon^{(r)}\kappa_r k_\mu^{(r)}G^{\mu\nu}A_\nu(k^{(r)})\right\} \\
& \qquad \times\prod_{r=1}^{M+N}\exp\left\{-\frac{i}{s^{(0)}}\left(T_r k_\mu^{(r)}-\epsilon^{(r)}\kappa_r A_\mu(k^{(r)})\right)\theta^{\mu\nu}K_\nu\right\} \\
& \qquad \times\prod_{r<s}^{M+N}\exp\left[G^{\mu\nu}\left(k_\mu^{(r)}-\epsilon^{(r)}\kappa_r A_\mu(k^{(r)})\frac{\partial}{\partial T_r}\right)\left(k_\nu^{(s)}-\epsilon^{(s)}\kappa_s A_\nu(k^{(s)})\frac{\partial}{\partial T_s}\right)\right. \\
& \qquad\qquad\qquad\qquad \left.\times\left\{-\frac{(T_r-T_s)^2}{s^{(o)}}+|T_s-T_r|\right\}\right], \qquad (8.2)
\end{aligned}
$$

where $\epsilon^{(r)}$ take values ± 1 such that $+1$ for $1\le r\le M$ and -1 for $M+1\le r\le M+N$, and the integral is performed over the region (4.4). If we neglect κ_r besides their differentiations the above zero-slope limit reduces to Eq.(4.3), modulo the factor α'^{-M-N}. This power has a simple origin in the scaling relations, $\kappa_r=\alpha' a_r$. It was discussed previously that $\exp\left\{\frac{s^{(o)}}{\alpha'}\right\}$ in the integral comes from the Schwinger representation of the open-string tachyon propagator, $\exp\left\{-s^{(o)}(p^2-\frac{1}{\alpha'})\right\}$. Presently the same term may be interpreted as an IR regularization of the amplitude by an analytic continuation of $-\frac{1}{\alpha'^2}$ to a small positive m^2.

The above zero-slope limit can be identified with the corresponding one-loop amplitude of the non-commutative $U(1)$ gauge theory. Some related calculation in the gauge theory may be found in [42][43] [8]. Similarly to the scalar field theory, $\exp\left\{\frac{K_\mu(\theta G\theta)^{\mu\nu}K_\nu}{4s^{(o)}}\right\}$ in the above integral is understood as a UV regularization [11] which depends on the external momentum K. In the gauge theory as well, this causes the problem of UV-IR mixing and makes the field theory description at high energy scale difficult.

---

[8] Eq.(8.2) may be compared with Eq.(4.9) of [43]. The integration measure used in that paper is different from ours.

As explained in Section 4, the open- and closed-string parameters used in the descriptions of the amplitude are two kinds of coordinates of the moduli space of conformal classes of cylinder with $M+N$ punctures at the boundaries. Two ends of the moduli space which are located at $\tau^{(o)} = 0$ and $\tau^{(o)} = +\infty$ play important roles in the zero-slope limits. The field theory amplitude (8.2) is obtained by a suitable magnification of the integral (7.51) on an infinitesimal neighbourhood around $\tau^{(o)} = +\infty$. We need to focus on the region $s^{(o)} \approx 0$ to know the UV behavior of the gauge theory. As is the case of the scalar field theory, a possible resolution is to take a zero-slope limit such that an infinitesimal neighbourhood around $\tau^{(o)} = 0$ is magnified. It becomes effective to use the closed-string parameters near this end. The zero-slope limit which we examine is essentially the same as that was examined for the open-string tachyons. Explicitly we take the limit by fixing the following parameters :

$$s^{(c)} = \alpha' |\tau^{(c)}|, \quad \sigma_r, \quad a_r. \tag{8.3}$$

In the limiting process we fix open-string tensors $G_{\mu\nu}$ and $\theta^{\mu\nu}$ as well to capture the world-volume theory. The auxiliary parameters in the above are the rescaled ones used in the integral (7.52). In contrast with the previous scaling we keep them intact. This is because the present limit is not expected to allow a naive world-line description. As can be observed in the expression (7.53) these parameters are nothing but the auxiliary parameters used for the description of gluon vertex operators in the boundary state formalism.

Relevant integral form of the one-loop amplitude is given by Eq.(7.52) or equivalently Eq.(7.53) using the boundary states. We slightly generalize the amplitude by dropping out the factors $\prod_{r=1}^{M+N} (-i\tau^{(c)})^{\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(r)}} e^{-\frac{a_r \epsilon^{(r)}}{(-i\tau^{(c)})}\alpha' k_\mu^{(r)} G^{\mu\nu} A_\nu(k^{(r)})}$ and introduce the following quantity :

$$\begin{aligned}
& J_A\left(\left(k^{(1)}, \cdots, k^{(M)}\right); \left(k^{(M+1)}, \cdots, k^{(M+N)}\right)\right) \\
&= 2\pi i (-)^{N+1} \int_{i0}^{i\infty} d\tau^{(c)} \prod_{r=1}^{M+N} \left.\frac{\partial}{\partial a_r}\right|_{a_r=0} \\
&\qquad \times \left[\int \prod_{r=M+1}^{M+N-1} d\sigma_r \, \langle B_D| \otimes \left\langle \hat{B}_N[A]; \{a_r\}; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right|\right] \\
&\qquad \times q_c^{\frac{L_0+\tilde{L}_0-2}{2}} \left[\int \prod_{r=1}^{M} d\sigma_r \left|\hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \otimes |B_D\rangle\right] ,
\end{aligned} \tag{8.4}$$

where the integration is performed on the region (3.66). The $U(1)$ symmetry of the one-loop diagram is fixed by putting $\sigma_{M+N} = 2\pi$. When the gluon vertices satisfy the on-shell conditions the above J_A coincides with the original amplitude. It is an off-shell generalization slightly different from the original one.

The zero-slope limit can be obtained by a calculation parallel to that of open-string tachyons. The operations $\prod_{r=1}^{M+N} \frac{\partial}{\partial a_r}\Big|_{a_r=0}$ are carried out on the dominant term of the α'-expansion without difficulty. The following integral turns out to be the zero-slope limit :

$$\begin{aligned}
& J_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(k^{(M+1)},\cdots,k^{(M+N)}\right)\right) \\
& \approx \frac{2\pi}{\alpha'}\delta^{26}\left(\sum_{r=1}^{M+N} k^{(r)}\right)\int ds^{(c)} \exp\left\{-\frac{\pi s^{(c)}}{2}\left(K_\mu g^{\mu\nu}K_\nu - \frac{4}{\alpha'}\right)\right\} \\
& \quad \times \left[\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{1/4}\int \prod_{r=1}^{M} d\sigma_r \prod_{1\le r<s\le M} e^{-\frac{i}{2}k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu}\right. \\
& \qquad\qquad \left.\times \prod_{r=1}^{M}\left\{-\frac{i}{2\pi}K_\mu\theta^{\mu\nu}A_\nu(k^{(r)})e^{-ik^{(r)}_\mu\theta^{\mu\nu}K_\nu\frac{\sigma_r}{2\pi}}\right\}\right] \\
& \quad \times \left[\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{1/4}\int \prod_{r=M+1}^{M+N-1} d\sigma_r \prod_{M+1\le r<s\le M+N} e^{\frac{i}{2}k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu}\right. \\
& \qquad\qquad \left.\times \prod_{r=M+1}^{M+N}\left\{\frac{i}{2\pi}K_\mu\theta^{\mu\nu}A_\nu(k^{(r)})e^{-ik^{(r)}_\mu\theta^{\mu\nu}K_\nu\frac{\sigma_r}{2\pi}}\right\}\right] \ ,
\end{aligned} \tag{8.5}$$

where the integral is performed over the region (4.11) and the closed-string metric is understood as $g^{\mu\nu} \approx -\frac{(\theta G\theta)^{\mu\nu}}{(2\pi\alpha')^2}$.

As was pointed out in the previous study of the scalar field theory, the zero-slope limit based on the closed-string parameters describes physics at the trans-string scale of the world-volume theory. All the perturbative stringy states of open-string contribute to the limit. They bring about a striking contrast between the two limits. We can find the propagator of closed-string tachyon of momentum K in the limit (8.5) while the counterpart in (8.2) is the curious regularization factor. In that limit there appear terms describing correlations between two gluons inserted at T_r and T_s. These correlations are caused by the kinetic energies of the propagating gluons. But they are lost in (8.5) and gluons become topological in this limit. The disappearance originates in the modular transforms (3.64).

It is also possible to rewrite the limit (8.5) in terms of the Moyal products. The translation can be made in the same manner as we translated the tachyon amplitude (4.10) into (4.15). We obtain

$$\int \prod_{r=1}^{M+N} \frac{dk^{(r)}}{(2\pi)^{13}}(-)^{M+N} J_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(-k^{(M+1)},\cdots,-k^{(M+N)}\right)\right)$$

$$\begin{aligned}
&\approx \frac{2\pi}{\alpha'}\int dp_0\left[\int_0^\infty ds^{(c)}\exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)\right\}\right]\\
&\quad\times\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}}\int\frac{dx}{(2\pi)^{26}}e^{ip_{0\mu}x^{\mu}}\star\left[\int\prod_{r=1}^{M}d\sigma_r\left\{i\mathcal{A}_\mu(x_0(\sigma_1))\frac{dx_0^\mu(\sigma_1)}{d\sigma_1}\right\}\star\cdots\right.\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\left.\cdots\star\left\{i\mathcal{A}_\mu(x_0(\sigma_M))\frac{dx_0^\mu(\sigma_M)}{d\sigma_M}\right\}\right]\\
&\quad\times\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}}\overline{\int\frac{dx}{(2\pi)^{26}}e^{ip_{0\mu}x^{\mu}}\star\left[\int\prod_{r=M+1}^{M+N-1}d\sigma_r\left\{i\mathcal{A}_\mu(x_0(\sigma_{M+1}))\frac{dx_0^\mu(\sigma_{M+1})}{d\sigma_{M+1}}\right\}\star\cdots\right.}\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\overline{\left.\cdots\star\left\{i\mathcal{A}_\mu(x_0(\sigma_{M+N}))\frac{dx_0^\mu(\sigma_{M+N})}{d\sigma_{M+N}}\right\}\right]},
\end{aligned}\tag{8.6}$$

where the integrations on σ_r are performed over the region (4.11). And $x_0(\sigma)$ is the straight line given by $x_0^\mu(\sigma) = x^\mu + \theta^{\mu\nu}p_{0\nu}\frac{\sigma}{2\pi}$. This allows us to write down the generating function of the amplitudes in terms of open Wilson lines. By summing up the above equation with respect to M and N the zero-slope limit becomes as follows :

$$\begin{aligned}
&\sum_{M=0}^{\infty}\sum_{N=1}^{\infty}\int\prod_{r=1}^{M+N}\frac{dk^{(r)}}{(2\pi)^{13}}(-)^{M+N}J_A\left(\left(k^{(1)},\cdots,k^{(M)}\right);\left(-k^{(M+1)},\cdots,-k^{(M+N)}\right)\right)\\
&\approx \frac{2\pi}{\alpha'}\int dp_0\left[\int_0^\infty ds^{(c)}\exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)\right\}\right]\\
&\quad\times\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}}\int\frac{dx}{(2\pi)^{26}}e^{ip_{0\mu}x^{\mu}}\star\left[\mathcal{P}_\star e^{i\int_0^{2\pi}d\sigma\mathcal{A}_\mu(x_0(\sigma))\frac{dx_0^\mu}{d\sigma}(\sigma)}\right]\\
&\quad\times\left(\frac{-\mathrm{det}G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}}\overline{\int\frac{dx}{(2\pi)^{26}}e^{ip_{0\mu}x^{\mu}}\star\left[\mathcal{P}_\star e^{i\int_0^{2\pi}d\sigma\mathcal{A}_\mu(x_0(\sigma))\frac{dx_0^\mu}{d\sigma}(\sigma)}\right]\star\left\{i\mathcal{A}_\mu(x_0(2\pi))\frac{dx_0}{d\sigma}(2\pi)\right\}}.
\end{aligned}\tag{8.7}$$

The last term, which makes the equation asymmetric, appears owing to fixing the $U(1)$ symmetry of the open-string diagrams. Modulo this asymmetry the generating function is factorized at the zero-slope limit into a sum of products of straight open Wilson lines. Open Wilson lines with the same velocity $\frac{\theta^{\mu\nu}p_{0\nu}}{2\pi}$ interact with each other by exchanging closed-string tachyon with momentum $p_{0\mu}$.

9 Open Wilson Lines in Closed-String Theory (II)

9.1 Straight open Wilson line and closed-string tachyon

It was shown in [23] disk amplitudes of a closed-string tachyon scattering with arbitrary number of gluons turn out to generate a *straight* open Wilson line in Seiberg-Witten's zero-slope limit. The path is a straight line connecting x^μ and $x^\mu + \theta^{\mu\nu} p_{0\nu}$, where p_0 is the momentum of the tachyon. The displacement $\theta^{\mu\nu} p_{0\nu}$ is required by the gauge invariance [24]. In this section we first reproduce the above result by using the boundary states constructed in Section 7. Computations provided below become also helpful for our subsequent investigations of generically curved open Wilson lines.

Closed-string tachyon state with momentum $p_{0\mu}$ and its BPZ dual state are given by

$$|p_{0N}\rangle = \lim_{\tau\to-\infty} V_T(\sigma,\tau;p_0)|\mathbf{0}\rangle = e^{ip_{0\mu}\hat{x}_0^\mu}|\mathbf{0}\rangle\ , \quad \langle -p_{0N}| = \langle \mathbf{0}|e^{ip_{0\mu}\hat{x}_0^\mu}\ . \tag{9.1}$$

Disk amplitude of the closed-string tachyon scattering with M gluons is obtained in the boundary state formalism by integrating the overlap $\left\langle -p_{0N}\left| B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right.\right\rangle$ on its moduli space. To evaluate this overlap, we start by computing the overlap with the auxiliary boundary states. This becomes

$$\begin{aligned}
&\left\langle -p_{0N}\left| \hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right.\right\rangle \\
&\quad= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \prod_{r<s}^{M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\delta^{(p+1)}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right) \\
&\qquad\times\prod_{r<s}^{M}\exp\left[2\alpha' G^{\mu\nu}\left(k_\mu^{(r)}-a_rA_\mu(k^{(r)})\partial_{\sigma_r}\right)\left(k_\nu^{(s)}-a_sA_\nu(k^{(s)})\partial_{\sigma_s}\right)\ln\left|e^{i\sigma_r}-e^{i\sigma_s}\right|\right] \\
&\qquad\times\exp\left[-\frac{i}{2\pi}\sum_{r=1}^{M}\left(a_rA_\mu(k^{(r)})-\sigma_r k_\mu^{(r)}\right)\theta^{\mu\nu}p_{0\nu}\right]\ .
\end{aligned}\tag{9.2}$$

The zero-slope limit can be read from the RHS as follows :

$$\begin{aligned}
&\left\langle -p_{0N}\left| \hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right.\right\rangle \\
&\quad\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \prod_{r<s}^{M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\delta^{(p+1)}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right) \\
&\qquad\times\exp\left[-\frac{i}{2\pi}\sum_{r=1}^{M}\left(a_rA_\mu(k^{(r)})-\sigma_r k_\mu^{(r)}\right)\theta^{\mu\nu}p_{0\nu}\right]\ .
\end{aligned}\tag{9.3}$$

This gives

$$\begin{aligned}
&\left\langle -p_{0N}\left|B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right.\right\rangle \\
&= \prod_{r=1}^{M}\left(i\frac{\partial}{\partial a_r}\right)\Bigg|_{a_r=0}\left\langle -p_{0N}\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right.\right\rangle \\
&\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}}\prod_{r<s}^{M} e^{\frac{i}{2}k^{(r)}_\mu\theta^{\mu\nu}k^{(s)}_\nu\epsilon(\sigma_r-\sigma_s)}\delta^{(p+1)}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right) \\
&\qquad\times\prod_{r=1}^{M}\left(\frac{dy_0^\mu(\sigma_r)}{d\sigma_r}A_\mu(k^{(r)})e^{ik^{(r)}_\nu y_0^\nu(\sigma_r)}\right) \\
&= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}}\int\frac{d^{p+1}x}{(2\pi)^{p+1}}e^{ip_{0\mu}x^\mu}e^{-\frac{i}{2}(k^{(1)}_\mu+\cdots+k^{(M)}_\mu)\theta^{\mu\nu}p_{0\nu}}\prod_{r<s}^{M}e^{\frac{i}{2}k^{(r)}_\mu\theta^{\mu\nu}k^{(s)}_\nu\epsilon(\sigma_r-\sigma_s)} \\
&\qquad\times\prod_{r=1}^{M}\left\{\frac{dy_0^\mu(\sigma_r)}{d\sigma_r}A_\mu(k^{(r)})e^{ik^{(r)}_\nu\left(x^\nu+y_0^\nu(\sigma_r)\right)}\right\}\ , \qquad (9.4)
\end{aligned}$$

where $y_0^\mu(\sigma)$ $(0\le\sigma\le 2\pi)$ is a straight line defined as

$$y_0^\mu(\sigma)=\theta^{\mu\nu}p_{0\nu}\frac{\sigma}{2\pi}\ . \qquad (9.5)$$

In the above we have used the following equations to obtain the last equality :

$$\begin{aligned}
\delta^{(p+1)}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right) &= \delta^{(p+1)}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right)\, e^{-\frac{i}{2}(k^{(1)}_\mu+\cdots+k^{(M)}_\mu)\theta^{\mu\nu}p_{0\nu}} \\
&= \int\frac{d^{p+1}x}{(2\pi)^{p+1}}e^{i(k^{(1)}_\mu+\cdots+k^{(M)}_\mu)x^\mu}e^{ip_{0\mu}x^\mu}e^{-\frac{i}{2}(k^{(1)}_\mu+\cdots+k^{(M)}_\mu)\theta^{\mu\nu}p_{0\nu}}\ . \qquad (9.6)
\end{aligned}$$

The relevant moduli parameters are σ_r with $0\le\sigma_1\le\cdots\le\sigma_M\le 2\pi$. Their integration will give us the amplitude. Eq.(9.4) yields

$$\begin{aligned}
&\int_{0\le\sigma_1\le\sigma_2\le\cdots\le\sigma_M\le 2\pi}\prod_{r=1}^{M}d\sigma_r\ \left\langle -p_{0N}\left|B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right.\right\rangle \\
&\approx\left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}}\int\frac{d^{p+1}x}{(2\pi)^{p+1}}\int_{0\le\sigma_1\le\sigma_2\le\cdots\le\sigma_M\le 2\pi}\prod_{r=1}^{M}d\sigma_r \\
&\qquad\left\{\frac{dy_0^\mu(\sigma_1)}{d\sigma_r}A_\mu(k^{(1)})e^{ik^{(1)}_\nu\left(x^\nu+y_0^\nu(\sigma_1)\right)}\right\}\star\left\{\frac{dy_0^\mu(\sigma_2)}{d\sigma_r}A_\mu(k^{(2)})e^{ik^{(2)}_\nu\left(x^\nu+y_0^\nu(\sigma_2)\right)}\right\}\star\cdots \\
&\qquad\qquad\cdots\star\left\{\frac{dy_0^\mu(\sigma_M)}{d\sigma_r}A_\mu(k^{(M)})e^{ik^{(M)}_\nu\left(x^\nu+y_0^\nu(\sigma_M)\right)}\right\}\star e^{ip_{0\mu}x^\mu}\ , \qquad (9.7)
\end{aligned}$$

where the Moyal products are taken with respect to x. The straight open Wilson line is obtained from the above equation by summing up with respect to M as follows :

$$\sum_{M=0}^{\infty} i^M \int \prod_{r=1}^{M} \frac{d^{p+1}k^{(r)}}{(2\pi)^{\frac{p+1}{2}}} \int_{0\leq\sigma_1\leq\sigma_2\leq\cdots\leq\sigma_M\leq 2\pi} \prod_{r=1}^{M} d\sigma_r \ \left\langle -p_{0N} \left| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right.\right\rangle$$

$$\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \int \frac{d^{p+1}x}{(2\pi)^{p+1}} \left[\mathcal{P}_\star \exp\left(i \int_0^{2\pi} d\sigma \frac{dy_0^\mu(\sigma)}{d\sigma} \mathcal{A}_\mu\left(x + y_0(\sigma)\right)\right)\right] \star e^{ip_{0\mu}\hat{x}^\mu} \ , \qquad (9.8)$$

where the path is taken along the straight line $x_0^\mu(\sigma) \equiv x^\mu + y_0^\mu(\sigma)$.

9.2 Curved open Wilson lines and closed-string states

Taking account of the fact that closed-string tachyons give rise to straight open Wilson lines, Dhar and Kitazawa suggested in [27] that fluctuations of open Wilson lines should originate in the massive states of closed-string. They conjectured a possible correspondence between the perturbative massive states of closed-string and the gauge theory operators obtained as the coefficients in a perturbative expansion of open Wilson line (the harmonic expansion at the straight line). In this subsection we prove their conjecture. We show, in a self-contained fashion, how one can obtain curved open Wilson lines and present an explicit correspondence between their fluctuations and the closed-string states.

Let $P_\mu(\sigma)$ be a loop in the momentum space. The harmonic expansion is given by $P_\mu(\sigma) = \frac{1}{2\pi} \left[p_{0\mu} + \frac{1}{\sqrt{2}} \sum_{n=1}^{\infty} \left(\psi_{n\mu} e^{-in\sigma} + \bar{\psi}_{n\mu} e^{in\sigma}\right)\right]$. We first introduce the following *out*-state of closed-string :

$$\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| \equiv \langle \mathbf{0} | :\exp\left(i \int_0^{2\pi} d\sigma P_\mu(\sigma) \hat{X}^\mu(\sigma)\right): \ . \qquad (9.9)$$

One may think of this state as a stringy extension, (or a generalization to include the massive modes), of the closed-string tachyon state. The oscillator representation can be read as

$$\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| = \langle -p_{0N} | \prod_{n=1}^{\infty} \exp\left[-\frac{\sqrt{\alpha'}}{2} \frac{1}{n} \left(\bar{\psi}_{n\mu} \alpha_n^\mu + \psi_{n\mu} \tilde{\alpha}_n^\mu\right)\right] \ . \qquad (9.10)$$

This tells us that we can write the state in terms of the coherent state given in Appendix B as follows :

$$\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| = \langle -p_{0N} | \otimes \prod_{n=1}^{\infty} \left(-\frac{\sqrt{\alpha'}}{2} \psi_n, -\frac{\sqrt{\alpha'}}{2} \bar{\psi}_n \right| \ , \qquad (9.11)$$

where $\left(-\frac{\sqrt{\alpha'}}{2} \psi_n, -\frac{\sqrt{\alpha'}}{2} \bar{\psi}_n \right|$ are the coherent states $(\lambda_n^+, \lambda_n^- |$ of the n-th levels defined in Eq.(B.5) with setting the complex variables $\lambda_n^+ = -\frac{\sqrt{\alpha'}}{2}\psi_n$ and $\lambda_n^- = -\frac{\sqrt{\alpha'}}{2}\bar{\psi}_n$. One can readily find that for $n \geq 1$

$$\begin{aligned} \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| \alpha^\mu_{-n} &= -\frac{\sqrt{\alpha'}}{2} g^{\mu\nu} \bar{\psi}_{n\nu} \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| \ , \\ \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| \tilde{\alpha}^\mu_{-n} &= -\frac{\sqrt{\alpha'}}{2} g^{\mu\nu} \psi_{n\nu} \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| \ . \end{aligned} \tag{9.12}$$

Recalling $\overline{\psi}_n$ is the complex conjugate of ψ_n, one can think of the state $\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right|$ as a real section of the coherent state of closed-string. Eq.(9.10) also yields the following equalities :

$$\begin{aligned} &\prod_{n=1}^{\infty} \prod_{\mu=0}^{p} \left(-\frac{2n}{\sqrt{\alpha'}} \frac{\partial}{\partial \bar{\psi}_{n\mu}} \right)^{m(n,\mu)} \left(-\frac{2n}{\sqrt{\alpha'}} \frac{\partial}{\partial \psi_{n\mu}} \right)^{m'(n,\mu)} \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| \Bigg|_{\psi, \bar{\psi}=0} \\ &\qquad = \langle -p_{0N} | \prod_{n=1}^{\infty} \prod_{\mu=0}^{p} \left(\alpha^\mu_n \right)^{m(n,\mu)} \left(\tilde{\alpha}^\mu_n \right)^{m'(n,\mu)} \ , \end{aligned} \tag{9.13}$$

where $m(n, \mu)$ and $m'(n, \mu)$ are integers greater than or equal to zero. This implies that the state $\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right|$ is a generating function of the closed-string states which are off-shell in general.

Now we wish to compute overlaps of the above state, instead of the tachyon, with the boundary states. Since it is a generating function of the closed-string states, these overlaps give us a generating function of amplitudes between closed-string states and gluons. We will show in the below that the zero-slope limit of this generating function is nothing but an open Wilson line taken along a curve parametrized by ψ_n and $\bar{\psi}_n$.

By using Eq.(9.12), we obtain

$$\begin{aligned} &\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \middle| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \\ &\quad = \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} \prod_{r<s}^{M} e^{\frac{i}{2} k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu \epsilon(\sigma_r - \sigma_s)} \delta^{(p+1)} \left(p_0 + \sum_{r=1}^{M} k^{(r)} \right) \\ &\qquad \times \prod_{r<s}^{M} \exp \left[2\alpha' G^{\mu\nu} \left(k^{(r)}_\mu - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k^{(s)}_\nu - a_s A_\nu(k^{(s)}) \partial_{\sigma_s} \right) \ln \left| e^{i\sigma_r} - e^{i\sigma_s} \right| \right] \\ &\qquad \times \prod_{n=1}^{\infty} \exp \left[-\frac{\alpha'}{4n} \bar{\psi}_{n\mu} \left(g^{-1} N g^{-1} \right)^{\mu\nu} \psi_{n\nu} \right] \\ &\qquad \times \exp \left[-i \sum_{r=1}^{M} a_r A_\mu(k^{(r)}) \right. \\ &\qquad\qquad\qquad \left. \times \left\{ \frac{\theta^{\mu\nu}}{2\pi} p_{0\nu} + \frac{\alpha'}{\sqrt{2}} \sum_{n=1}^{\infty} \left(\left(\frac{1}{E} \right)^{\mu\nu} \psi_{n\nu} e^{-in\sigma_r} - \left(\frac{1}{E^T} \right)^{\mu\nu} \bar{\psi}_{n\nu} e^{in\sigma_r} \right) \right\} \right] \end{aligned}$$

$$\times \exp\left[i \sum_{r=1}^{M} k_{\mu}^{(r)} \left\{ \frac{\theta^{\mu\nu}}{2\pi} p_{0\nu}\sigma_r + \frac{\alpha'}{\sqrt{2}} \sum_{n=1}^{\infty} \frac{i}{n} \left(\left(\frac{1}{E}\right)^{\mu\nu} \psi_{n\nu} e^{-in\sigma_r} + \left(\frac{1}{E^T}\right)^{\mu\nu} \bar{\psi}_{n\nu} e^{in\sigma_r} \right) \right\} \right] . \tag{9.14}$$

The zero-slope limit can be read from the RHS. In the present limiting procedure, we have

$$\alpha' \left(\frac{1}{E}\right)^{\mu\nu} \approx \frac{\theta^{\mu\nu}}{2\pi} \ , \quad \alpha' \left(\frac{1}{E^T}\right)^{\mu\nu} \approx -\frac{\theta^{\mu\nu}}{2\pi} \ , \quad \alpha' \left(g^{-1} N g^{-1}\right)^{\mu\nu} \approx \frac{1}{4\pi^2 \alpha'} \left(\theta G \theta\right)^{\mu\nu} \ . \tag{9.15}$$

This enables us to find out that

$$\begin{aligned} &\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \left| \hat{B}_N[A]; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \right. \\ &\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} \prod_{r<s}^{M} e^{\frac{i}{2} k_{\mu}^{(r)} \theta^{\mu\nu} k_{\nu}^{(s)} \epsilon(\sigma_r - \sigma_s)} \delta^{(p+1)} \left(p_0 + \sum_{r=1}^{M} k^{(r)} \right) \\ &\quad \times \prod_{n=1}^{\infty} \exp\left[-\frac{\bar{\psi}_{n\mu} \left(\theta G \theta\right)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n \alpha'} \right] \exp\left[-i \sum_{r=1}^{M} \left\{ a_r \frac{dy^{\mu}(\sigma_r)}{d\sigma_r} A_{\mu}(k^{(r)}) - k_{\mu}^{(r)} y^{\mu}(\sigma_r) \right\} \right] . \end{aligned} \tag{9.16}$$

Here we introduce $y^{\mu}(\sigma)$ with

$$y^{\mu}(\sigma) = y_0^{\mu}(\sigma) + \tilde{y}^{\mu}(\sigma) \ , \tag{9.17}$$

where

$$\tilde{y}^{\mu}(\sigma) \equiv \theta^{\mu\nu} \frac{1}{2\pi} \frac{1}{\sqrt{2}} \sum_{n=1}^{\infty} \frac{i}{n} \left(\psi_{n\nu} e^{-in\sigma} - \bar{\psi}_{n\nu} e^{in\sigma} \right) \ . \tag{9.18}$$

It is a curve deviating from the straight line $y_0^{\mu}(\sigma)$ and the deviation is denoted by $\tilde{y}^{\mu}(\sigma)$.

It follows from Eq.(9.16) that

$$\begin{aligned} &\left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \left| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \right. \\ &= \prod_{r=1}^{M} \left(i \frac{\partial}{\partial a_r} \right) \Bigg|_{a_r = 0} \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \left| \hat{B}_N[A]; \{a_r\} (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \right. \\ &\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} \prod_{r<s}^{M} e^{\frac{i}{2} k_{\mu}^{(r)} \theta^{\mu\nu} k_{\nu}^{(s)} \epsilon(\sigma_r - \sigma_s)} \delta^{(p+1)} \left(p_0 + \sum_{r=1}^{M} k^{(r)} \right) \\ &\quad \times \prod_{n=1}^{\infty} \exp\left[-\frac{\bar{\psi}_{n\mu} \left(\theta G \theta\right)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n \alpha'} \right] \prod_{r=1}^{M} \left\{ \frac{dy^{\mu}(\sigma_r)}{d\sigma_r} A_{\mu}(k^{(r)}) e^{i k_{\nu}^{(r)} y^{\nu}(\sigma_r)} \right\} . \end{aligned} \tag{9.19}$$

This takes essentially the same form as Eq.(9.4) with the straight line $y_0^\mu(\sigma)$ replaced by the curved one $y^\mu(\sigma)$. Therefore, performing the same rearrangement as carried out in the last subsection, we obtain

$$\begin{aligned}
&\sum_{M=0}^{\infty} i^M \int \prod_{r=1}^{M} \frac{d^{p+1}k^{(r)}}{(2\pi)^{\frac{p+1}{2}}} \int_{0\le\sigma_1\le\cdots\le\sigma_M\le 2\pi} \prod_{r=1}^{M} d\sigma_r \\
&\qquad\qquad \times \left\langle \Omega(\psi_n,\bar{\psi}_n;p_0) \,\middle|\, B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)}) \right\rangle \\
&\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \prod_{n=1}^{\infty} \exp\left[-\frac{\bar{\psi}_{n\mu}\left(\theta G\theta\right)^{\mu\nu}\psi_{n\nu}}{16\pi^2 n\alpha'}\right] \\
&\qquad \times \int \frac{d^{p+1}x}{(2\pi)^{p+1}} \left[\mathcal{P}_\star \exp\left(i\int_0^{2\pi} d\sigma \frac{dy^\mu(\sigma)}{d\sigma}\mathcal{A}_\mu(x+y(\sigma))\right)\right] \star e^{ip_{0\mu}x^\mu} \ . \qquad (9.20)
\end{aligned}$$

This is the open Wilson line taken along the curve $x^\mu(\sigma) \equiv x^\mu + y^\mu(\sigma)$.

Thus we have shown that the zero-slope limit of the generating function of the amplitudes between closed-string states and gluons becomes the open Wilson line multiplied by a Gaussian weight. The path is curved by ψ_n and $\overline{\psi}_n$. As can be seen in Eq.(9.13), these variables originally measure condensations of the n-th massive modes of closed-string.

This completes the proof of the conjecture.

Overlap with graviton

As has been mentioned, the overlap $\left\langle \Omega(\psi_n,\bar{\psi}_n;p_0) \,\middle|\, B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)}) \right\rangle$ serves as a generating function of the amplitudes between closed-string states and gluons. As an illustration, let us consider the case of gravitons.

The graviton states can be written as follows :

$$\langle -p_{0N}|\, h_{\mu\nu}(p_0)\alpha_1^\mu\tilde{\alpha}_1^\nu = \frac{4}{\alpha'} h_{\mu\nu}(p_0) \frac{\partial^2}{\partial\bar{\psi}_{1\mu}\partial\psi_{1\nu}} \left\langle \Omega\left(\psi_n,\bar{\psi}_n;p_0\right)\right| \Big|_{\psi,\bar{\psi}=0} \ , \qquad (9.21)$$

where $h_{\mu\nu}(p_0)$ denotes the polarization tensor of graviton which is symmetric and traceless. Using Eq.(9.19), we obtain

$$\begin{aligned}
&\left\langle -p_{0N}\right|\alpha_1^\mu\tilde{\alpha}_1^\nu \left| B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)}) \right\rangle \\
&\quad = \frac{4}{\alpha'} \frac{\partial^2}{\partial\bar{\psi}_{1\mu}\partial\psi_{1\nu}} \left\langle \Omega\left(\psi_n,\bar{\psi}_n;p_0\right) \,\middle|\, B_N;(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)}) \right\rangle \Big|_{\psi,\bar{\psi}=0} \\
&\quad \approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \prod_{r=1}^{M} e^{ik_\mu^{(r)}\theta^{\mu\nu}p_{0\nu}\frac{\sigma_r}{2\pi}} \prod_{r<s}^{M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)} \delta^{(p+1)}\left(p_{0\mu}+\sum_{r=1}^{M}k_\mu^{(r)}\right)
\end{aligned}$$

$$
\begin{aligned}
&\times\frac{2}{\alpha'}\left(-\frac{1}{2\pi}\right)^{M}\Bigg[-\frac{(\theta G\theta)^{\mu\nu}}{8\pi^2\alpha'}\prod_{u=1}^{M}\left(p_0\theta A(k^{(u)})\right) \\
&\quad+\sum_{r=1}^{M}\Bigg\{\left(\theta A(k^{(r)})\right)^{\mu}\frac{\left(\theta k^{(r)}\right)^{\nu}}{2\pi}-\frac{\left(\theta k^{(r)}\right)^{\mu}}{2\pi}\left(\theta A(k^{(r)})\right)^{\nu} \\
&\qquad\qquad+i\frac{\left(\theta k^{(r)}\right)^{\mu}}{2\pi}i\frac{\left(\theta k^{(r)}\right)^{\nu}}{2\pi}\left(p_0\theta A(k^{(r)})\right)\Bigg\}\prod_{u\neq r}\left(p_0\theta A(k^{(u)})\right) \\
&\quad+\sum_{r\neq s}\Bigg\{\left(i\frac{\left(\theta k^{(r)}\right)^{\mu}\left(p_0\theta A(k^{(r)})\right)}{2\pi}-i\left(\theta A(k^{(r)})\right)^{\mu}\right)e^{i\sigma_r} \\
&\qquad\times\left(i\frac{\left(\theta k^{(s)}\right)^{\nu}\left(p_0\theta A(k^{(s)})\right)}{2\pi}+i\left(\theta A(k^{(s)})\right)^{\nu}\right)e^{-i\sigma_s}\Bigg\}\prod_{u\neq r,s}\left(p_0\theta A(k^{(u)})\right)\Bigg]\ .
\end{aligned}
\tag{9.22}
$$

This reproduces the result obtained in Appendix A of [23]. In fact, combining the above equation with Eq.(9.21), we obtain

$$
\begin{aligned}
&\langle -p_{0N}|h_{\mu\nu}(p_0)\alpha_1^{\mu}\tilde{\alpha}_1^{\nu}\left|B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle \\
&\quad\approx\left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}}\prod_{r=1}^{M}e^{ik_\mu^{(r)}\theta^{\mu\nu}p_{0\nu}\frac{\sigma_r}{2\pi}}\prod_{r<s}^{M}e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\prod_{\mu=0}^{p}\delta\left(p_{0\mu}+\sum_{r=1}^{M}k_\mu^{(r)}\right) \\
&\qquad\times\frac{2}{\alpha'}\left(-\frac{1}{2\pi}\right)^{M}\Bigg[\sum_{r=1}^{M}\frac{1}{(2\pi)^2}k_\mu^{(r)}\left(\theta h(p_0)\theta\right)^{\mu\nu}k_\nu^{(r)}\prod_{u=1}^{M}\left(p_0\theta A(k^{(u)})\right) \\
&\qquad\quad+\sum_{r\neq s}\Bigg\{\left(\frac{\left(p_0\theta A(k^{(r)})\right)}{2\pi}k_\mu^{(r)}-A_\mu(k^{(r)})\right)\left(\theta h(p_0)\theta\right)^{\mu\nu}\left(\frac{\left(p_0\theta A(k^{(s)})\right)}{2\pi}k_\nu^{(s)}+A_\nu(k^{(s)})\right) \\
&\qquad\qquad\qquad\times e^{i(\sigma_r-\sigma_s)}\prod_{u\neq r,s}\left(p_0\theta A(k^{(u)})\right)\Bigg\}\Bigg]\ ,
\end{aligned}
\tag{9.23}
$$

where the properties of $h_{\mu\nu}(p_0)$ are used in the following manner :

$$
0=\alpha' g^{\mu\nu}h_{\mu\nu}(p_0)=\left(\alpha' G^{\mu\nu}-\frac{(\theta G\theta)^{\mu\nu}}{(2\pi)^2\alpha'}\right)h_{\mu\nu}(p_0)\approx-\frac{(\theta G\theta)^{\mu\nu}}{(2\pi)^2\alpha'}h_{\mu\nu}(p_0)\ . \tag{9.24}
$$

It is worth emphasizing that Eq.(9.23) is derived without using the on-shell conditions of graviton, i.e. neither $g^{\mu\nu}p_{0\mu}h_{\nu\lambda}(p_0)=0$ or $\alpha' g^{\mu\nu}p_{0\mu}p_{0\nu}=0$.

9.3 Reparametrization invariance of open Wilson lines

Wilson lines in gauge theories are invariant under reparametrizations of the paths. Transformations analogous to the reparametrizations are generated by $L_n-\tilde{L}_{-n}$ in closed-string theory.

In fact, $L_n - \tilde{L}_{-n}$ can be identified with vector fields $z^{n+1}\partial_z - \bar{z}^{-n+1}\partial_{\bar{z}}$ on the world-sheet [9] and at $\tau = 0$, where the boundary states reside, these vector fields have the forms of $e^{in\sigma}\partial_\sigma$. We have observed that the action of diffS^1 on the boundary states is identified with the action of the closed-string BRST charge. The boundary states are in general not BRST-closed. Hence the reparametrization invariance of open Wilson lines indicates that the action of diffS^1 or the BRST charge Q_c becomes null at the zero-slope limit.

Let us verify the above observation. By making use of Eqs.(2.11) and (7.19), we obtain for $\forall n \in \mathbb{Z}$

$$\begin{aligned} &\left(L_n - \tilde{L}_{-n}\right) \left|B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \\ &\quad = \sum_{r=1}^{M} e^{in\sigma_r} \left\{ -i\frac{\partial}{\partial \sigma_r} + n\left(\alpha' k^{(r)}_\mu G^{\mu\nu} k^{(r)}_\nu + 1\right) \right\} \left|B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \\ &\qquad + \sum_{r=1}^{M} e^{in\sigma_r} \alpha' k^{(r)}_\mu G^{\mu\nu} A_\nu(k^{(r)}) \left|B_N[A]_{\check{r}}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \ , \end{aligned} \tag{9.25}$$

where $\left|B_N[A]_{\check{r}}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle$ denotes

$$\lim_{\forall \tau_r \to 0+} V^{ren}_A(\sigma_1, \tau_1; k^{(1)}) \cdots V^{ren}_T(\sigma_r, \tau_r; k^{(r)}) \cdots V^{ren}_A(\sigma_M, \tau_M; k^{(M)}) \left|B_N\right\rangle \ . \tag{9.26}$$

Combining Eq.(9.25) with Eq.(9.20), we find that in the zero-slope limit

$$\begin{aligned} &\int \prod_{r=1}^{M} \frac{d^{p+1}k^{(r)}}{(2\pi)^{\frac{p+1}{2}}} \int_{0 \le \sigma_1 \le \cdots \le \sigma_M \le 2\pi} \prod_{r=1}^{M} d\sigma_r \\ &\qquad \times \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0)\right| i\epsilon_n \left(L_n - \tilde{L}_{-n}\right) \left|B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \\ &\approx \int \prod_{r=1}^{M} \frac{d^{p+1}k^{(r)}}{(2\pi)^{\frac{p+1}{2}}} \int_{0 \le \sigma_1 \le \cdots \le \sigma_M \le 2\pi} \prod_{s=1}^{M} d\sigma_s \\ &\qquad \times \sum_{r=1}^{M} \frac{\partial}{\partial \sigma_r} \epsilon_n e^{in\sigma_r} \left\langle \Omega(\psi_n, \bar{\psi}_n; p_0) \right| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\rangle \\ &\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}} \right)^{\frac{1}{4}} \prod_{n=1}^{\infty} \exp\left[-\frac{\bar{\psi}_{n\mu} \left(\theta G \theta\right)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n\alpha'} \right] \int \frac{d^{p+1}x}{(2\pi)^{p+1}} \int_{0 \le \sigma_1 \le \cdots \le \sigma_M \le 2\pi} \prod_{s=1}^{M} d\sigma_s \\ &\quad \sum_{r=1}^{M} \frac{\partial}{\partial \sigma_r} \epsilon_n e^{in\sigma_r} \left\{ \frac{dy^\mu(\sigma_1)}{d\sigma_1} \mathcal{A}_\mu(x + y(\sigma_1)) \right\} \star \cdots \star \left\{ \frac{dy^\mu(\sigma_M)}{d\sigma_M} \mathcal{A}_\mu(x + y(\sigma_M)) \right\} \star e^{ip_{0\mu}x^\mu} \ , \end{aligned} \tag{9.27}$$

---

[9] $z = e^{\tau + i\sigma}$.

where ϵ_n denotes an infinitesimal parameter. This shows that at the zero-slope limit $L_n - \tilde{L}_{-n}$ give rise to infinitesimal reparametrizations $\sigma \mapsto \sigma' = \sigma + \delta_\epsilon^{(n)}\sigma$, where $\delta_\epsilon^{(n)}\sigma \equiv -\epsilon_n e^{in\sigma}$. This can be checked by recalling that the pull-back $\tilde{\mathcal{A}}(\sigma) = \frac{dy^\mu(\sigma)}{d\sigma}\mathcal{A}_\mu(x+y(\sigma))$ on the path transforms under the infinitesimal reparametrizations as $\tilde{\mathcal{A}}(\sigma) \mapsto \tilde{\mathcal{A}}(\sigma) + \delta_\epsilon^{(n)}\tilde{\mathcal{A}}(\sigma)$, where

$$\delta_\epsilon^{(n)}\tilde{\mathcal{A}}(\sigma) = \frac{\partial}{\partial\sigma}\left(\epsilon_n e^{in\sigma}\frac{dy^\mu(\sigma)}{d\sigma}\mathcal{A}_\mu(x+y(\sigma))\right) \; . \tag{9.28}$$

From this, we can write the integrand in the RHS of Eq.(9.27) as follows :

$$\begin{aligned}&\sum_{r=1}^{M}\frac{\partial}{\partial\sigma_r}\epsilon_n e^{in\sigma_r}\left\{\frac{dy^\mu(\sigma_1)}{d\sigma_1}\mathcal{A}_\mu(x+y(\sigma_1))\right\}\star\cdots\star\left\{\frac{dy^\mu(\sigma_M)}{d\sigma_M}\mathcal{A}_\mu(x+y(\sigma_M))\right\}\\ &\quad = \delta_\epsilon^{(n)}\left(\left\{\frac{dy^\mu(\sigma_1)}{d\sigma_1}\mathcal{A}_\mu(x+y(\sigma_1))\right\}\star\cdots\star\left\{\frac{dy^\mu(\sigma_M)}{d\sigma_M}\mathcal{A}_\mu(x+y(\sigma_M))\right\}\right) \; .\end{aligned} \tag{9.29}$$

The above integrand is a total derivative with respect to each σ_r. The σ_r integrations in (9.27) give rise to surface terms. Let us show that these surface terms actually cancel out. Taking account of the integration region of σ_r being $[\sigma_{r-1}, \sigma_{r+1}]$ for $2 \le r \le M-1$, the following equality holds for arbitrary $\mathcal{F}$:

$$\begin{aligned}&\int_{0\le\sigma_1\le\cdots\le\sigma_M\le 2\pi}\prod_{s=1}^{M}d\sigma_s\left(\frac{\partial}{\partial\sigma_r}e^{in\sigma_r}+\frac{\partial}{\partial\sigma_{r+1}}e^{in\sigma_{r+1}}\right)\mathcal{F}(\sigma_1,\cdots,\sigma_M)\\ &= \int_{0\le\tilde\sigma_1\le\cdots\le\tilde\sigma_{M-1}}\prod_{s=1}^{M-1}d\tilde\sigma_s\Big\{e^{in\tilde\sigma_{r+1}}\mathcal{F}\left(\tilde\sigma_1,\cdots,\tilde\sigma_r,\tilde\sigma_{r+1}\tilde\sigma_{r+1},\tilde\sigma_{r+2},\cdots,\tilde\sigma_{M-1}\right)\\ &\qquad\qquad -e^{in\tilde\sigma_{r-1}}\mathcal{F}\left(\tilde\sigma_1,\cdots,\tilde\sigma_{r-2},\tilde\sigma_{r-1},\tilde\sigma_{r-1},\tilde\sigma_r,\cdots,\tilde\sigma_{M-1}\right)\Big\} \; .\end{aligned} \tag{9.30}$$

This leads us to find

$$\begin{aligned}&\int\frac{d^{p+1}x}{(2\pi)^{p+1}}\int_{0\le\sigma_1\le\cdots\le\sigma_M\le 2\pi}\prod_{s=1}^{M}d\sigma_s\\ &\sum_{r=1}^{M}\frac{\partial}{\partial\sigma_r}\epsilon_n e^{in\sigma_r}\left\{\frac{dy^\mu(\sigma_1)}{d\sigma_1}\mathcal{A}_\mu(x+y(\sigma_1))\right\}\star\cdots\star\left\{\frac{dy^\mu(\sigma_M)}{d\sigma_M}\mathcal{A}_\mu(x+y(\sigma_M))\right\}\star e^{ip_{0\mu}x^\mu}\\ &= \int\frac{d^{p+1}x}{(2\pi)^{p+1}}\int_{0\le\sigma_1\le\cdots\le\sigma_{M-1}\le 2\pi}\prod_{s=1}^{M-1}d\sigma_s\\ &\quad\left\{\frac{dy^\mu(\sigma_1)}{d\sigma_1}\mathcal{A}_\mu(x+y(\sigma_1))\right\}\star\cdots\star\left\{\frac{dy^\mu(\sigma_{M-1})}{d\sigma_{M-1}}\mathcal{A}_\mu(x+y(\sigma_{M-1}))\right\}\\ &\quad\star\epsilon_n\left[\left\{\frac{dy^\mu(2\pi)}{d\sigma}\mathcal{A}_\mu(x+y(2\pi))\right\}\star e^{ip_{0\mu}x^\mu}-e^{ip_{0\mu}x^\mu}\star\left\{\frac{dy^\mu(0)}{d\sigma}\mathcal{A}_\mu(x+y(0))\right\}\right] \; .\end{aligned} \tag{9.31}$$

Here we have used the cyclic property of the Moyal product inside the integration: $\int d^{p+1}x\, f(x)\star g(x) = \int d^{p+1}x\, g(x)\star f(x)$. Let us recall that $e^{ip_{0\mu}x^\mu}$ plays a role of a translation generator on the non-commutative space-time:

$$f(x)\star e^{ip_{0\mu}x^\mu} = e^{ip_{0\mu}x^\mu}\star f(x-\theta p_0)\ . \tag{9.32}$$

Combined with the relations $\frac{dy^\mu}{d\sigma}(2\pi) = \frac{dy^\mu}{d\sigma}(0)$ and $y^\mu(2\pi) - y^\mu(0) = \theta^{\mu\nu}p_{0\nu}$, this tells us that Eq.(9.31) is vanishing and so is Eq.(9.27).

Therefore we have shown that the action of diffS^1 on the boundary states reduces to infinitesimal reparametrizations of paths of open Wilson lines and becomes null at the zero-slope limit. The situation is summarized as follows :

$$\begin{aligned}
&\sum_{M=0}^{\infty} i^M \int \prod_{r=1}^{M} \frac{d^{p+1}k^{(r)}}{(2\pi)^{\frac{p+1}{2}}} \int_{0\le\sigma_1\le\cdots\le\sigma_M\le 2\pi} \prod_{r=1}^{M} d\sigma_r \\
&\qquad\qquad \times \left\langle \Omega(\psi_n,\bar{\psi}_n;p_0)\right|\, i\epsilon_n \left(L_n - \tilde{L}_{-n}\right) \left|B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle \\
&\quad \approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{p+1}}\right)^{\frac{1}{4}} \prod_{n=1}^{\infty} \exp\left[-\frac{\bar{\psi}_{n\mu}\,(\theta G\theta)^{\mu\nu}\,\psi_{n\nu}}{16\pi^2 n\alpha'}\right] \\
&\qquad \delta_\epsilon^{(n)} \int \frac{d^{p+1}x}{(2\pi)^{p+1}} \mathcal{P}_\star \left[\exp\left(i\int_0^{2\pi} d\sigma \frac{y^\mu(\sigma)}{d\sigma}\mathcal{A}_\mu(x+y(\sigma))\right)\right] \star e^{ip_{0\mu}x^\mu} \\
&\quad = 0\ .
\end{aligned} \tag{9.33}$$

9.4 Factorization by closed-string momentum eigenstates

We have seen in Section 6 that analogues of open Wilson line, $\mathcal{P}_\star\left[\exp\left\{\int_0^{2\pi} d\sigma\phi(x(\sigma))\right\}\right]$, factorize the generating function of one-loop amplitudes of open-string tachyons at the zero-slope limit. Decompositions of the string amplitudes made by insertions of the unity, $1 = \int [dP_N] |P_N\rangle\langle P_N|$, play an important role to gain the factorization (6.12). With the same manipulation one can expect that a similar factorization is also obtainable for the amplitudes of gluons. Prior to the actual computations, let us explain briefly why the decompositions via the momentum eigenstates give us open Wilson lines or their analogues.

To make the discussion transparent we start with the momentum eigenstate $\langle -P_N|$. As described in Eq.(5.42), we can write the state as

$$\langle -P_N| = \prod_{n=1}^{\infty} \exp\left[-\frac{\alpha'}{4n}\bar{\psi}_{n\mu}g^{\mu\nu}\psi_{n\nu}\right] \times {}_{B=0}\langle B_N| : \exp\left(i\int_0^{2\pi} d\sigma P_\mu(\sigma)\hat{X}^\mu(\sigma)\right) :\ . \tag{9.34}$$

Taking account of its use in the decompositions of the amplitudes we should consider the state $\langle -P_N| q_c^{\frac{1}{4}(L_0+\tilde{L}_0-2)}$ rather than $\langle -P_N|$. Let us also recall that the above ψ_n and $\overline{\psi}_n$ are rescaled appropriately in order to obtain the factorization (6.12) and that the rescaled variables are kept intact under taking the limit. To be explicit, we denote the rescaled ψ_n and $\bar{\psi}_n$ by the following ψ'_n and $\overline{\psi}'_n$:

$$(\psi_{n\mu}, \bar{\psi}_{n\mu}) \mapsto (\psi'_{n\mu}, \bar{\psi}'_{n\mu}) = (2q_c^{\frac{n}{4}}\psi_{n\mu}, 2q_c^{\frac{n}{4}}\bar{\psi}_{n\mu}) \ . \tag{9.35}$$

Oscillator representation of the state $\langle -P_N| q_c^{\frac{1}{4}(L_0+\tilde{L}_0-2)}$ can be read as follows by using these rescaled variables :

$$\begin{aligned}\langle -P_N|\, q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)} &= q_c^{\frac{\alpha'}{8}g^{\mu\nu}p_{0\mu}p_{0\nu}-\frac{1}{2}} \times \\ \times \langle -p_{0N}| \prod_{n=1}^{\infty} \exp&\left[-\frac{q_c^{\frac{n}{2}}}{n}\alpha_n^{\mu}g_{\mu\nu}\tilde{\alpha}_n^{\nu} - \frac{\sqrt{\alpha'}}{2n}\left(\bar{\psi}'_{n\mu}\alpha_n^{\mu} + \psi'_{n\mu}\tilde{\alpha}_n^{\mu}\right) - \frac{\alpha'}{8nq_c^{\frac{n}{2}}}\bar{\psi}'_{n\mu}g^{\mu\nu}\psi'_{n\nu}\right]\end{aligned} \tag{9.36}$$

Now it is clear that this state reduces to the state $\left\langle \Omega(\psi'_n, \bar{\psi}'_n; p_0)\right|$ at the zero-slop limit :

$$\begin{aligned}&\langle -P_N|\, q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)} \\ &\approx e^{-\frac{\pi s(c)}{4}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)}\langle -p_{0N}| \prod_{n=1}^{\infty} \exp\left[-\frac{\sqrt{\alpha'}}{2n}\left(\bar{\psi}'_{n\mu}\alpha_n^{\mu} + \psi'_{n\mu}\tilde{\alpha}_n^{\mu}\right) + \frac{\psi'_{n\mu}\left(\theta G\theta\right)^{\mu\nu}\bar{\psi}'_{n\nu}}{32\pi^2 nq_c^{\frac{n}{2}}\alpha'}\right] \\ &= e^{-\frac{\pi s(c)}{4}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)} \prod_{n=1}^{\infty} \exp\left\{\frac{\psi'_{n\mu}\left(\theta G\theta\right)^{\mu\nu}\bar{\psi}'_{n\nu}}{32\pi^2 nq_c^{\frac{n}{2}}\alpha'}\right\}\left\langle \Omega(\psi'_n, \bar{\psi}'_n; p_0)\right| \ .\end{aligned} \tag{9.37}$$

As we have seen in subsection 9.2 the above state is a generating function of the closed-string states and its overlaps with the boundary states lead the open Wilson line taken along the corresponding path.

We will compute the factorization of the generating function of one-loop amplitudes of gluons. We will restrict ourselves to 25-brane in the critical dimensions. Computations become parallel to those given in Section 6 but much complicated. The factorization at the zero-slope limit by open Wilson lines are given in Eq.(9.44).

Let us study factorizations of the following amplitudes :

$$\begin{aligned}&\left\langle B_N[A]; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right|\, q_c^{\frac{1}{2}(L_0+\tilde{L}_0-2)} \\ &\qquad\qquad \times \left|B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right\rangle \ .\end{aligned} \tag{9.38}$$

Oscillator representations of the boundary states of gluons are unsuitable for computations of the amplitudes. We instead consider the corresponding amplitudes in auxiliary forms. Let us decompose the string amplitudes (7.36) by using the momentum eigenstates as follows :

$$\left\langle \hat{B}_N[A]; \{a_r\}; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)})\right|$$

$$\times q_c^{\frac{1}{2}\left(L_0+\tilde{L}_0-2\right)}\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle$$

$$= \int [dP_N]\left\langle \hat{B}_N[A];\{a_r\};(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right| q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|-P_N\right\rangle$$

$$\times\left\langle -P_N\right| q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle \ . \qquad (9.39)$$

Here we have changed the integration variables from $P_\mu(\sigma)$ to $-P_\mu(\sigma)$ for the later convenience. The amplitudes (9.38) are obtained from the above by the operation $\prod_{r=1}^{M} i\frac{\partial}{\partial a_r}\Big|_{a_r=0}$.

Each factor in the above decomposition can be evaluated by using the oscillator representations. These are given in Eq.(7.23) (and Eq.(7.33)) for the boundary states, and Eq.(5.21) (and its hermitian conjugate) for the momentum eigenstates. We then need to compute matrix elements similar to Eq.(6.2). Eq.(B.7) enables us to calculate them. After the rescaling (9.35) of ψ_n and $\overline{\psi}_n$ we obtain the following expressions :

$$\int [dP_N]\left\langle \hat{B}_N[A];\{a_r\};(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right| q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|-P_N\right\rangle$$

$$\times\left\langle -P_N\right| q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle$$

$$= \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{2}}\int d^{26}p_0\ \delta^{26}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right)\delta^{26}\left(p_0-\sum_{r=M+1}^{M+N}k^{(r)}\right)\ q_c^{\frac{\alpha'}{4}g^{\mu\nu}p_{0\mu}p_{0\nu}-1}$$

$$\times\prod_{1\le r<s\le M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\prod_{M+1\le r<s\le M+M} e^{-\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}$$

$$\times\prod_{1\le r<s\le M}\exp\left[\alpha' G^{\mu\nu}\left(k_\mu^{(r)}-a_rA_\mu(k^{(r)})\partial_{\sigma_r}\right)\left(k_\nu^{(s)}-a_sA_\nu(k^{(s)})\partial_{\sigma_s}\right)\ln\left|e^{i\sigma_r}-e^{i\sigma_s}\right|^2\right]$$

$$\times\prod_{M+1\le r<s\le M+N}\exp\left[\alpha' G^{\mu\nu}\left(k_\mu^{(r)}-a_rA_\mu(k^{(r)})\partial_{\sigma_r}\right)\left(k_\nu^{(s)}-a_sA_\nu(k^{(s)})\partial_{\sigma_s}\right)\ln\left|e^{i\sigma_r}-e^{i\sigma_s}\right|^2\right]$$

$$\times\prod_{n=1}^{\infty}\exp\left[-\frac{2\alpha' q_c^{\frac{n}{2}}}{n}\left(\frac{1}{E^T-q_c^{\frac{n}{2}}E}g\frac{1}{E^T}\right)^{\mu\nu}\right.$$

$$\times\left\{\sum_{r,s=1}^{M}\left(k_\mu^{(r)}-ina_rA_\mu(k^{(r)})\right)\left(k_\nu+ina_sA_\nu(k^{(s)})\right)e^{in(\sigma_r-\sigma_s)}\right.$$

$$\left.\left.+\sum_{r,s=M+1}^{M+N}\left(k_\mu^{(r)}+ina_rA_\mu(k^{(r)})\right)\left(k_\nu-ina_sA_\nu(k^{(s)})\right)e^{-in(\sigma_r-\sigma_s)}\right\}\right]$$

$$\times\int\prod_{n=1}^{\infty}\left[\frac{d^{26}\bar{\psi}_n d^{26}\psi_n}{(2i)^{26}}\left(\frac{\alpha'}{4n\pi q_c^{\frac{n}{2}}}\right)^{26}\frac{-\det g_{\mu\nu}}{\det^2\left(g-q_c^{\frac{n}{2}}N\right)_{\mu\nu}}\right]$$

$$
\begin{aligned}
&\times \exp\left[-\sum_{n=1}^{\infty}\frac{\alpha'}{4n}\bar{\psi}_{n\mu}\left(\frac{1}{q_c^{\frac{n}{2}}g}+\frac{1}{g-q_c^{\frac{n}{2}}N^T}N^T\frac{1}{g}+\frac{1}{g}N\frac{1}{g-q_c^{\frac{n}{2}}N}\right)^{\mu\nu}\psi_{n\nu}\right]\\
&\times \exp\Bigg[-i\sum_{r=1}^{M}a_rA_\mu(k^{(r)})\Bigg[\frac{\theta^{\mu\nu}}{2\pi}p_{0\nu}+\frac{\alpha'}{\sqrt{2}}\sum_{n=1}^{\infty}\Bigg\{\left(\frac{1}{E-q_c^{\frac{n}{4}}E^T}\right)^{\mu\nu}\psi_{n\nu}e^{-in\sigma_r}\\
&\qquad\qquad -\left(\frac{1}{E^T-q_c^{\frac{n}{4}}E}\right)^{\mu\nu}\bar{\psi}_{n\nu}e^{in\sigma_r}\Bigg\}\Bigg]\Bigg]\\
&\times \exp\Bigg[i\sum_{r=1}^{M}k_\mu^{(r)}\Bigg[\frac{\theta^{\mu\nu}}{2\pi}p_{0\nu}\sigma_r+\frac{\alpha'}{\sqrt{2}}\sum_{n=1}^{\infty}\frac{i}{n}\Bigg\{\left(\frac{1}{E-q_c^{\frac{n}{4}}E^T}\right)^{\mu\nu}\psi_{n\nu}e^{-in\sigma_r}\\
&\qquad\qquad +\left(\frac{1}{E^T-q_c^{\frac{n}{4}}E}\right)^{\mu\nu}\bar{\psi}_{n\nu}e^{in\sigma_r}\Bigg\}\Bigg]\Bigg]\\
&\times \exp\Bigg[-i\sum_{r=M+1}^{M+N}a_rA_\mu(k^{(r)})\Bigg[\frac{\theta^{\mu\nu}}{2\pi}p_{0\nu}-\frac{\alpha'}{\sqrt{2}}\sum_{n=1}^{\infty}\Bigg\{\left(\frac{1}{E^T-q_c^{\frac{n}{4}}E}\right)^{\mu\nu}\psi_{n\nu}e^{-in\sigma_r}\\
&\qquad\qquad -\left(\frac{1}{E-q_c^{\frac{n}{4}}E^T}\right)^{\mu\nu}\bar{\psi}_{n\nu}e^{in\sigma_r}\Bigg\}\Bigg]\Bigg]\\
&\times \exp\Bigg[i\sum_{r=M+1}^{M+N}k_\mu^{(r)}\Bigg[\frac{\theta^{\mu\nu}}{2\pi}p_{0\nu}\sigma_r-\frac{\alpha'}{\sqrt{2}}\sum_{n=1}^{\infty}\frac{i}{n}\Bigg\{\left(\frac{1}{E^T-q_c^{\frac{n}{4}}E}\right)^{\mu\nu}\psi_{n\nu}e^{-in\sigma_r}\\
&\qquad\qquad +\left(\frac{1}{E-q_c^{\frac{n}{4}}E^T}\right)^{\mu\nu}\bar{\psi}_{n\nu}e^{in\sigma_r}\Bigg\}\Bigg]\Bigg]\ , \qquad (9.40)
\end{aligned}
$$

where we have newly written $(\psi'_{n\mu},\bar{\psi}'_{n\mu})$ as $(\psi_{n\mu},\bar{\psi}_{n\mu})$.

We examine the zero-slope limit of the above expression. The limiting procedure we consider is the same that was investigated in Section 8 to capture the UV behavior of the noncommutative gauge theory. It is taken by fixing parameters $s^{(c)}=\alpha'|\tau^{(c)}|$, σ_r and a_r besides the open-string tensors. ψ_n and $\overline{\psi}_n$ in Eq.(9.40) are also left intact. This yields

$$
\begin{aligned}
&\int[dP_N]\left\langle\hat{B}_N[A];\{a_r\};(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right|q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|-P_N\right\rangle\\
&\qquad\times\left\langle-P_N\right|q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle\\
&\approx\left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{2}}\int d^{26}p_0\exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)\right\}\\
&\quad\times\int\prod_{n=1}^{\infty}\left[\frac{d^{26}\bar{\psi}_n d^{26}\psi_n}{(2i)^{26}}\left(\frac{1}{\pi}\right)^{26}\left\{-\det\left(-\frac{(\theta G\theta)^{\mu\nu}}{16\pi^2nq_c^{\frac{n}{2}}\alpha'}\right)\right\}\exp\left(\frac{\bar{\psi}_{n\mu}\left(\theta G\theta\right)^{\mu\nu}\psi_{n\nu}}{16\pi^2nq_c^{\frac{n}{2}}\alpha'}\right)\right]
\end{aligned}
$$

$$
\begin{aligned}
&\times\delta^{26}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right)\prod_{1\le r<s\le M}e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\\
&\qquad\times\exp\left[-i\sum_{r=1}^{M}\left\{a_r\frac{dy^\mu(\sigma_r)}{d\sigma_r}A_\mu(k^{(r)})-k_\mu^{(r)}y^\mu(\sigma_r)\right\}\right]\\
&\times\delta^{26}\left(p_0-\sum_{r=M+1}^{M+N}k^{(r)}\right)\prod_{M+1\le r<s\le M+N}e^{-\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\\
&\qquad\times\exp\left[-i\sum_{r=M+1}^{M+N}\left\{a_r\frac{dy^\mu(\sigma_r)}{d\sigma_r}A_\mu(k^{(r)})-k_\mu^{(r)}y^\mu(\sigma_r)\right\}\right]\ ,
\end{aligned}
\tag{9.41}
$$

where $y^\mu(\sigma)$ is the curve given by Eq.(9.17).

The zero-slope limit of the amplitudes (9.38) can be obtained from the above by the operations $\prod_{r=1}^{M+N}i\frac{\partial}{\partial a_r}\Big|_{a_r=0}$, which are carried out without difficulty. The zero-slope limits turn out to be as follows :

$$
\begin{aligned}
&\int[dP_N]\left\langle B_N[A];(\sigma_{M+1},k^{(M+1)}),\cdots,(\sigma_{M+N},k^{(M+N)})\right|q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|-P_N\right\rangle\\
&\qquad\times\left\langle -P_N\right|q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)}\left|B_N[A];(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle\\
&\approx\left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{2}}\int d^{26}p_0\exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu}g^{\mu\nu}p_{0\nu}-\frac{4}{\alpha'}\right)\right\}\\
&\quad\times\int\prod_{n=1}^{\infty}\left[\frac{d^{26}\bar{\psi}_n d^{26}\psi_n}{(2i)^{26}}\left(\frac{1}{\pi}\right)^{26}\left\{-\det\left(-\frac{(\theta G\theta)^{\mu\nu}}{16\pi^2 n q_c^{\frac{n}{2}}\alpha'}\right)\right\}\exp\left(\frac{\bar{\psi}_{n\mu}\left(\theta G\theta\right)^{\mu\nu}\psi_{n\nu}}{16\pi^2 n q_c^{\frac{n}{2}}\alpha'}\right)\right]\\
&\qquad\times\delta^{26}\left(p_0+\sum_{r=1}^{M}k^{(r)}\right)\prod_{1\le r<s\le M}e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\\
&\qquad\qquad\times\prod_{r=1}^{M}\left\{\frac{dy^\mu(\sigma_r)}{d\sigma_r}A_\mu(k^{(r)})e^{ik_\nu^{(r)}y^\nu(\sigma_r)}\right\}\\
&\qquad\times\delta^{26}\left(p_0-\sum_{r=M+1}^{M+N}k^{(r)}\right)\prod_{M+1\le r<s\le M+N}e^{-\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)}\\
&\qquad\qquad\times\prod_{r=M+1}^{M+N}\left\{\frac{dy^\mu(\sigma_r)}{d\sigma_r}A_\mu(k^{(r)})e^{ik_\nu^{(r)}y^\nu(\sigma_r)}\right\}\ .
\end{aligned}
\tag{9.42}
$$

Factorized form of the $M+N$ gluon amplitude is obtained by integrating the above amplitude over the moduli (4.11), where we set $\sigma_{M+N}=2\pi$ in order to fix the $U(1)$ symmetry. This gives rise to the asymmetric term of the factorization (8.7) at the zero-slope limit. In order to avoid complexity of expressions we ignore this gauge fixing in the below and integrate σ_{M+N}

over $\sigma_{M+N-1} \leq \sigma_{M+N} \leq 2\pi$. The integrations over σ_r turn out to be written by using the Moyal products :

$$\begin{aligned}
&\int_{0\leq\sigma_1\leq\cdots\leq\sigma_M\leq 2\pi} \prod_{r=1}^{M} d\sigma_r \int_{0\leq\sigma_{M+1}\leq\cdots\leq\sigma_{M+N}\leq 2\pi} \prod_{s=M+1}^{M+N} d\sigma_s \int [dP_N] \\
&\quad \left\langle B_N[A]; (\sigma_{M+1}, k^{(M+1)}), \cdots, (\sigma_{M+N}, k^{(M+N)}) \right| q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)} \left|-P_N\right\rangle \\
&\qquad \times \left\langle -P_N \right| q_c^{\frac{1}{4}\left(L_0+\tilde{L}_0-2\right)} \left| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \\
&\approx \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{2}} \int d^{26}p_0 \exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu} g^{\mu\nu} p_{0\nu} - \frac{4}{\alpha'}\right)\right\} \\
&\quad \times \int \prod_{n=1}^{\infty} \left[\frac{d^{26}\bar{\psi}_n d^{26}\psi_n}{(2i)^{26}} \left(\frac{1}{\pi}\right)^{26} \left\{-\det\left(-\frac{(\theta G \theta)^{\mu\nu}}{16\pi^2 n q_c^{\frac{n}{2}} \alpha'}\right)\right\} \exp\left(\frac{\bar{\psi}_{n\mu} (\theta G \theta)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n q_c^{\frac{n}{2}} \alpha'}\right)\right] \\
&\quad \times \int \frac{d^{26}x}{(2\pi)^{26}} \left\{\frac{dy^\mu(\sigma_1)}{d\sigma_1} A_\mu(k^{(1)}) e^{ik_\nu^{(1)}(x^\nu + y^\nu(\sigma_1))}\right\} \\
&\qquad\qquad \star \cdots \star \left\{\frac{dy^\mu(\sigma_M)}{d\sigma_M} A_\mu(k^{(M)}) e^{ik_\nu^{(M)}(x^\nu + y^\nu(\sigma_M))}\right\} \star e^{ip_{0\mu}x^\mu} \\
&\quad \times \int \frac{d^{26}\tilde{x}}{(2\pi)^{26}}\, e^{-ip_{0\mu}\tilde{x}^\mu} \star \left\{\frac{dy^\mu(\sigma_{M+N})}{d\sigma_1} A_\mu(k^{(M+N)}) e^{ik_\nu^{(M+N)}(\tilde{x}^\nu + y^\nu(\sigma_{M+N}))}\right\} \\
&\qquad\qquad \star \cdots \star \left\{\frac{dy^\mu(\sigma_{M+1})}{d\sigma_{M+1}} A_\mu(k^{(M+1)}) e^{ik_\nu^{(M+1)}(\tilde{x}^\nu + y^\nu(\sigma_{M+1}))}\right\} \ .
\end{aligned} \tag{9.43}$$

Factorization of the generating function of the amplitudes (9.38) at the zero-slope limit can be obtained from the above by integrating out the gluon momenta k_r and then summing up with respect to M and N. It turns out to have the following form :

$$\begin{aligned}
&2\pi \int_0^{+\infty} d|\tau^{(c)}| \sum_{M=0}^{\infty} \sum_{N=0}^{\infty} i^M (-i)^N \int_{0\leq\sigma_1\leq\cdots\leq\sigma_M\leq 2\pi} \prod_{r=1}^{M} d\sigma_r \int_{0\leq\sigma_{M+1}\leq\cdots\leq\sigma_{M+N}\leq 2\pi} \prod_{s=M+1}^{M+N} d\sigma_s \\
&\quad \int \prod_{r=1}^{M} \frac{d^{26}k^{(r)}}{(2\pi)^{13}} \int \prod_{s=M+1}^{M+N} \frac{d^{26}k^{(s)}}{(2\pi)^{13}} \left\langle B_N[A]; (\sigma_{M+1}, -k^{(M+1)}), \cdots, (\sigma_{M+N}, -k^{(M+N)}) \right| \\
&\qquad\qquad\qquad \times q_c^{\frac{1}{2}\left(L_0+\tilde{L}_0-2\right)} \left| B_N[A]; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle \\
&\approx \int d^{26}p_0 \left[\frac{2\pi}{\alpha'} \int_0^{+\infty} ds^{(c)} \exp\left\{-\frac{\pi s^{(c)}}{2}\left(p_{0\mu} g^{\mu\nu} p_{0\nu} - \frac{4}{\alpha'}\right)\right\}\right] \\
&\quad \times \int \prod_{n=1}^{\infty} \left[\frac{d^{26}\bar{\psi}_n d^{26}\psi_n}{(2\pi i)^{26}} \left\{-\det\left(-\frac{\theta G \theta}{16\pi^2 n q_c^{\frac{n}{2}} \alpha'}\right)\right\} \exp\left(\frac{\bar{\psi}_{n\mu} (\theta G \theta)^{\mu\nu} \psi_{n\nu}}{16\pi^2 n q_c^{\frac{n}{2}} \alpha'}\right)\right] \\
&\quad \times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}} \int \frac{d^{26}x}{(2\pi)^{26}} \left[\mathcal{P}_\star \exp\left(i \int_0^{2\pi} d\sigma \frac{dy^\mu(\sigma)}{d\sigma} \mathcal{A}_\mu\left(x + y(\sigma)\right)\right)\right] \star e^{ip_{0\mu}x^\mu}
\end{aligned}$$

$$\times \left(\frac{-\det G_{\mu\nu}}{(2\alpha')^{26}}\right)^{\frac{1}{4}} \overline{\int \frac{d^{26}x}{(2\pi)^{26}} \left[\mathcal{P}_\star \exp\left(i\int_0^{2\pi} d\sigma \frac{dy^\mu(\sigma)}{d\sigma} \mathcal{A}_\mu\left(x+y(\sigma)\right)\right)\right] \star e^{ip_{0\mu}x^\mu}} \ . \tag{9.44}$$

10 Duality between Open and Closed Strings

So far, momentum eigenstates of closed-string play a crucial role in our study. Their overlaps with the boundary states provide open Wilson lines at the zero-slope limit. The eigenvalues, that is, loops in the momentum space become paths of open Wilson lines after suitable rescalings. In this section we wish to deliver the other side of the story. The following discussions include much speculation and therefore they are incomplete.

Let us first observe that the auxiliary boundary states of gluons are eigenstates of the momentum operator $\hat{P}^{(B)}_\mu(\sigma)$. The boundary state $\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\ \cdots,\ (\sigma_M,k^{(M)})\right\rangle$ has the eigenvalue equal to[10] $P^{(B)}_\mu(\sigma)=\sum_{r=1}^M\left(k^{(r)}_\mu - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right)\delta\left(\sigma-\sigma_r\right)$. Therefore this state is proportional to the eigenstate $\left|P^{(B)}_N\right\rangle$ of the corresponding eigenvalues. This can be seen by a comparison between oscillator representations (7.23) and (5.37) of the boundary state and the momentum eigenstate. In terms of the parametrization (5.26) the above eigenvalue corresponds to

$$\begin{aligned} \varrho_{n\mu} &= \sqrt{2}\sum_{r=1}^M \left(k^{(r)}_\mu - ina_rA_\mu(k^{(r)})\right)e^{in\sigma_r}\ , \\ \bar{\varrho}_{n\mu} &= \sqrt{2}\sum_{r=1}^M \left(k^{(r)} + ina_rA_\mu(k^{(r)})\right)e^{-in\sigma_r}\ , \quad p_{0\mu}=\sum_{r=1}^M k^{(r)}_\mu\ . \end{aligned} \tag{10.1}$$

By using these values of ϱ_n and $\overline{\varrho}_n$, the precise relation between the two states can be written as follows :

$$\begin{aligned} &\left|\hat{B}_N[A];\{a_r\};(\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right\rangle \\ &\quad = \prod_{r=1}^M \exp\left[\alpha' G^{\mu\nu}\sum_{n=1}^\infty \left(\frac{1}{n}k^{(r)}_\mu k^{(r)}_\nu + (a_r)^2 nA_\mu(k^{(r)})A_\nu(k^{(r)})\right)\right] \\ &\qquad \times e^{\frac{i}{2\pi}\sum_{r=1}^M\left(a_rA_\mu(k^{(r)})-\sigma_rk^{(r)}_\mu\right)\theta^{\mu\nu}\sum_{s=1}^M k^{(s)}_\nu} \prod_{r<s}^M e^{\frac{i}{2}k^{(r)}_\mu\theta^{\mu\nu}k^{(s)}_\nu\epsilon(\sigma_r-\sigma_s)} \end{aligned}$$

---

[10] The eigenvalue becomes complex but this causes no trouble in the subsequent arguments at least formally. Such subtlety does not occur in the tachyon case.

$$\times \left| P^{(B)}_\mu(\sigma) = \sum_{r=1}^{M} \left(k^{(r)}_\mu - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \delta(\sigma - \sigma_r) \right\rangle \ , \tag{10.2}$$

where we have used $\mathcal{C}_{P^{(B)}_N}$ in (5.38) as the normalization constant of the momentum eigenstate and represented it in terms of open-string tensors.

Let us comment on the multiplicative factors appearing in the above relation. As can be seen from Eq.(3.30), the second line on the RHS comes from the terms proportional to θ of disk Green's functions at the world-sheet boundary (the boundary circle). As regard the exponential in the first line on the RHS, we might think of it as being related to short distance singularities between the gluons on the boundary circle. In fact, by recasting the exponent into the following form,

$$\begin{aligned} &\alpha' G^{\mu\nu} \sum_{n=1}^{\infty} \left(\frac{1}{n} k^{(r)}_\mu k^{(r)}_\nu + (a_r)^2 n A_\mu(k^{(r)}) A_\nu(k^{(r)}) \right) \\ &\qquad = -\alpha' G^{\mu\nu} \lim_{s \to r} \left(k^{(r)}_\mu - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k^{(s)}_\nu - a_s A_\nu(k^{(s)}) \partial_{\sigma_s} \right) \ln \left| e^{i\sigma_r} - e^{i\sigma_s} \right| \ , \end{aligned} \tag{10.3}$$

we find a singularity similar to that appearing in the OPE between auxiliary gluon vertex operators.

In the same way, concerning the dual boundary states, we obtain the following relation as well:

$$\begin{aligned} &\left\langle \hat{B}_N; \{a_r\}; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right| \\ &\quad = \prod_{r=1}^{M} \left[\alpha' G^{\mu\nu} \sum_{n=1}^{\infty} \left(\frac{1}{n} k^{(r)}_\mu k^{(r)}_\nu + (a_r)^2 n A_\mu(k^{(r)}) A_\nu(k^{(r)}) \right) \right] \\ &\qquad \times e^{-\frac{i}{2} \sum_{r=1} \left(a_r A_\mu(k^{(r)}) - \sigma_r k^{(r)}_\mu \right) \theta^{\mu\nu} \sum_{s=1}^{M} k^{(s)}_\nu} \prod_{r<s}^{M} e^{-\frac{i}{2} k^{(r)}_\mu \theta^{\mu\nu} k^{(s)}_\nu \epsilon(\sigma_r - \sigma_s)} \\ &\qquad \times \left\langle P^{(B)}_\mu(\sigma) = -\sum_{r=1}^{M} \left(k^{(r)}_\mu - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \delta(\sigma - \sigma_r) \right| \ , \end{aligned} \tag{10.4}$$

where $\langle P^{(B)}_N |$ denotes the hermitian conjugate of $| P^{(B)}_N \rangle$.

Boundary states of open-string tachyons are obtained from auxiliary boundary states of gluons by letting their auxiliary parameters vanish. Hence these boundary states are also eigenstates of $\hat{P}^{(B)}_\mu(\sigma)$. The boundary state $\left| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle$ has the eigenvalue $P^{(B)}_\mu(\sigma) = \sum_{r=1}^{M} k^{(r)}_\mu \delta(\sigma - \sigma_r)$ and we can write the state as follows :

$$\left| B_N; (\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)}) \right\rangle$$

$$
\begin{aligned}
&= \prod_{r=1}^{M} \exp\left(\alpha' k_\mu^{(r)} G^{\mu\nu} k_\nu^{(r)} \sum_{n=1}^{\infty} \frac{1}{n}\right) \times e^{-\frac{i}{2\pi}\sum_{r=1}^{M}\sigma_r k_\mu^{(r)}\theta^{\mu\nu}\sum_{s=1}^{M} k_\nu^{(s)}} \prod_{r<s}^{M} e^{\frac{i}{2}k_\mu^{(r)}\theta^{\mu\nu}k_\nu^{(s)}\epsilon(\sigma_r-\sigma_s)} \\
&\quad \times \left| P_\mu^{(B)}(\sigma) = \sum_{r=1}^{M} k_\mu^{(r)}\delta(\sigma-\sigma_r) \right\rangle \ . \qquad (10.5)
\end{aligned}
$$

We would like to discuss physical meanings of the momentum eigenstates appearing in Eqs.(10.2) and (10.5). We henceforth concentrate on gluons. It is because analyses for the tachyon can be carried out in a parallel way to the gluon case and the formulae for gluons reduce to those of tachyons by setting the auxiliary parameters $a_r = 0$. Let us regard the momentum eigenstate $\left\langle P_\mu^{(B)}(\sigma) = -\sum_r \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right)\delta(\sigma-\sigma_r)\right|$ as a boundary state. The associated boundary action $S_b\left[X;P^{(B)}\right]$ can be obtained by following the prescription in [30]. It is given by

$$
e^{-S_b\left[X;P^{(B)}\right]} = \left\langle P_\mu^{(B)}(\sigma) = -\sum_{r=1}^{M}\left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right)\delta(\sigma-\sigma_r) \right| X_N \Big\rangle \ , \qquad (10.6)
$$

where $|X_N\rangle$ is the coordinate eigenstate in (5.31). Overlaps $\left\langle P_N^{(B)} \middle| X_N \right\rangle$ can be computed for arbitrary eigenvalues by using the formulae (B.7) in the oscillator representations of the eigenstates. The overlaps take the following forms :

$$
\begin{aligned}
\left\langle P_N^{(B)} \middle| X_N \right\rangle &= \frac{1}{(8\pi)^{\frac{p+1}{2}}}\mathcal{C}_X^{(N)}\mathcal{C}_{P_N^{(B)}}\frac{\det E_{\mu\nu}}{\det g_{\mu\nu}}\prod_{n=1}^{\infty} e^{-\frac{1}{4\pi n}\bar{\varrho}_{n\mu}\theta^{\mu\nu}\varrho_{n\nu}} \\
&\quad \times \exp\left[i\int_0^{2\pi} d\sigma\left(\frac{1}{2}B_{\mu\nu}X^\mu(\sigma)\partial_\sigma X^\nu(\sigma) - X^\mu(\sigma)P_\mu^{(B)}(\sigma)\right)\right] \ . \qquad (10.7)
\end{aligned}
$$

We apply the above formula to Eq.(10.6) with putting

$$
P_\mu^{(B)}(\sigma) = -\sum_{r=1}^{M}\left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right)\delta(\sigma-\sigma_r) \ .
$$

This enables us to find that the boundary action $S_b\left[X;P^{(B)}\right]$ becomes, modulo constant terms,

$$
\begin{aligned}
&S_b\left[X;P^{(B)}\right] \\
&= -i\int_0^{2\pi} d\sigma\left(\frac{1}{2}B_{\mu\nu}X^\mu(\sigma)\partial_\sigma X^\nu(\sigma) + X^\mu(\sigma)\sum_{r=1}^{M}\left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right)\delta(\sigma-\sigma_r)\right) \ . \\
&\qquad (10.8)
\end{aligned}
$$

The above boundary action appears naturally in the path-integral formalism of world-sheet theory of string. Let $\hat{\mathcal{G}}_A\left((\sigma_1,k^{(1)}),\cdots,(\sigma_M,k^{(M)})\right)$ be the correlation function of M gluons in the auxiliary forms with momenta $k_\mu^{(r)}$ and polarization vectors $A_\mu(k^{(r)})$ on the world-sheet disk. In the path-integral approach the correlation function can be expressed as

$$\begin{aligned}
&\hat{\mathcal{G}}_A\left((\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right) \\
&\quad = \int [dX^\mu] \left[e^{i\left(k_\mu^{(1)} - a_1 A_\mu(k^{(1)})\partial_{\sigma_1}\right) X^\mu(\sigma_1)} \cdots e^{i\left(k_\mu^{(M)} - a_M A_\mu(k^{(M)})\partial_{\sigma_M}\right) X^\mu(\sigma_M)} \right] e^{-S[X]} , \qquad (10.9)
\end{aligned}$$

where $S[X]$ is the world-sheet action (2.3). We can recast the RHS of this equation into

$$\hat{\mathcal{G}}_A\left((\sigma_1, k^{(1)}), \cdots, (\sigma_M, k^{(M)})\right) = \int [dX^\mu] e^{-S_{\text{eff}}[X]} , \qquad (10.10)$$

where $S_{\text{eff}}[X]$ is the sum of the bulk free action $S_0[X]$ and the boundary action $S_b[X]$: $S_{\text{eff}}[X] = S_0[X] + S_b[X]$ with

$$\begin{aligned}
S_0[X] &= \frac{1}{4\pi\alpha'} \int_\Sigma d\tau d\sigma \partial_a X^M(\sigma, \tau) \partial^a X^N(\sigma, \tau) g_{MN} , \qquad (10.11) \\
S_b[X] &= -i \int_{\partial\Sigma} d\sigma \left[\frac{1}{2} B_{\mu\nu} X^\mu(\sigma) \partial_\sigma X^\nu(\sigma) + X^\mu(\sigma) \sum_{r=1}^{M} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right) \delta(\sigma - \sigma_r) \right] .
\end{aligned}$$

Here we have used $\partial_\sigma X^\mu(\sigma) \cdot \delta(\sigma - \sigma_r) = -X^\mu(\sigma)\partial_\sigma \delta(\sigma - \sigma_r) = X^\mu(\sigma)\partial_{\sigma_r}\delta(\sigma - \sigma_r)$. We find that the above boundary action $S_b[X]$ is identical with $S_b\left[X; P^{(B)}\right]$ given in Eq.(10.8). Boundary conditions of X^μ can be read from variation of $S_{\text{eff}}[X]$. It turns out to be $P_\mu^{(B)}(\sigma, 0) = -\sum_{r=1}^{M} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right) \delta(\sigma - \sigma_r)$, as expected.

These observations on the boundary states and the momentum eigenstates seem to indicate a chance to interpret boundary states of open-string legs as momentum eigenstates of closed-string with eigenvalues being delta functions on the boundary circle. Our expectation is, in fact, beyond this. Let us recall that the momentum eigenstates are expressed in Eqs.(5.40) and (5.42) by using operators of the form

$$: \exp\left(i \int_{\partial\Sigma} d\sigma P_\mu(\sigma) \hat{X}^\mu(\sigma) \right) :$$

where $\hat{X}(\sigma)$ are the closed-string coordinate operators and : : denotes the standard normal-ordering of closed-string. If we forget about $\hat{X}(\sigma)$ being quantum operators, we could write down the exponential, without any hesitation, in the path-ordered form along the boundary circle, $\mathcal{P}\left[\exp\left(i \int_{\partial\Sigma} d\sigma P_\mu(\sigma) \hat{X}^\mu(\sigma)\right)\right]$. Since $\hat{X}(\sigma)$ are quantum operators the path-ordered exponential becomes vague without a prescription. Namely we need to regularize the path-ordered integral. Relevant regularization in place of the above normal-ordering will be discretization of the boundary circle. In such a regularization scheme the eigenvalues $P(\sigma)$ are regarded as sums of delta functions on the original boundary circle. This leads us to the conjecture : *Momentum eigenstates of closed-string have expansions by means of boundary states with open-string legs* .

Finally we would like to mention some related issues. As we carried out to a certain extent in this paper, our boundary states enable us to perform off-shell calculations. This suggests that these states can be used in a covariant formulation of a field theory of interacting open- and closed-strings. String field theory that has prediction power for the open-closed mixed systems has been required also in our understanding of unstable D-branes and tachyon condensations associated with them [44]. It seems probable that boundary states with open-string legs constructed here are generalized to vertex functions between open- and closed-strings in such a field theory. We plan to discuss these issues elsewhere.

ACKNOWLEDGEMENTS

T.N. would like to thank members of particle theory group of KEK for their hospitality during his stay in summer, 2002. The final part of this work was done there. K.M. would like to thank Y. Kitazawa, N. Ishibashi, E. Sezgin, T. Suyama, T. Masuda and P. Sundell for discussions and comments. The work of K.M. is supported in part by NSF Grant PHY-0070964.

A Open-String and Closed-String Tensors

Let $g_{\mu\nu}$ and $B_{\mu\nu}$ be a flat space-time metric and a constant two-form gauge field of closed-string. These are called closed-string tensors in the text. $E_{\mu\nu}$ is given by their combination as $E_{\mu\nu} \equiv g_{\mu\nu} + 2\pi\alpha' B_{\mu\nu}$. In [10], open-string metric $G_{\mu\nu}$ and non-commutativity parameter $\theta^{\mu\nu}$ are obtained respectively from symmetric and anti-symmetric parts of $\left(\frac{1}{E}\right)^{\mu\nu}$:

$$\left(\frac{1}{E}\right)^{\mu\nu} = G^{\mu\nu} + \frac{\theta^{\mu\nu}}{2\pi\alpha'} , \tag{A.1}$$

where $G^{\mu\nu} = (G^{-1})^{\mu\nu}$. Tensors $G_{\mu\nu}$ and $\theta^{\mu\nu}$ are called open-string tensors in the text. The above relation implies :

$$\begin{aligned} G^{\mu\nu} &= \frac{1}{2}\left(\frac{1}{E} + \frac{1}{E^T}\right)^{\mu\nu} = \left(\frac{1}{E} g \frac{1}{E^T}\right)^{\mu\nu} = \left(\frac{1}{E^T} g \frac{1}{E}\right)^{\mu\nu} , \\ \frac{\theta^{\mu\nu}}{2\pi\alpha'} &= \frac{1}{2}\left(\frac{1}{E} - \frac{1}{E^T}\right)^{\mu\nu} = -2\pi\alpha' \left(\frac{1}{E} B \frac{1}{E^T}\right)^{\mu\nu} = -2\pi\alpha' \left(\frac{1}{E^T} B \frac{1}{E}\right)^{\mu\nu} . \end{aligned} \tag{A.2}$$

It is also possible to express the closed-string tensors by means of the open-string ones :

$$\begin{aligned} g^{\mu\nu} &= \left(g^{-1}\right)^{\mu\nu} = G^{\mu\nu} - \frac{1}{(2\pi\alpha')^2} \left(\theta G \theta\right)^{\mu\nu} , \\ B_{\mu\nu} &= -\frac{1}{(2\pi\alpha')^2} \left(\frac{1}{G^{-1} + \frac{\theta}{2\pi\alpha'}} \theta \frac{1}{G^{-1} - \frac{\theta}{2\pi\alpha'}}\right)_{\mu\nu} . \end{aligned} \tag{A.3}$$

The tensor $N_{\mu\nu} \equiv \left(g \frac{1}{E} E^T\right)_{\mu\nu}$ is used frequently in the text. This tensor enjoys the following relations :

$$\left(\frac{1}{E^T} N \frac{1}{E^T}\right)^{\mu\nu} = G^{\mu\nu} , \tag{A.4}$$

$$\left(g^{-1} N g^{-1}\right)^{\mu\nu} = G^{\mu\nu} + \frac{1}{\left(2\pi\alpha'\right)^2} \left(\theta G \theta\right)^{\mu\nu} + \frac{\theta^{\mu\nu}}{\pi\alpha'} , \tag{A.5}$$

$$\left(N g^{-1} N^T\right)_{\mu\nu} = \left(N^T g^{-1} N\right)_{\mu\nu} = g_{\mu\nu} . \tag{A.6}$$

Eq.(A.5) implies

$$\left(g^{-1}\left(N - N^T\right) g^{-1}\right)^{\mu\nu} = \frac{2}{\pi\alpha'} \theta^{\mu\nu} . \tag{A.7}$$

Combination of Eq.(A.5) with Eq.(A.2) gives the following equalities :

$$\begin{aligned} g^{\mu\nu} + \left(g^{-1} N g^{-1}\right)^{\mu\nu} &= 2\left(\frac{1}{E}\right)^{\mu\nu} = 2G^{\mu\nu} + \frac{\theta^{\mu\nu}}{\pi\alpha'} , \\ g^{\mu\nu} + \left(g^{-1} N^T g^{-1}\right)^{\mu\nu} &= 2\left(\frac{1}{E^T}\right)^{\mu\nu} = 2G^{\mu\nu} - \frac{\theta^{\mu\nu}}{\pi\alpha'} . \end{aligned} \tag{A.8}$$

B Some Formulae of Creation and Annihilation Modes

We present formulae which become very useful for computations of string amplitudes in the text. As an illustration we derive F_A in Eq.(7.39) (and F in Eq.(3.47)).

B.1 The formulae

Formula B.1 *Let u_μ, v_μ, w_μ and y_μ be arbitrary complex $(p+1)$-vectors. Let $\Omega^{(1)}_{\mu\nu}$ and $\Omega^{(2)}_{\mu\nu}$ be any complex $(p+1)\times(p+1)$-matrices. The following equality holds :*

$$\begin{aligned}
&\exp\left[\frac{1}{n}\alpha^\mu_n\Omega^{(1)}_{\mu\nu}\tilde\alpha^\nu_n + u_\mu\alpha^\mu_n + v_\mu\tilde\alpha^\mu_n\right]\exp\left[\frac{1}{n}\alpha^\mu_{-n}\Omega^{(2)}_{\mu\nu}\tilde\alpha^\nu_{-n} + w_\mu\alpha^\mu_{-n} + y_\mu\tilde\alpha^\mu_{-n}\right]|\mathbf{0}\rangle \\
&\quad = \frac{\det g_{\mu\nu}}{\det\left(g-\Omega^{(2)T}g^{-1}\Omega^{(1)}\right)_{\mu\nu}} \\
&\qquad \times\exp\Bigg[n u_\mu\left(\frac{1}{g-\Omega^{(2)}g^{-1}\Omega^{(1)T}}\right)^{\mu\nu} w_\nu + n v_\mu\left(\frac{1}{g-\Omega^{(2)T}g^{-1}\Omega^{(1)}}\right)^{\mu\nu} y_\nu \\
&\qquad\qquad + n u_\mu\left(g^{-1}\Omega^{(2)}\frac{1}{g-\Omega^{(1)T}g^{-1}\Omega^{(2)}}\right)^{\mu\nu} v_\nu \\
&\qquad\qquad + n w_\mu\left(g^{-1}\Omega^{(1)}\frac{1}{g-\Omega^{(2)T}g^{-1}\Omega^{(1)}}\right)^{\mu\nu} y_\nu\Bigg] \\
&\qquad \times\exp\Bigg[\frac{1}{n}\alpha^\mu_{-n}\left(\Omega^{(2)}\frac{1}{g-\Omega^{(1)T}g^{-1}\Omega^{(2)}}g\right)_{\mu\nu}\tilde\alpha^\nu_{-n} \\
&\qquad\qquad + \left\{v_\lambda\left(g^{-1}\Omega^{(2)T}\right)^\lambda{}_\mu + w_\mu\right\}\left(\frac{1}{g-\Omega^{(1)}g^{-1}\Omega^{(2)T}}g\right)^\mu{}_\nu\alpha^\nu_{-n} \\
&\qquad\qquad + \left\{u_\lambda\left(g^{-1}\Omega^{(2)}\right)^\lambda{}_\mu + y_\mu\right\}\left(\frac{1}{g-\Omega^{(1)T}g^{-1}\Omega^{(2)}}g\right)^\mu{}_\nu\tilde\alpha^\nu_{-n}\Bigg]|\mathbf{0}\rangle\ . \qquad \text{(B.1)}
\end{aligned}$$

Use of coherent states becomes convenient to see the above equality. Let $\lambda^\pm_{n\mu}$ be complex $(p+1)$-vectors $(n=1,2,\ldots)$. Coherent state $|\lambda^+_n,\lambda^-_n)$ of the n-th level oscillators is defined as

$$\left|\lambda^+_n,\lambda^-_n\right) = \exp\left[\frac{1}{n}\left(\lambda^+_{n\mu}\alpha^\mu_{-n} + \lambda^-_{n\mu}\tilde\alpha^\mu_{-n}\right)\right]|0\rangle\ . \qquad \text{(B.2)}$$

$\lambda^\pm_{n\mu}$ becomes eigenvalues of the annihilation operators α^μ_n and $\tilde\alpha^\mu_n$ respectively :

$$\alpha^\mu_n\left|\lambda^+_n,\lambda^-_n\right) = g^{\mu\nu}\lambda^+_{n\nu}\left|\lambda^+_n,\lambda^-_n\right)\ ,\quad \tilde\alpha^\mu_n\left|\lambda^+_n,\lambda^-_n\right) = g^{\mu\nu}\lambda^-_{n\nu}\left|\lambda^+_n,\lambda^-_n\right)\ . \qquad \text{(B.3)}$$

Foe each n, the coherent states $|\lambda_n^+, \lambda_n^-)$ constitute a (over)complete basis of the Fock space built by α_{-n}^μ and $\tilde{\alpha}_{-n}^\mu$. The completeness relation reads

$$\begin{aligned} 1 \;=\; & \int \frac{1}{(\det g_{\mu\nu})^2} \prod_{\mu=0}^{p} \left(\frac{d\bar{\lambda}_{n\mu}^+ d\lambda_{n\mu}^+}{2\pi i n} \frac{d\bar{\lambda}_{n\mu}^- d\lambda_{n\mu}^-}{2\pi i n} \right) \\ & \times \left|\lambda_n^+, \lambda_n^-\right) \exp\left[-\frac{1}{n} \left(\bar{\lambda}_{n\mu}^+ g^{\mu\nu} \lambda_{n\nu}^+ + \bar{\lambda}_{n\mu}^- g^{\mu\nu} \lambda_{n\nu}^- \right) \right] \left(\lambda_n^+, \lambda_n^-\right| \ . \end{aligned} \tag{B.4}$$

Here $(\lambda_n^+, \lambda_n^-|$ denotes the hermitian conjugate of the state $|\lambda_n^+, \lambda_n^-)$. It takes the form of

$$\left(\lambda_n^+, \lambda_n^-\right| = \langle \mathbf{0} | \exp\left[\frac{1}{n} \left(\bar{\lambda}_{n\mu}^+ \alpha_n^\mu + \bar{\lambda}_{n\mu}^- \tilde{\alpha}_n^\mu \right) \right] \ , \tag{B.5}$$

where $\bar{\lambda}_{n\mu}^\pm$ are complex conjugate to $\lambda_{n\mu}^\pm$ and become eigenvalues of the creation operators α_{-n}^μ and $\tilde{\alpha}_{-n}^\mu$.

Formula B.1 can be shown by making use of the above partition of unity. We only describe an outline of the proof. We first insert the unity given in Eq.(B.4) between the two exponentials on the LHS of Eq.(B.1). This makes the LHS into Gaussian integrals with respect to $(\lambda_n^+, \bar{\lambda}_n^+)$ and $(\lambda_n^-, \bar{\lambda}_n^-)$. These Gaussian integrals are performed successively by using the relation

$$\int \prod_{\mu=0}^{p} \left(\frac{d\bar{\lambda}_\mu d\lambda_\mu}{2i} \right) \exp\left[-\left(\bar{\lambda}_\mu + \alpha_\mu \right) M^{\mu\nu} \left(\lambda_\nu + \beta_\nu \right) \right] = \frac{\pi^{p+1}}{\det M^{\mu\nu}} \ , \tag{B.6}$$

for $\forall \alpha_\mu, \beta_\mu \in \mathbb{C}^{p+1}$ and any $(p+1) \times (p+1)$ matrix $M^{\mu\nu}$. Then we obtain the RHS of Eq.(B.1).

The following formula is a corollary of Formula B.1:

Formula B.2

$$\begin{aligned} & \langle \mathbf{0} | \exp\left[\frac{1}{n} \alpha_n^\mu \Omega_{\mu\nu}^{(1)} \tilde{\alpha}_n^\nu + u_\mu \alpha_n^\mu + v_\mu \tilde{\alpha}_n^\mu \right] \exp\left[\frac{1}{n} \alpha_{-n}^\mu \Omega_{\mu\nu}^{(2)} \tilde{\alpha}_{-n}^\nu + w_\mu \alpha_{-n}^\mu + y_\mu \tilde{\alpha}_{-n}^\mu \right] | \mathbf{0} \rangle \\ & \quad = \frac{\det g_{\mu\nu}}{\det \left(g - \Omega^{(2)T} g^{-1} \Omega^{(1)} \right)_{\mu\nu}} \\ & \qquad \times \exp\left[n u_\mu \left(\frac{1}{g - \Omega^{(2)} g^{-1} \Omega^{(1)T}} \right)^{\mu\nu} w_\nu + n v_\mu \left(\frac{1}{g - \Omega^{(2)T} g^{-1} \Omega^{(1)}} \right)^{\mu\nu} y_\nu \right. \\ & \qquad \qquad + n u_\mu \left(g^{-1} \Omega^{(2)} \frac{1}{g - \Omega^{(1)T} g^{-1} \Omega^{(2)}} \right)^{\mu\nu} v_\nu \\ & \qquad \qquad \left. + n w_\mu \left(g^{-1} \Omega^{(1)} \frac{1}{g - \Omega^{(2)T} g^{-1} \Omega^{(1)}} \right)^{\mu\nu} y_\nu \right] \ . \end{aligned} \tag{B.7}$$

It is worth noting that a similar formula to Formula B.1 is used in open string field theory (see e.g. Eq.(B.2) of [45]). In particular, when we restrict $\Omega^{(1)}_{\mu\nu}$ and $\Omega^{(2)}_{\mu\nu}$ to symmetric matrices and change the variables from $\alpha^{\mu}_{\pm n}$ and $\tilde{\alpha}^{\mu}_{\pm n}$ to $\hat{a}^{(i)\mu}_{n}$ and $\hat{a}^{\dagger(i)\mu}_{n}$ ($i = \mathrm{I}, \mathrm{II}$) defined in Eq.(C.8), Formula B.1 reduces to the formula (B.2) of [45]. Formula B.1 can be regarded as a closed-string extension of it.

B.2 Applications

In this appendix, we derive Eqs.(3.47) and (7.39). This also serves as an illustration of usage of the formulae.

Contributions of the massive states of closed-string propagating between boundary states of tachyons are denoted by $F\left(q_c, \{\sigma_r\}, \{k^{(r)}\}\right)$ in Eq.(3.45). Those between boundary states of gluons are denoted by $F_A\left(q_c, \{\sigma_r\}, \{k^{(r)}\}; \{a_r\}\right)$ in Eq.(7.37). Their oscillator representations (3.46) and (7.38) imply that F_A reduces to F by setting $\forall a_r = 0$. We therefore focus on Eq.(7.39).

The representation (7.38) allows us to evaluate F_A by applying Formula B.2 with the following substitution :

$$\begin{aligned}
\Omega^{(1)}_{\mu\nu} &= -q_c^n N_{\mu\nu} \ , \qquad \Omega^{(2)}_{\mu\nu} = -N_{\mu\nu} \ , \\
u_\mu &= -\frac{\sqrt{2\alpha'}}{n} q_c^{\frac{n}{2}} \sum_{r=M+1}^{M+N} \left(k^{(r)}_{\mu} + i n a_r A_\nu(k^{(r)})\right) \left(\frac{1}{E^T} g\right)^{\nu}_{\ \mu} e^{-in\sigma_r} \ , \\
v_\mu &= -\frac{\sqrt{2\alpha'}}{n} q_c^{\frac{n}{2}} \sum_{r=M+1}^{M+N} \left(k^{(r)}_{\nu} - i n a_r A_\nu(k^{(r)})\right) \left(\frac{1}{E} g\right)^{\nu}_{\ \mu} e^{in\sigma_r} \ , \\
w_\mu &= \frac{\sqrt{2\alpha'}}{n} \sum_{r=1}^{M} \left(k^{(r)}_{\nu} - i n a_r A_\mu(k^{(r)})\right) \left(\frac{1}{E^T} g\right)^{\nu}_{\ \mu} e^{in\sigma_r} \ , \\
y_\mu &= \frac{\sqrt{2\alpha'}}{n} \sum_{r=1}^{M} \left(k^{(r)}_{\nu} + i n a_r A_\nu(k^{(r)})\right) \left(\frac{1}{E} g\right)^{\nu}_{\ \mu} e^{-in\sigma_r} \ .
\end{aligned} \tag{B.8}$$

In what follows, we will calculate each term in Eq.(B.7) with the above substitution. In this course the following relations will be used implicitly :

$$\left(g - \Omega^{(2)} g^{-1} \Omega^{(1)T}\right)_{\mu\nu} = \left(g - \Omega^{(2)T} g^{-1} \Omega^{(1)}\right)_{\mu\nu} = \left(1 - q_c^n\right) g_{\mu\nu} \ . \tag{B.9}$$

First, the determinants in the RHS of Eq.(B.7) become as follows :

$$\frac{\det g_{\mu\nu}}{\det\left(g - \Omega^{(2)T} g^{-1} \Omega^{(1)}\right)_{\mu\nu}} = \left(\frac{1}{1 - q_c^n}\right)^{p+1} \ . \tag{B.10}$$

As regards the exponential in the RHS of Eq.(B.7), the first two terms of the exponent are translated to

$$
\begin{aligned}
& nu_\mu \left(\frac{1}{g - \Omega^{(2)} g^{-1} \Omega^{(1)T}} \right)^{\mu\nu} w_\nu + nv_\mu \left(\frac{1}{g - \Omega^{(2)T} g^{-1} \Omega^{(1)}} \right)^{\mu\nu} y_\nu \\
& = -2\alpha' G^{\mu\nu} \sum_{r=1}^{M} \sum_{s=M+1}^{M+N} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\mu^{(s)} - a_s A_\mu(k^{(s)}) \partial_{\sigma_s} \right) \\
& \qquad \times \frac{q_c^{\frac{n}{2}}}{n\left(1 - q_c^n\right)} \left\{ e^{in(\sigma_r - \sigma_s)} + e^{-in(\sigma_r - \sigma_s)} \right\} \ . \qquad \text{(B.11)}
\end{aligned}
$$

We can recast the third term as follows :

$$
\begin{aligned}
& nu_\mu \left(g^{-1} \Omega^{(2)} \frac{1}{g - \Omega^{(1)T} g^{-1} \Omega^{(2)}} \right)^{\mu\nu} v_\nu \\
& = -2\alpha' G^{\mu\nu} \sum_{r,s=M+1}^{M+N} \left(k_\mu^{(r)} + ina_r A_\mu(k^{(r)}) \right) \left(k_\nu^{(s)} - ina_s A_\nu(k^{(s)}) \right) \frac{q_c^n e^{-in(\sigma_r - \sigma_s)}}{n\left(1 - q_c^n\right)} \\
& = -2\alpha' G^{\mu\nu} \sum_{M+1 \leq r < s \leq M+N} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\nu^{(s)} - a_s A_\nu(k^{(s)}) \partial_{\sigma_s} \right) \\
& \qquad \times \frac{q_c^n}{n\left(1 - q_c^n\right)} \left\{ e^{in(\sigma_r - \sigma_s)} + e^{-in(\sigma_r - \sigma_s)} \right\} \\
& \quad -2\alpha' G^{\mu\nu} \sum_{r=M+1}^{M+N} \left\{ k_\mu^{(r)} k_\nu^{(r)} + n^2 \left(a_r\right)^2 A_\mu(k^{(r)}) A_\nu(k^{(r)}) \right\} \frac{q_c^n}{n\left(1 - q_c^n\right)} \ . \qquad \text{(B.12)}
\end{aligned}
$$

The last term of the exponent turns out to be the same as (B.12) with shifting the indices r and s to $1 \leq r, s \leq M$.

Contribution of the n-th level oscillators is obtained by gathering all the above results. Contributions of the massive states are given by the infinite products taken over all the levels. These turn out to be written as follows :

$$
\begin{aligned}
& F_A\left(q_c, \{\sigma_r\} \left\{k^{(r)}\right\}; \{a_r\}\right) \\
& \quad = \left(\frac{1}{1 - q_c^n} \right)^{p+1} \\
& \quad \times \prod_{1 \leq r < s \leq M} \exp \Bigg[-2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\nu^{(s)} - a_s A\nu(k^{(s)}) \partial_{\sigma_s} \right) \\
& \qquad \times \sum_{n=1}^{\infty} \frac{\left(q_c e^{i(\sigma_r - \sigma_s)}\right)^n + \left(q_c e^{-i(\sigma_r - \sigma_s)}\right)^n - 2q_c^n}{n\left(1 - q_c^n\right)} \Bigg] \\
& \quad \times \prod_{M+1 \leq r < s \leq M+N} \exp \Bigg[-2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)}) \partial_{\sigma_r} \right) \left(k_\nu^{(s)} - a_s A_\nu(k^{(s)}) \partial_{\sigma_s} \right)
\end{aligned}
$$

$$
\begin{aligned}
&\qquad\qquad\qquad\qquad \times \sum_{n=1}^{\infty} \frac{\left(q_c e^{i(\sigma_r-\sigma_s)}\right)^n + \left(q_c e^{-i(\sigma_r-\sigma_s)}\right)^n - 2q_c^n}{n\left(1-q_c^n\right)} \Bigg] \\
&\times \prod_{r=1}^{M} \prod_{s=M+1}^{M+N} \exp \Bigg[-2\alpha' G^{\mu\nu} \left(k_\mu^{(r)} - a_r A_\mu(k^{(r)})\partial_{\sigma_r}\right) \left(k_\nu^{(s)} - a_s A_\nu(k^{(s)})\partial_{\sigma_s}\right) \\
&\qquad\qquad\qquad \times \sum_{n=1}^{\infty} \frac{\left(q_c^{\frac{1}{2}} e^{i(\sigma_r-\sigma_s)}\right)^n + \left(q_c^{\frac{1}{2}} e^{-i(\sigma_r-\sigma_s)}\right)^n - 2q_c^n}{n\left(1-q_c^n\right)} \Bigg] \\
&\times \exp \Bigg[-2\alpha' G^{\mu\nu} \left\{ \left(\sum_{r=1}^{M+N} k_\mu^{(r)} \right) \left(\sum_{s=1}^{M+N} k_\nu^{(s)} \right) + \sum_{r=1}^{M+N} \left(a_r\right)^2 A_\mu(k^{(r)}) A_\nu(k^{(r)}) \right\} \\
&\qquad \times \sum_{n=1}^{\infty} \frac{q_c^n}{n\left(1-q_c^n\right)} \Bigg] \ , \qquad\qquad\qquad\qquad\qquad\qquad \text{(B.13)}
\end{aligned}
$$

where we have used the following rearrangement :

$$
\begin{aligned}
\sum_{r=1}^{M+N} k_\mu^{(r)} G^{\mu\nu} k_\nu^{(r)} \quad = \quad & \left(\sum_{r=1}^{M+N} k_\mu^{(r)} \right) G^{\mu\nu} \left(\sum_{s=1}^{M+N} k_\nu^{(s)} \right) - 2 \sum_{1\le r<s\le M} k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)} \\
& -2 \sum_{r=1}^{M} \sum_{s=M+1}^{M+N} G^{\mu\nu} k_\mu^{(r)} k_\nu^{(s)} - 2 \sum_{M+1\le r<s\le M+N} k_\mu^{(r)} G^{\mu\nu} k_\nu^{(s)} \ . \quad \text{(B.14)}
\end{aligned}
$$

The infinite sums in Eq.(B.13) can be translated into infinite products by using the following relation :

$$
\sum_{n=1}^{\infty} \frac{x^n}{n\left(1-y^n\right)} = -\ln \prod_{m=0}^{\infty} \left(1 - xy^m\right) \ . \qquad \text{(B.15)}
$$

After these translations we obtain Eq.(7.39).

C Eigenstates

We provide oscillator realizations of eigenstates of the closed-string operators $\hat{X}^\mu(\sigma)$, $\hat{P}_\mu(\sigma)$ and $\hat{P}_\mu^{(B)}(\sigma)$. These are used in the text.

To start with, it is useful to recall coordinate and momentum eigenstates of a harmonic oscillator in quantum mechanics. Description of this system is made by an annihilation and a creation operators $\hat{a}$ and $\hat{a}^\dagger$ satisfying $[\hat{a}, \hat{a}^\dagger] = 1$. Let $\hat{q}$ and $\hat{p}$ be the coordinate and the momentum operators satisfying $[\hat{q}, \hat{p}] = 1$. The operators $(\hat{a}, \hat{a}^\dagger)$ are related with the canonical pair $(\hat{q}, \hat{p})$ as

$$
\hat{a} = \frac{1}{\sqrt{2}} \left(\frac{1}{\Gamma} \hat{q} + i\bar{\Gamma}\hat{p} \right) \ , \quad \hat{a}^\dagger = \frac{1}{\sqrt{2}} \left(\frac{1}{\bar{\Gamma}} \hat{q} - i\Gamma\hat{p} \right) \ ,
$$

$$\Longleftrightarrow \quad \hat{q} = \frac{1}{\sqrt{2}}\left(\Gamma\hat{a} + \bar{\Gamma}\hat{a}\right) \ , \quad \hat{p} = \frac{-i}{\sqrt{2}}\left(\frac{1}{\bar{\Gamma}}\hat{a} - \frac{1}{\Gamma}\hat{a}^\dagger\right) \ , \tag{C.1}$$

where $\Gamma \in \mathbb{C}$ is chosen so that $1/\omega|\Gamma|^2$ becomes the mass of harmonic oscillator. Here ω is the frequency. Let $|q\rangle$ and $|p\rangle$ be the eigenstates of $\hat{q}$ and $\hat{p}$. They are normalized by $\langle q'|q\rangle = \delta(q'-q)$ and $\langle p'|p\rangle = \delta(p'-p)$. It is possible to realize these states on the Fock vacuum $|0\rangle$ by using $\hat{a}^\dagger$. They are given by :

$$\begin{aligned} |q\rangle &= \left(\frac{1}{\pi|\Gamma|^2}\right)^{\frac{1}{4}} \exp\left[-\frac{\bar{\Gamma}}{2\Gamma}\hat{a}^\dagger\hat{a}^\dagger + \frac{\sqrt{2}}{\Gamma}q\hat{a}^\dagger - \frac{q^2}{2|\Gamma|^2}\right]|0\rangle \ , \\ |p\rangle &= \left(\frac{|\Gamma|^2}{\pi}\right)^{\frac{1}{4}} \exp\left[\frac{\bar{\Gamma}}{2\Gamma}\hat{a}^\dagger\hat{a}^\dagger + i\sqrt{2}\bar{\Gamma}p\hat{a}^\dagger - \frac{|\Gamma|^2}{2}p^2\right]|0\rangle \ . \end{aligned} \tag{C.2}$$

The above realizations of coordinate and momentum eigenstates are generalized to the case of string. We first expand the coordinate and the momentum operators of closed-string by suitable canonical pairs, typically denoted by $(\hat{\phi}^{(i)}_n, \hat{\pi}^{(i)}_n)$ $(i = \mathrm{I}, \mathrm{II};\ n = 1, 2, ..)$. For the each pair, we introduce creation and annihilation operators, typically $(\hat{a}^{(i)}_n, \hat{a}^{(i)\dagger}_n)$. We then realize eigenstates of the canonical operators $\hat{\phi}^{(i)}_n$ and $\hat{\pi}^{(i)}_n$ by using the creation operators $\hat{a}^{(i)\dagger}_n$. Eigenstates of the coordinate and the momentum operators of closed-string are given by their infinite products.

In order to obtain canonical pairs, we expand $\hat{X}^\mu(\sigma)$, $\hat{P}_\mu(\sigma)$ and $\hat{P}^{(B)}_\mu(\sigma)$ by a real basis of the periodic functions on a circle: $\{1, \cos n\sigma, \sin n\sigma\}_{n=1}^\infty$,

$$\begin{aligned} \hat{X}^\mu(\sigma) &= \hat{x}^\mu_0 + \sqrt{2}\sum_{n=1}^\infty \left(\hat{\phi}^{(\mathrm{I})\mu}_n \cos n\sigma + \hat{\phi}^{(\mathrm{II})\mu}_n \sin n\sigma\right) \ , \\ \hat{P}_\mu(\sigma) &= \frac{1}{2\pi}\left[\hat{p}_{0\mu} + \sqrt{2}\sum_{n=1}^\infty \left(\hat{\pi}^{(\mathrm{I})}_{n\mu}\cos n\sigma + \hat{\pi}^{(\mathrm{II})}_{n\mu}\sin n\sigma\right)\right] \ , \\ \hat{P}^{(B)}_\mu(\sigma) &= \frac{1}{2\pi}\left[\hat{p}_{0\mu} + \sqrt{2}\sum_{n=1}^\infty \left(\hat{\varpi}^{(\mathrm{I})}_{n\mu}\cos n\sigma + \hat{\varpi}^{(\mathrm{II})}_{n\mu}\sin n\sigma\right)\right] \ . \end{aligned} \tag{C.3}$$

The canonical commutation relations between $\hat{X}^\mu(\sigma)$ and $\hat{P}_\mu(\sigma)$ $(\hat{P}^{(B)}_\mu(\sigma))$ are converted into the following relations among the hermitian operators $\hat{\phi}^{(i)\mu}_n$, $\hat{\pi}^{(i)}_{n\mu}$ and $\hat{\varpi}^{(i)}_{n\mu}$ $(i = \mathrm{I}, \mathrm{II};\ n = 1, 2, \ldots)$:

$$[\hat{x}^\mu_0, \hat{p}_{0\nu}] = i\delta^\mu_\nu \ , \quad [\hat{\phi}^{(i)\mu}_m, \hat{\pi}^{(j)}_{n\nu}] = i\delta^{\mu\nu}\delta^{i,j}\delta_{m,n} \ , \quad [\hat{\phi}^{(i)\mu}_m, \hat{\varpi}^{(j)}_{n\nu}] = i\delta^\mu_\nu \ , \tag{C.4}$$

and the others are vanishing. These can be derived from the following relations as well:

$$\left\{\begin{aligned} \hat{\phi}^{(\mathrm{I})\mu}_n &= \frac{\sqrt{\alpha'}}{2n}i\left(\alpha^\mu_n + \tilde{\alpha}^\mu_n - \alpha^\mu_{-n} - \tilde{\alpha}^\mu_{-n}\right) \\ \hat{\phi}^{(\mathrm{II})\mu}_n &= \frac{\sqrt{\alpha'}}{2n}\left(\alpha^\mu_n - \tilde{\alpha}^\mu_n + \alpha^\mu_{-n} - \tilde{\alpha}^\mu_{-n}\right) \end{aligned}\right. \ , \quad \left\{\begin{aligned} \hat{\pi}^{(\mathrm{I})}_{n\mu} &= \frac{1}{2\sqrt{\alpha'}}g_{\mu\nu}\left(\alpha^\nu_n + \tilde{\alpha}^\nu_n + \alpha^\nu_{-n} + \tilde{\alpha}^\nu_{-n}\right) \\ \hat{\pi}^{(\mathrm{II})}_{n\mu} &= \frac{-i}{2\sqrt{\alpha'}}g_{\mu\nu}\left(\alpha^\nu_n - \tilde{\alpha}^\nu_n - \alpha^\nu_{-n} + \tilde{\alpha}^\nu_{-n}\right) \end{aligned}\right. \ ,$$

$$\begin{cases} \hat{\varpi}^{(\mathrm{I})}_{n\mu} = \dfrac{1}{2\sqrt{\alpha'}}\left(E_{\mu\nu}\alpha^{\nu}_{n} + E^{T}_{\mu\nu}\tilde{\alpha}^{\nu}_{n} + E_{\mu\nu}\alpha^{\nu}_{-n} + E^{T}_{\mu\nu}\tilde{\alpha}^{\nu}_{-n}\right) \\ \hat{\varpi}^{(\mathrm{II})}_{n\mu} = \dfrac{-i}{2\sqrt{\alpha'}}\left(E_{\mu\nu}\alpha^{\nu}_{n} - E^{T}_{\mu\nu}\tilde{\alpha}^{\nu}_{n} - E_{\mu\nu}\alpha^{\nu}_{-n} + E^{T}_{\mu\nu}\tilde{\alpha}^{\nu}_{-n}\right) \end{cases} . \tag{C.5}$$

Let us begin with the canonical pairs $\{\hat{\phi}^{(i)\mu}_{n}, \hat{\pi}^{(i)}_{n\mu}\}$. For these pairs, we introduce annihilation and creation modes $(\hat{a}^{(i)\mu}_{n}, \hat{a}^{\dagger(i)\mu}_{n})$ by

$$\hat{\phi}^{(i)\mu}_{n} = \sqrt{\frac{\alpha'}{2n}}\, i\left(\hat{a}^{(i)\mu}_{n} - \hat{a}^{\dagger(i)\mu}_{n}\right) \ , \quad \hat{\pi}^{(i)}_{n\mu} = \sqrt{\frac{n}{2\alpha'}}\, g_{\mu\nu}\left(\hat{a}^{(i)\nu}_{n} + \hat{a}^{\dagger(i)\nu}_{n}\right) \ . \tag{C.6}$$

They satisfy

$$[\hat{a}^{(i)\mu}_{m}, \hat{a}^{\dagger(j)\nu}_{n}] = g^{\mu\nu}\delta^{i,j}\delta_{m,n} \ . \tag{C.7}$$

It follows from Eqs.(C.5) that these modes are expressed as

$$\begin{aligned} \hat{a}^{(\mathrm{I})\mu}_{n} &= \frac{1}{\sqrt{2n}}\left(\alpha^{\mu}_{n} + \tilde{\alpha}^{\mu}_{n}\right) \ , \qquad \hat{a}^{\dagger(\mathrm{I})\mu} = \frac{1}{\sqrt{2n}}\left(\alpha^{\mu}_{-n} + \tilde{\alpha}^{\mu}_{-n}\right) \ , \\ \hat{a}^{(\mathrm{II})\mu}_{n} &= \frac{-i}{\sqrt{2n}}\left(\alpha^{\mu}_{n} - \tilde{\alpha}^{\mu}_{n}\right) \ , \qquad \hat{a}^{\dagger(\mathrm{II})\mu}_{n} = \frac{i}{\sqrt{2n}}\left(\alpha^{\mu}_{-n} - \tilde{\alpha}^{\mu}_{-n}\right) \ . \end{aligned} \tag{C.8}$$

Eqs.(C.6) take the same forms as Eqs.(C.1) with $\Gamma = i\sqrt{\frac{\alpha'}{n}}$. Therefore, Eqs.(C.2) enable us to write eigenstates $\left|\phi^{(\mathrm{I})}_{n}, \phi^{(\mathrm{II})}_{n}\right\rangle$ of $\hat{\phi}^{(\mathrm{I})\mu}_{n}$ and $\hat{\phi}^{(\mathrm{II})\mu}_{n}$ with eigenvalues $(\phi^{(\mathrm{I})\mu}_{n}, \phi^{(\mathrm{II})\mu}_{n}) \in \mathbb{R}^{p+1} \times \mathbb{R}^{p+1}$ as follows :

$$\begin{aligned} \left|\phi^{(\mathrm{I})}_{n}, \phi^{(\mathrm{II})}_{n}\right\rangle &= \left(\frac{n}{\pi\alpha'}\right)^{\frac{p+1}{2}} \sqrt{-\det g_{\mu\nu}} \\ &\times \prod_{i=\mathrm{I,II}} \exp\left[\frac{1}{2}\hat{a}^{\dagger(i)\mu}_{n} g_{\mu\nu}\hat{a}^{\dagger(i)\nu}_{n} - i\sqrt{\frac{2n}{\alpha'}}\phi^{(i)\mu}_{n} g_{\mu\nu}\hat{a}^{\dagger(i)\nu}_{n} - \frac{n}{2\alpha'}\phi^{(i)\mu}_{n} g_{\mu\nu}\phi^{(i)\nu}_{n}\right]|\mathbf{0}\rangle \ . \end{aligned} \tag{C.9}$$

In the same way, eigenstates $\left|\pi^{(\mathrm{I})}_{n}, \pi^{(\mathrm{II})}_{n}\right\rangle$ of $\hat{\pi}^{(\mathrm{I})\mu}_{n}$ and $\hat{\pi}^{(\mathrm{II})\mu}_{n}$ with eigenvalues $(\pi^{(\mathrm{I})\mu}_{n}, \pi^{(\mathrm{II})\mu}_{n}) \in \mathbb{R}^{p+1} \times \mathbb{R}^{p+1}$ become :

$$\begin{aligned} \left|\pi^{(\mathrm{I})}_{n}, \pi^{(\mathrm{II})}_{n}\right\rangle &= \left(\frac{\alpha'}{\pi n}\right)^{\frac{p+1}{2}} \frac{1}{\sqrt{-\det g_{\mu\nu}}} \\ &\times \prod_{i=\mathrm{I,II}} \exp\left[-\frac{1}{2}\hat{a}^{\dagger(i)\mu}_{n} g_{\mu\nu}\hat{a}^{\dagger(i)\nu}_{n} + \sqrt{\frac{\alpha'}{2n}}\pi^{(i)}_{n\mu}\hat{a}^{\dagger(i)\mu}_{n} - \frac{\alpha'}{2n}\pi^{(i)}_{n\mu} g^{\mu\nu}\pi^{(i)}_{n\nu}\right]|\mathbf{0}\rangle \ . \end{aligned} \tag{C.10}$$

By the construction, these states are normalized as follows :

$$\begin{aligned} \left\langle \phi'^{(\mathrm{I})}_{n}, \phi'^{(\mathrm{II}}_{n} \,\middle|\, \phi^{(\mathrm{I})}_{n}, \phi^{(\mathrm{II})}_{n}\right\rangle &= \prod_{\mu=0}^{p} \delta\left(\phi'^{(\mathrm{I})\mu}_{n} - \phi^{(\mathrm{I})\mu}_{n}\right)\delta\left(\phi'^{(\mathrm{II})\mu}_{n} - \phi^{(\mathrm{II})\mu}_{n}\right) \ , \\ \left\langle \pi'^{(\mathrm{I})}_{n}, \pi'^{(\mathrm{II}}_{n} \,\middle|\, \pi^{(\mathrm{I})}_{n}, \pi^{(\mathrm{II})}_{n}\right\rangle &= \prod_{\mu=0}^{p} \delta\left(\pi'^{(\mathrm{I})}_{n\mu} - \pi^{(\mathrm{I})}_{n\mu}\right)\delta\left(\pi'^{(\mathrm{II})}_{n\mu} - \pi^{(\mathrm{II})}_{n\mu}\right) \ . \end{aligned} \tag{C.11}$$

Eqs.(C.8) make it possible to write these eigenstates in the forms of Eqs.(5.14) and (5.23). In the text we use complex variables $(\chi_n^\mu, \bar{\chi}_n^\mu) = (\phi_n^{(\mathrm{I})\mu} + i\phi_n^{(\mathrm{II})\mu}, \phi_n^{(\mathrm{I})\mu} - i\phi_n^{(\mathrm{II})\mu})$ and write the coordinate eigenstates as $|\chi_n, \bar{\chi}_n\rangle = \left|\phi_n^{(\mathrm{I})}, \phi_n^{(\mathrm{II})}\right\rangle$. As for the momentum eigenstates, we use complex variables $(\psi_{n\mu}, \bar{\psi}_{n\mu}) = (\pi_{n\mu}^{(\mathrm{I})} + i\pi_{n\mu}^{(\mathrm{II})}, \pi_{n\mu}^{(\mathrm{I})} - i\pi_{n\mu}^{(\mathrm{II})})$ and write them as $\left|\psi_n, \bar{\psi}_n\right\rangle = \left|\pi_n^{(\mathrm{I})}, \pi_n^{(\mathrm{II})}\right\rangle$.

Next we consider eigenstates of $\hat{P}_\mu^{(B)}(\sigma)$. Taking account of Eqs.(C.5), let us write the expansion modes $\hat{\varpi}_{n\mu}^{(i)}$ $(i = \mathrm{I}, \mathrm{II};\ n = 1, 2, \ldots)$ as

$$\hat{\varpi}_{n\mu}^{(i)} = \sqrt{\frac{n}{2\alpha'}} G_{\mu\nu} \left(\hat{\eta}_n^{(i)\mu} + \hat{\eta}_n^{\dagger(i)\mu}\right) \ , \tag{C.12}$$

where $\hat{\eta}_n^{(i)\mu}$ and $\hat{\eta}_n^{\dagger(i)\mu}$ denote

$$\begin{aligned}
\hat{\eta}_n^{(\mathrm{I})\mu} &= \frac{1}{\sqrt{2n}} \left\{ \left(\frac{1}{E^T} g\right)^\mu_{\ \nu} \alpha_n^\nu + \left(\frac{1}{E} g\right)^\mu_{\ \nu} \tilde{\alpha}_n^\nu \right\} \ , \\
\hat{\eta}_n^{\dagger(\mathrm{I})\mu} &= \frac{1}{\sqrt{2n}} \left\{ \left(\frac{1}{E^T} g\right)^\mu_{\ \nu} \alpha_{-n}^\nu + \left(\frac{1}{E} g\right)^\mu_{\ \nu} \tilde{\alpha}_{-n}^\nu \right\} \ , \\
\hat{\eta}_n^{(\mathrm{II})\mu} &= \frac{-i}{\sqrt{2n}} \left\{ \left(\frac{1}{E^T} g\right)^\mu_{\ \nu} \alpha_n^\nu - \left(\frac{1}{E} g\right)^\mu_{\ \nu} \tilde{\alpha}_n^\nu \right\} \ , \\
\hat{\eta}_n^{\dagger(\mathrm{II})\mu} &= \frac{i}{\sqrt{2n}} \left\{ \left(\frac{1}{E^T} g\right)^\mu_{\ \nu} \alpha_{-n}^\nu - \left(\frac{1}{E} g\right)^\mu_{\ \nu} \tilde{\alpha}_{-n}^\nu \right\} \ .
\end{aligned} \tag{C.13}$$

They turn out to satisfy

$$[\hat{\eta}_m^{(i)\mu}, \hat{\eta}_n^{\dagger(j)\nu}] = G^{\mu\nu} \delta^{i,j} \delta_{m,n} \ . \tag{C.14}$$

Comparing Eqs.(C.12) and (C.14) with Eqs.(C.6) and (C.7), we can find that eigenstates $\left|\varpi_n^{(\mathrm{I})}, \varpi_n^{(\mathrm{II})}\right\rangle$ of $\hat{\varpi}_{n\mu}^{(\mathrm{I})}$ and $\hat{\varpi}_{n\mu}^{(\mathrm{II})}$ are obtained from $\left|\pi_n^{(\mathrm{I})}, \pi_n^{(\mathrm{II})}\right\rangle$ by the following replacements :

$$g_{\mu\nu} \to G_{\mu\nu} \ ; \quad \left(\hat{a}_n^{(i)\mu}, \hat{a}_n^{\dagger(i)\mu}\right) \to \left(\hat{\eta}_n^{(i)\mu}, \hat{\eta}_n^{\dagger(i)\mu}\right) \ . \tag{C.15}$$

Thus we have :

$$\begin{aligned}
\left|\varpi_n^{(\mathrm{I})}, \varpi_n^{(\mathrm{II})}\right\rangle &= \left(\frac{\alpha'}{\pi n}\right)^{\frac{p+1}{2}} \frac{1}{\sqrt{-\det G_{\mu\nu}}} \\
&\quad \times \prod_{i=\mathrm{I},\mathrm{II}} \exp\left[-\frac{1}{2} \hat{\eta}^{\dagger(i)\mu} G_{\mu\nu} \hat{\eta}_n^{\dagger(i)\nu} + \sqrt{\frac{2\alpha'}{n}} \varpi_{n\mu}^{(i)} \hat{\eta}_n^{\dagger(i)\mu} - \frac{\alpha'}{2n} \varpi_{n\mu}^{(i)} G^{\mu\nu} \varpi_{n\nu}^{(i)} \right] |\mathbf{0}\rangle \ .
\end{aligned} \tag{C.16}$$

Substitutions of Eqs.(C.13) into the above realizations make the states in the forms (5.29). We use complex variables $(\varrho_{n\mu}, \bar{\varrho}_{n\mu}) = (\varpi_{n\mu}^{(\mathrm{I})} + i\varpi_{n\mu}^{(\mathrm{II})}, \bar{\varpi}_{n\mu}^{(\mathrm{I})} - i\varpi_{n\mu}^{(\mathrm{II})})$ in the text and write $|\varrho_n, \bar{\varrho}_n\rangle = \left|\varpi_n^{(\mathrm{I})}, \varpi_n^{(\mathrm{II})}\right\rangle$.

References

[1] J. Polchinski, "*Dirichlet-Branes and Ramond-Ramond Charges*", Phys. Rev. Lett. **75** (1995) 4724, [hep-th/9510017].

[2] J. Maldacena, "*The Large N Limit of Superconformal Field Theories and Supergravity*", Adv. Theor. Math. Phys. **2** (1998) 231, [hep-th/9711200].

[3] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, "*Gauge Theory Correlators from Non-Critical String Theory*", Phys. Lett. **B428** (1998) 105, [hep-th/9802109].

[4] E. Witten, "*Anti De Sitter Space and Holography*", Adv. Theor. Math. Phys. **2** (1998) 253, [hep-th/9802150].

[5] A. Jevicki, Y. Kazama and T. Yoneya, "*Quantum Metamorphosis of Conformal Transformation in D3-Brane Yang-Mills Theory*", Phys. Rev. Lett. **81** (1998) 5072, [hep-th/9808039].

[6] A. Connes, M.R. Douglas and A. Schwarz, "*Noncommutative Geometry and Matrix Theory: Compactification on Tori*", J. High Energy Phys. **9802** (1998) 003, [hep-th/9711162].

[7] M.R. Douglas and C. Hull, "*D-branes and the Noncommutative Torus*", J. High Energy Phys. **9802** (1998) 008, [hep-th/9711165].

[8] C.-S. Chu and P.-M. Ho, "*Noncommutative Open String and D-brane*", Nucl. Phys. **B550** (1999) 151, [hep-th/9812219];
"*Constrained Quantization of Open String in Background B Field and Noncommutative D-brane*", Nucl. Phys. **B568** (2000) 447, [hep-th/9906192].

[9] F. Ardalan, H. Arfaei and M.M. Sheikh-Jabbari, "*Noncommutative Geometry from Strings and Branes*", J. High Energy Phys. **9902** (1999) 016, [hep-th/9810072];

"*Dirac Quantization of Open Strings and Noncommutativity in Branes*", Nucl. Phys. **B576** (2000) 578, [hep-th/9906161].

[10] N. Seiberg and E. Witten, "*String Theory and Noncommutative Geometry*", J. High Energy Phys. **9909** (1999) 032, [hep-th/9908142].

[11] A. Gonzalez-Arroyo and M. Okawa, "*Twisted-Eguchi-Kawai Model: A Reduced Model for Large-N Lattice Gauge Theory*", Phys. Rev. **D27** (1983) 2397.

S. Minwalla, M.V. Raamsdonk and N. Seiberg, "*Noncommutative Perturbative Dynamics*", J. High Energy Phys. **0002** (2000) 020, [hep-th/9912072].

M.V. Raamsdonk and N. Seiberg, "*Comments on Noncommutative Perturbative Dynamics*", J. High Energy Phys. **0003** (2000) 035, [hep-th/0002186].

[12] O. Andreev and H. Dorn, "*Diagrams of Noncommutative* Φ^3 *Theory from String Theory*", Nucl. Phys. **B583** (2000) 145, [hep-th/0003113].

[13] Y. Kiem and S. Lee, "*UV/IR Mixing in Noncommutative Field Theory via Open String Loops*", Nucl. Phys. **B586** (2000) 303, [hep-th/0003145]

[14] A. Bilal, C.-S. Chu and R. Russo, "*String Theory and Noncommutative Field Theories at One Loop*", Nucl. Phys. **B582** (2000) 65, [hep-th/0003180].

[15] J. Gomis, M. Kleban, T. Mehen, M. Rangamani and S. Shenker, "*Noncommutative Gauge Dynamics From The String Worldsheet*", J. High Energy Phys. **0008** (2000) 011, [hep-th/0003215].

[16] H. Liu and J. Michelson, "*Stretched Strings in Noncommutative Field Theory*", Phys. Rev. **D62** (2000) 066003, [hep-th/0004013].

[17] C.-S. Chu, R. Russo and S. Sciuto, "*Multiloop String Amplitudes with B-Field and Noncommutative QFT*", Nucl. Phys. **B585** (2000) 193, [hep-th/0004183].

Y. Kiem, S.-S. Kim, S.-J. Rey and H.-T. Sato, "*Anatomy of Two-Loop Effective Action in Noncommutative Field Theories*", Nucl. Phys. **B641** (2002) 256, [hep-th/0110066].

Y. Kiem, S. Lee, S.-J. Rey and H.-T. Sato, "*Interacting Open Wilson Lines from Noncommutative Field Theories*", Phys. Rev. **D65** (2002) 046003, [hep-th/0110215].

[18] A. Hashimoto and N. Itzhaki, "*Non-Commutative Yang-Mills and the AdS/CFT Correspondence*", Phys. Lett. **B465** (1999) 142, [hep-th/9907166].

[19] J.M. Maldacena and J.G. Russo, "*Large N Limit of Non-Commutative Gauge Theories*", J. High Energy Phys. **9909** (1999) 025, [hep-th/9908134].

[20] S. Hyun, Y. Kiem, S. Lee and C.-Y. Lee, "*Closed Strings Interacting with Noncommutative D-branes*", Nucl. Phys. **B569** (2000) 262, [hep-th/9909059].

[21] M.R. Garousi, "*Non-commutative world-volume interactions on D-brane and Dirac-Born-Infeld action*", Nucl. Phys. **B579** (2000) 209, [hep-th/9909214].

[22] T. Mehen and M.B. Wise, "*Generalized $\star$-Products, Wilson Lines and the Solution of the Seiberg-Witten Equations*", J. High Energy Phys. **0012** (2000) 008, [hep-th/0010204].

[23] Y. Okawa and H. Ooguri, "*How Noncommutative Gauge Theories Couple to Gravity*", Nucl. Phys. **B599** (2001) 55, [hep-th/0012218].

[24] N. Ishibashi, S. Iso, H. Kawai and Y. Kitazawa, "*Wilson Loops in Noncommutative Yang Mills*", Nucl. Phys. **B573** (2000) 573, [hep-th/9910004].

[25] D.J. Gross, A. Hashimoto and N. Itzhaki, "*Observables of Non-Commutative Gauge Theories*", Adv. Theor. Math. Phys. **4** (2000) 893, [hep-th/0008075].

[26] S. Das and S.-J. Rey, "*Open Wilson Lines in Noncommutative Gauge Theory and Tomography of Holographic Dual Supergravity*", Nucl. Phys. **B590** (2000) 453, [hep-th/0008042].

[27] A. Dhar and Y. Kitazawa, "*Loop Equation and Wilson Line Correlators in Noncommutative Gauge Theories*", Nucl. Phys. **B613** (2001) 105, [hep-th/0104021];

"*Non-commutative Gauge Theory, Open Wilson Lines and Closed Strings*", J. High Energy Phys. **0108** (2001) 044, [hep-th/0106217].

[28] C.G. Callan, C. Lovelace, C.R. Nappi and S.A. Yost, "*String Loop Corrections to Beta Functions*", Nucl. Phys. **B288** (1987) 525.

[29] C.G. Callan, C. Lovelace, C.R. Nappi and S.A. Yost, "*Adding Holes and Crosscaps to the Superstring*", Nucl. Phys. **B293** (1987) 83.

[30] C.G. Callan, C. Lovelace, C.R. Nappi and S.A. Yost, "*Loop Corrections to Superstring Equations of Motion*", Nucl. Phys. **B308** (1988) 221.

[31] N. Ishibashi, "*The Boundary and Crosscap States in Conformal Field Theories*", Mod. Phys. Lett. **4** (1989) 251.

[32] Y. Kiem, S.-J. Rey, H.-T. Sato and J.-T. Yee, "*Anatomy of One-Loop Effective Action in Noncommutative Scalar Field Theories*", Eur. Phys. J. **C22** (2002) 757, [hep-th/0107106].

[33] B. Zwiebach, "*Closed String Field Theory: Quantum Action and the BV Master Equation*", Nucl. Phys. **B390** (1993) 33, [hep-th/9206084].

[34] T. Nakatsu, "*Classical Open-String Field Theory: A_∞-Algebra, Renormalization Group and Boundary States*", Nucl. Phys. **B642** (2002) 13, [hep-th/0105272].

[35] M.B. Green, J.H. Schwarz and E. Witten, "*Superstring theory*", Vol. 2, (Cambridge University Press, 1987).

[36] F. Sugino, "*Witten's Open String Field Theory in Constant B-Field Background*", J. High Energy Phys. **0003** (2000) 017, [hep-th/9912254].

T. Kawano and T. Takahashi, "*Open String Field Theory on Noncommutative Space*", Prog. Theor. Phys. **104** (2000) 459, [hep-th/9912274].

[37] S.S. Gubser and S.L. Sondhi, "*Phase Structure of Non-commutative Scalar Field Theories*", Nucl. Phys. **B605** (2001) 395, [hep-th/0006119].

I. Chepelev and R. Roiban, "*Convergence Theorem for Non-commutative Feynman Graphs and Renormalization*", J. High Energy Phys. **0103** (2001) 001, [hep-th/0008090].

G.-H. Chen and Y.-S. Wu, "*On Critical Phenomena in a Noncommutative Space*", [hep-th/0103020].

[38] Y. Kiem, S.-J. Rey, H.-T. Sato and J.-T. Yee, "*Open Wilson Lines and Generalized Star Product in Noncommutative Scalar Field Theories*", Phys. Rev. **D65** (2002) 026002, [hep-th/0106121].

[39] C. Itzykson and J.-B. Zuber, "*Quantum Field Theory*", (McGraw-Hill, 1980).

[40] D.J. Gross and A. Jevicki, "*Operator Formulation of Interacting String Field Theory (I)*", Nucl. Phys. **B283** (1987) 1.

[41] H. Itoyama and P. Moxhay, "*Multiparticle Superstring Tree Amplitudes*", Nucl. Phys. **B293** (1987) 685.

[42] A. Armoni and E. Lopez, "*UV/IR Mixing via Closed Strings and Tachyonic Instabilities*", Nucl. Phys. **B632** (2002) 240, [hep-th/0110113].

[43] Y. Kiem, Y. Kim, C. Ryou and H.-T. Sato, "*One-loop Noncommutative* $U(1)$ *Gauge Theory from Bosonic Worldline Approach*", Nucl. Phys. **B630** (2002) 55, [hep-th/0112176].

[44] A. Sen, "*Descent Relations among Bosonic D-branes*", Int. J. Mod. Phys. **A14** (1999) 4061, [hep-th/9902105].

[45] I. Kishimoto, "*Some Properties of String Field Algebra*", J. High Energy Phys. **0112** (2001) 007, [hep-th/0110124].